

**A new species of *Cratiria* (Caliciaceae, Ascomycota)  
from Ascension Island, South Atlantic Ocean**

**John A. Elix**

Research School of Chemistry, Building 137,  
Australian National University, Canberra, ACT 2601, Australia  
**e-mail:** John.Elix@anu.edu.au

**Helmut Mayrhofer**

Institute of Biology, Division of Plant Sciences, NAWI Graz,  
University of Graz, Holteigasse 6, 8010 Graz, Austria  
**e-mail:** helmut.mayrhofer@uni-graz.at

**Abstract**

*Cratiria jamesiana* Elix & H. Mayrhofer, a saxicolous species with *Physconia*- then *Buellia*-type ascospores and bacilliform conidia, and containing thuringione and arthothelin, is described as new to science.

**Introduction**

This paper is a continuation of our investigations into *Buellia*-like lichens in the Southern Hemisphere (Elix 2018; Elix & McCarthy 2018; Elix *et al.* 2018; Elix 2019a, b and references therein). The genus *Cratiria* Marbach includes species that are characterized by relatively large, submuriform or 1-septate ascospores, 15–28 × 7–13 µm, with apical wall-thickenings, short, bacilliform conidia 4–6 µm long, a hymenium that can be interspersed with oil droplets or not and an excipulum containing lichen substances (Marbach 2000; Elix 2014). In this paper we describe a new saxicolous species of *Cratiria* from Ascension Island in the South Atlantic Ocean. Methods are as described in previous papers cited above.

*Cratiria jamesiana* Elix & H. Mayrhofer, sp. nov.  
MycoBank number: **MB 832570**

Figs 1, 2

Similar to *Cratiria chloraceus* Marbach, but differs in being saxicolous and in having a K-excipulum and hypothecium.

*Type:* Ascension Island, Green Mountain, Monkey Rock, [07.9500°S, 14.3500°W], 460 m alt., on volcanic rock, *P.W. James s.n.*, 3.xi.1976 (BM – holotype).

*Thallus* crustose, continuous, to 55 mm wide and up to 1 mm thick; upper surface pale grey to pale yellow-grey, verruculose, matt; warts 0.1–1 mm wide; prothallus black, marginal when abutting other lichens; medulla white, lacking calcium oxalate (H<sub>2</sub>SO<sub>4</sub>–), I–; photobiont cells 8–14 µm diam. *Apothecia* 0.4–1.5 mm wide, lecideine, scattered, sessile; disc black, pale grey-pruinose, concave to ± plane; proper exciple thick, concolorous with the disc, entire, persistent, cupuliform, in section 75–100 µm thick; outer zone opaque brown-black with crystals, K–, N–; inner zone brown. *Hypothecium* 190–240 µm thick, dark brown to brown-black, K–. *Epithymenium* 10–15 µm thick, pale brown to grey-brown, with crystals soluble in K, K–, N–. *Hymenium* 90–110 µm thick, colourless, interspersed with oil droplets; subhymenium 25–35 µm thick, pale brown; paraphyses 1.7–2.0 µm wide, simple to branched, capitate, with apices dark brown, 3–3.5 µm wide; asci approximating the *Bacidia*-type, with 8 or fewer spores. *Ascospores* at first of the *Physconia*-type, then of the *Buellia*-type, 1-septate, olive-brown to brown, ellipsoid, 15–[16.6]–20 × 8–[8.9]–10 µm, rarely constricted at the septum, with apical wall-thickenings; outer spore-wall rugulate. *Pycnidia* immersed; conidia bacilliform, straight, 4–5 × 0.7–1 µm.

*Chemistry:* Thallus K–, C–, P–, UV+ orange; containing thuringione (major), arthothelin (major), 3-*O*-methylthiophanic acid (minor).

**OFF-PRINT**

A new species of *Cratiria* (Caliciaceae, Ascomycota)  
from Ascension Island, South Atlantic Ocean

John A. Elix & Helmut Mayrhofer

*Australasian Lichenology* **86** (January 2020), 87–89

**Etymology:** the species is named after the collector of the type specimen, the late British lichenologist Peter James.

### Remarks

This new species is characterized by the verrucose-areolate, pale grey to pale yellow-grey crustose thallus, with a non-amyloid medulla lacking calcium oxalate, the sessile, lecideine apothecia with pruinose discs, an inspersed hymenium, *Physconia*- then *Buellia*-type ascospores,  $15\text{--}20 \times 8\text{--}10\text{ }\mu\text{m}$ , the bacilliform conidia,  $4\text{--}5\text{ }\mu\text{m}$  long, and the presence of thuringione and arthothelin. Its chemistry and anatomy closely resemble those of *C. chloraceus*, a corticolous species known from Australia, Papua New Guinea and New Caledonia (Elix *et al.* 2017). However, the thallus of *C. chloraceus* differs in being corticolous and significantly thinner, and the hypothecium reacts K+ intense crimson-purple. *Cratiria jamesiana* could be confused with *Buellia halonia* (Ach.) Tuck., which has been reported from Ascension Island (Aptroot 2008). Both species have similar-sized *Physconia*- then *Buellia*-type ascospores, and both contain xanthonenes. However, *B. halonia* has a non-inspersed hymenium, an epihymenium that is often aeruginose in part, and ascospores lacking apical wall-thickenings. In addition, it contains isoarthothelin (major), roccellic acid (major) and minor quantities of atranorin. *Cratiria jamesiana* is the first saxicolous species of *Cratiria* known to contain xanthonenes.

At present, the new species is known only from the type collection.

### Acknowledgement

H.M. acknowledges financial support from the Austrian Science Fund (FWF-projects P8500-BIO, P10514-BIO and P25237-B16).

### References

- Aptroot, A (2008): Lichens of St Helena and Ascension Island. *Botanical Journal of the Linnean Society* **158**, 147–171.
- Elix, JA (2014): New species and new records of the lichen genus *Cratiria* (Physciaceae, Ascomycota) in Australia. *Telopea* **16**, 141–148.
- Elix, JA (2018): A key to the buellioid lichens (Ascomycota, Caliciaceae) in New Zealand. *Australasian Lichenology* **83**, 26–35.
- Elix, JA (2019a): A new species and new records of buellioid lichens (Caliciaceae, Ascomycota) from the Kerguelen Islands. *Australasian Lichenology* **84**, 16–25.
- Elix, JA (2019b): Four new species and new records of buellioid lichens (Caliciaceae, Ascomycota) from Antarctica. *Australasian Lichenology* **84**, 33–43.
- Elix, JA; Kantvilas, G; McCarthy, PM (2017): Thirteen new species and a key to buellioid lichens (Caliciaceae, Ascomycota) in Australia. *Australasian Lichenology* **81**, 26–67.
- Elix, JA; Mayrhofer, H; Rodriguez, JM (2018): Two new species, a new combination and four new records of saxicolous buellioid lichens (Ascomycota, Caliciaceae) from southern South America. *Australasian Lichenology* **83**, 3–13.
- Elix, JA; McCarthy, PM (2018): Three new species and four new records of buellioid lichens (Caliciaceae, Ascomycota) from south-eastern Australia. *Herzogia* **31**, 444–452.
- Marbach, B (2000): Corticole und lignicole Arten der Flechtengattung *Buellia* sensu lato in den Subtropen und Tropen. *Bibliotheca Lichenologica* **74**, 1–384.



Figure 1. *Cratiria jamesiana* (holotype in BM). Bar = 2 mm.




Figure 2. Ascospore ontogeny of *C. jamesiana*. Bar = 10  $\mu\text{m}$ .