

Continuous Differentiability of the Value Function of Semilinear Parabolic Infinite Time Horizon Optimal Control Problems on $L^2(\Omega)$ under Control Constraints *

Karl Kunisch [†]

Buddhika Priyasad [‡]

February 8, 2023

Abstract

An abstract framework guaranteeing the local continuous differentiability of the value function associated with optimal stabilization problems subject to abstract semilinear parabolic equations subject to a norm constraint on the controls is established. It guarantees that the value function satisfies the associated Hamilton-Jacobi-Bellman equation in the classical sense. The applicability of the developed framework is demonstrated for specific semilinear parabolic equations.

1 Introduction.

Continuous differentiability of the value function with respect to the initial datum is an important problem in optimal feedback control theory. Indeed, if the value function is C^1 then it is the solution of a Hamilton Jacobi Bellman (HJB) equation and its negative gradient can be used to define an optimal state feedback law. The subject matter of this paper addresses local continuous differentiability of the value function \mathcal{V} for infinite horizon optimal control problems subject to semilinear parabolic control problems and norm constraints on the control. Such problems are intimately related to stabilization problems which are often cast as infinite horizon optimal control problems. Investigating infinite horizon problems constitutes one of the specificities of this paper. Another one is the fact that we focus on the differentiability of \mathcal{V} on (subsets of) $L^2(\Omega)$. Thus we need to consider the semilinear equations with initial data $y_0 \in L^2(\Omega)$. As a consequence the solutions of the semilinear equations only enjoy low Sobolev-space regularity. This restricts the class of nonlinearities, compared to those which are admissible if the states are in $L^\infty((0, \infty) \times \Omega)$, which is the situation typically addressed in the literature on optimal control [Cas] and [Tro2]. The latter necessitates to take the initial conditions in spaces strictly smaller than $L^2(\Omega)$. Here we consider $L^2(\Omega)$, first due to intrinsic interest, secondly because ultimately the HJB equation should be solved numerically, which is easier in an $L^2(\Omega)$ setting than in other topologies, like $H^1(\Omega)$. Let

*The authors were supported by the ERC advanced grant 668998 (OCLOC) under the EU's H2020 research program.

[†]Institute for Mathematics and Scientific Computing, University of Graz, Heinrichstrasse 36, A-8010 Graz, Austria, and Radon Institute, Austrian Academy of Science, Linz, Austria. (karl.kunisch@uni-graz.at).

[‡]Institute for Mathematics and Scientific Computing, University of Graz, Heinrichstrasse 36, A-8010 Graz, Austria. (b.sembukutti-liyanage@uni-graz.at).

us also recall that one of the approaches to solve the HJB equation is given by the policy iteration. It assumes that the value function is C^1 .

The underlying analysis demands stability and sensitivity analysis of infinite dimensional optimal control problems subject to nonlinear equations. For this purpose we utilize the theory of generalized equations as established by [Don] and [Rob]. It involves first order approximations of the state and adjoint equations, which lead to restrictions on the class of nonlinearities which can be admitted. We refer to the section on examples in this respect.

The current investigations are to some degree a continuation of work the first author's work on optimal feedback control for infinite dimensional systems. In [BKP1, BKP2, BKP3] Taylor approximations of the value function for problems with a concrete structure, namely, bilinear control systems, and the Navier Stokes equations were investigated and differentiability of the value function was obtained as a by product. In these investigations norm constraints were not considered. Here we admit norm constraints and we focus on semilinear equations. Let us also notice that the systems investigated in [BKP1, BKP2, BKP3] share the property that the second derivatives with respect to the state variable of the nonlinearity in the state equation do not depend on the state itself anymore.

Let us also compare our work to the developments in the field of parametric sensitivity analysis of semilinear parabolic equations under control constraints. There are many papers focusing on stability and sensitivity analysis of finite time horizon problems along with pointwise control constraints, see e.g. [BM, GHH, Gri, GV, Mal, MT, Tro1, Wac], and the literature there. First, of these papers, except for [GV, Tro1], consider the case with initial data in $H^1(\Omega)$ or $C(\bar{\Omega})$. In [GV] again the third derivative of the nonlinearity is zero. Secondly, all of them consider the finite horizon case. Since we treat infinite horizon problems we have to guarantee stabilizability (for small initial data) under control constraints. Then we use a fixed point argument to obtain well-posedness of the system. Well-posedness and stability with respect to parameters of the adjoint equation is significantly more involved for infinite horizon problems than for finite horizon problems. It requires techniques, differently from those used in the finite horizon case. Another aspect is the proper characterization of the adjoint state at $t = \infty$.

In the finite dimensional case, there is, of course a tremendous amount of work on the treatment of the value function if it is not C^1 . Fewer papers concentrate on the case where the value function enjoys smoothness properties. We mention [Goe] and [CF] in this respect.

In order to achieve the goal we desire, we lay out the following setup. In Section 2, we consider an abstract parametric optimization problem with an equality constraint and another convex constraint. Existence of an optimal solution, of a multiplier associate to the equality constraint, and Lipschitz stability of the component of the state variable which lies in the complement of the kernel of the linearized constraint will be established. This result is necessary but not sufficient for the further developments, since stability is obtained in a norm which is too weak and since the stability estimate does not involve the component in the kernel of the linearized constraint and the multiplier, i.e. the adjoint states, yet. At the level of Section 2 this remains as Assumption (H7). In Section 3 we specify the concrete optimal stabilization problem and a set of conditions, most importantly on the nonlinearity of the state equation, under which Assumption (H7) can be established, for initial data $y_0 \in L^2(\Omega)$. Section 3 also contains a summary of the main results of this paper. They are stated as theorems with a little stronger assumptions than eventually necessary, for the saker of easing the presentation. Section 4 is dedicated to the proof of verifying the assumptions of the general setup of Section 2 for the concrete optimal control problem stated in Section 3. As conclusion we obtain the Lipschitz continuity in the appropriate norms of the all

variables appearing in the optimality system with respect to the parameter of interest, which is the initial condition y_0 , in our case. Since our analysis is a local one involving second order optimality conditions, solutions to the optimality system are related to local solutions to the optimal control problem. As a corollary to these results we obtain that the local value function is Fréchet differentiable. In Section 5, we show that in the neighborhood of global solutions the value function \mathcal{V} satisfies the Hamilton-Jacobi-Bellman (HJB) equation in the strong sense. Finally, Section 6 is devoted to demonstrating that the developed framework is applicable for some concrete examples, namely for linear systems, Fisher's equations, and parabolic equations with global Lipschitz nonlinearities. All our results require a smallness assumption on the initial conditions y_0 . Two aspects need to be taken into consideration in this respect. First y_0 has to be sufficiently small so that the controlled system is stable. Secondly a second order optimality condition is needed. For this to hold a sufficient condition is provided by smallness of the adjoint state, which in turn can be implied by smallness of y_0 . We stress that these two issues are of related, but independent nature.

2 Lipschitz stability for an abstract optimization problem.

Here we present a stability result for an abstract, infinite dimensional optimization problem which will be the building block for the results below. This result is geared towards exploiting the specific nature of optimization problem with differential equations as constraints. First existence of a dual variable will result from a regular point condition. Subsequently the Lipschitz stability result is obtained in two steps. In the first one, we rely on the relationship between the linearized optimality conditions and an associated linear-quadratic optimal optimization problem, with an extra convex constraint. This approach is useful since it provides the existence of solutions to the linearized system on the basis of variational techniques. However it dictates a certain norms for the involved quantities. These norms are too weak for our goal of obtaining Lipschitz continuity of the adjoint variables in such a manner that differentiability of the cost with respect to the initial conditions can be argued. Therefore, in a second step we exploit the specific structure of the optimality systems, using the fact that it is related to a parabolic optimal control problem, to obtain the Lipschitz continuity in the stronger norms. This two step approach is also present in some of the earlier work on stability and sensitivity analysis which was quoted in the introduction. But due to that fact these papers considered finite horizon problems it came as a byproduct which improved the regularity of the adjoints. In our work it is essential to reach our goal. This is why we decided to formalize this two step approach which was not done in earlier work.

Concretely, we consider the optimization problem

$$\begin{cases} \min f(x) \\ e(x, q) = 0, x \in C. \end{cases} \quad (P_q)$$

with a parameter dependent equality constraint, and a general constraint described by $x \in C$, where C is a closed convex subset of a real Hilbert space X . Further, W is a real Hilbert space and P is a normed linear space. In the application that we have in mind, the parameter q will appear as the initial condition in the dynamical system. The following Assumption (H1) is assumed to hold throughout.

Assumption H1.

$q_0 \in P$ is a nominal reference parameter,

x_0 is a local solution (P_{q_0}) ,

$f : X \rightarrow \mathbb{R}^+$ is twice continuously differentiable in a neighborhood of x_0 ,

$e : X \times P \rightarrow W$ is continuous, and twice continuously differentiable w.r.t. x , with first and second derivative Lipschitz continuous in a neighborhood of (x_0, q_0) .

The derivatives with respect to x will be denoted by primes and the derivatives w.r.t. y and u later on, are denoted by subscripts. They are all considered in the sense of Lebesgue derivatives.

We introduce the Lagrangian $\mathcal{L} : X \times P \times W^* \rightarrow \mathbb{R}$ associated to (P_q) by

$$\mathcal{L}(x, q, \lambda) = f(x) + \langle \lambda, e(x, q) \rangle_{W^*, W}. \quad (2.1)$$

Next further relevant assumptions are introduced:

Assumption H2 (regular point condition).

$$0 \in \text{int } e'(x_0, q_0)(C - x_0),$$

where *int* denotes the interior in the W topology. This regularity condition implies the existence of a Lagrange multiplier $\lambda_0 \in W^*$, see e.g. [MZ] such that the following first order condition holds:

$$\begin{cases} \langle \mathcal{L}'(x_0, q_0, \lambda_0), c - x_0 \rangle_{X^*, X} \geq 0, \quad \forall c \in C, \\ e(x_0, q_0) = 0. \end{cases} \quad (2.2)$$

It is equivalent to

$$\begin{cases} 0 \in \mathcal{L}'(x_0, q_0, \lambda_0) + \partial \mathbf{I}_C(x_0), & \text{in } X^*, \\ e(x_0, q_0) = 0, & \text{in } W, \end{cases} \quad (2.3)$$

where $\partial \mathbf{I}_C(x)$ denotes the subdifferential of the indicator function of the set C at $x \in X$. Let $A \in \mathcal{L}(X, X^*)$ denote the operator representation of $\mathcal{L}''(x_0, q_0, \lambda_0)$, i.e.

$$\langle Ax_1, x_2 \rangle_{X^*, X} = \mathcal{L}''(x_0, q_0, \lambda_0)(x_1, x_2) \quad (2.4)$$

and define

$$E = e'(x_0, q_0) \in \mathcal{L}(X, W). \quad (2.5)$$

We further require

Assumption H3 (positive definiteness).

$$\exists \kappa > 0 : \langle Ax, x \rangle_{X^*, X} \geq \kappa \|x\|_X^2, \quad \forall x \in \ker E.$$

Condition (H3) is a bit stronger than a second order sufficient optimality condition, since it does not take into consideration the activity or inactivity of the constraints. Such weaker second order conditions typically allow to derive quadratic positive definite lower bounds on the cost and Hölder continuity with respect to perturbations. For Lipschitz continuity and differentiability stronger assumptions, such as (H3) are typically assumed. We refer exemplarily to [Gri, GHH, GV, Wac], and [IK, Section 2.3]. The constraints in these references, however, are not identical with those of the present paper.

The stability result of (x_0, λ_0) with respect to perturbation of q at q_0 will be based on Robinson's strong regularity condition which involves the following linearized form of the optimality condition,

$$\begin{cases} 0 \in \mathcal{L}'(x_0, q_0, \lambda_0) + A(x - x_0) + E^*(\lambda - \lambda_0) + \partial \mathbf{I}_C(x) & \text{in } X^*, \\ 0 = e(x_0, q_0) + E(x - x_0) & \text{in } W. \end{cases} \quad (2.6)$$

We define a multivalued operator $\mathcal{T} : X \times W^* \rightarrow X^* \times W$ by

$$\mathcal{T} \begin{pmatrix} x \\ \lambda \end{pmatrix} = \begin{pmatrix} A & E^* \\ E & 0 \end{pmatrix} \begin{pmatrix} x \\ \lambda \end{pmatrix} + \begin{pmatrix} f'(x_0) - Ax_0 \\ -Ex_0 \end{pmatrix} + \begin{pmatrix} \partial \mathbf{I}_C(x) \\ 0 \end{pmatrix}, \quad (2.7)$$

and observe that (2.6) is equivalent to

$$0 \in \mathcal{T} \begin{pmatrix} x \\ \lambda \end{pmatrix}.$$

Here it is understood that \mathcal{T} is evaluated at $(x_0, q_0, \lambda_0) \in X \times P \times W^*$. But \mathcal{T} is not yet the mapping for which we need to verify the Robinson-Dontchev strong regularity condition in our context. It relates to the fact that we must to treat the multiplier λ in smaller space than W^* . Before we can properly specify this condition some additional preparation is necessary. We first introduce Banach spaces:

$$\underline{X} \subset X, \quad \underline{W}^* \subset W^*, \quad \underline{X}^* \subset X^*, \quad (2.8)$$

with continuous injections. We emphasize that \underline{X}^* should not be confused with $(\underline{X})^*$. A restriction of \mathcal{T} will be defined as multivalued operator $\underline{\mathcal{T}} : \underline{X} \times \underline{W}^* \rightarrow \underline{X}^* \times W$. Indeed, in applications to optimal control problems extra regularity of multipliers can be obtained by investigating the solutions (2.3), see e.g. Section 3. In the context of optimal stabilization problems this structural property will become transparent in Proposition 4.1 and Proposition 4.2, see also [BKP3, Proposition 15]. It will turn out to be essential for our purposes. But this situation where the multiplier has extra regularity is also of abstract interest. When studying stability in this setting this means that the second coordinate of the domain of \mathcal{T} needs to be changed from W^* to \underline{W}^* . This entails that the range space of \mathcal{T} has to be modified appropriately, in order to obtain stability of the λ coordinate. For this purpose we introduce $\underline{X}^* \subset X^*$. The reason for further restricting X to \underline{X} will become evident in the proof of Proposition 4.2. It is related to the fact that we consider infinite horizon problems. A concrete use of these space is elaborated in detailed in subsection 3.2.2.

Now we adapt the conditions on f and e to the choice of the spaces in (2.8).

Assumption H4.

There exists a neighborhood $\tilde{U}_1 \times \tilde{U}_2 \subset \underline{X} \times P$ of (x_0, q_0) such that

- (i) the restriction of $x \mapsto f'(x)$ to \underline{X} defines a mapping $\underline{f}'(x)$ from $\tilde{U}_1 \subset \underline{X}$ to \underline{X}^* ,
- (ii) the restriction $e'(x, q)^* \in \mathcal{L}(W^*, X^*)$ to \underline{W}^* defines operators $\underline{e}'(x, q)^* \in \mathcal{L}(\underline{W}^*, \underline{X}^*)$ for every $(x, q) \in \tilde{U}_1 \times \tilde{U}_2$.

With these assumption holding we define the restricted linearized Lagrangian

$$\underline{\mathcal{L}}' : \tilde{U}_1 \times \tilde{U}_2 \times \underline{W}^* \subset \underline{X} \times P \times \underline{W}^* \rightarrow \underline{X}^* \quad \text{by} \quad \underline{\mathcal{L}}'(x, q, \lambda) = \underline{f}'(x) + \underline{e}'(x, q)^* \lambda. \quad (2.9)$$

Next we adapt $\partial\mathbf{I}_C \subset X^*$ to the situation of (2.8) and define for $x \in \underline{X}$ the set valued mapping

$$\underline{\partial\mathbf{I}_C}(x) = \{y \in \underline{X}^* : \langle y, v - x \rangle_{X^*, X} \leq 0, \forall v \in C \cap \underline{X}\} \subset \underline{X}^*. \quad (2.10)$$

We henceforth assume that $(x_0, \lambda_0) \in \underline{X} \times \underline{W}^*$, it will also follow as a special case of (H7) below. The following assumption will guarantee that the restriction $\underline{\mathcal{T}}$ of \mathcal{T} is well-defined as operator from $\underline{X} \times \underline{W}^*$ to $\underline{X}^* \times W$, and the one beyond is needed for Lipschitz continuous dependence of local solutions to (P_q) with respect to q .

Assumption H5.

$\underline{\mathcal{L}}' : \tilde{U}_1 \times \tilde{U}_2 \times \underline{W}^* \subset \underline{X} \times P \times \underline{W}^* \rightarrow \underline{X}^*$ is Fréchet differentiable with respect to x , and $(\underline{\mathcal{L}})'$, as a mapping $(x, q, \lambda) \mapsto (\underline{\mathcal{L}})'(x, q, \lambda)$, is continuous at $(x_0, q_0, \lambda_0) \in \underline{X} \times P \times \underline{W}^*$.

Assumption H6. There exists $\nu > 0$ such that:

$$\|e(x, q_1) - e(x, q_2)\|_W \leq \nu \|q_1 - q_2\|_P, \quad \forall (x, q_1) \text{ and } (x, q_2) \in \tilde{U}_1 \times \tilde{U}_2, \quad (2.11a)$$

$$\left\| \underline{e}'(x, q_1)^* - \underline{e}'(x, q_2)^* \right\|_{\mathcal{L}(W^*, X^*)} \leq \nu \|q_1 - q_2\|_P, \quad \forall (x, q_1) \text{ and } (x, q_2) \in \tilde{U}_1 \times \tilde{U}_2. \quad (2.11b)$$

Let us further set

$$\underline{E}^* = \underline{e}'(x_0, q_0)^* \in \mathcal{L}(W^*, X^*) \text{ and } \underline{A} = (\underline{\mathcal{L}}'(x_0, q_0, \lambda_0))' \in \mathcal{L}(\underline{X}, X^*).$$

With Assumptions (H1)-(H5) holding (2.3) can be expressed as

$$\begin{cases} 0 \in \underline{\mathcal{L}}'(x_0, q_0, \lambda_0) + \underline{\partial\mathbf{I}_C}(x_0), & \text{in } \underline{X}^*, \\ e(x_0, q_0) = 0, & \text{in } W. \end{cases} \quad (2.12)$$

Moreover (2.6) restricted to $\underline{X} \times \underline{X}^*$ result in:

$$0 \in \begin{cases} \underline{\mathcal{L}}'(x_0, q_0, \lambda_0) + \underline{A}(x - x_0) + \underline{E}^*(\lambda - \lambda_0) + \underline{\partial\mathbf{I}_C}(x) & \text{in } \underline{X}^*, \\ e(x_0, q_0) + E(x - x_0) & \text{in } W, \end{cases} \quad (2.13)$$

and the multivalued operator $\underline{\mathcal{T}} : \underline{X} \times \underline{W}^* \rightarrow \underline{X}^* \times W$ related to (2.7) is defined as

$$\underline{\mathcal{T}} \begin{pmatrix} x \\ \lambda \end{pmatrix} = \begin{pmatrix} \underline{A} & \underline{E}^* \\ E & 0 \end{pmatrix} \begin{pmatrix} x \\ \lambda \end{pmatrix} + \begin{pmatrix} f'(x_0) - \underline{A}x_0 \\ -Ex_0 \end{pmatrix} + \begin{pmatrix} \underline{\partial\mathbf{I}_C}(x) \\ 0 \end{pmatrix}. \quad (2.14)$$

Observe that (2.13) is equivalent to

$$0 \in \underline{\mathcal{T}} \begin{pmatrix} x \\ \lambda \end{pmatrix}.$$

Existence and Lipschitz continuity of solutions in a neighborhood (x_0, q_0, λ_0) will follow from the strong regularity assumption which requires us to show that there exist neighborhoods $\hat{V} \subset \underline{X}^* \times W$ of 0 and $\hat{U} = \hat{U}_1 \times \hat{U}_2 \subset \underline{X} \times \underline{W}^*$ of (x_0, q_0) such that $\underline{\mathcal{T}}^{-1}$ has the properties that $\underline{\mathcal{T}}^{-1}(\hat{V}) \cap \hat{U}$ is single-valued and that it is Lipschitz continuous from \hat{V} to \hat{U} , see [Don], (and also [Rob], [IK, Definition 2.2, p 31], in case $\underline{X} = X$, $\underline{W}^* = W^*$, $\underline{X}^* = X^*$). We approach the strong regularity assumption in two steps. In the first one we argue invertibility of \mathcal{T} and Lipschitz continuity of the variable x in X . For this purpose we exploit the symmetry of \mathcal{T} and consider an associated variational problem. In our specific situation the inverse of \mathcal{T} - and consequently of $\underline{\mathcal{T}}$ - is single-valued and thus the restriction to the neighborhood \hat{U} is not needed. Existence and Lipschitz continuity of λ as well as Lipschitz continuity of x in the small space $\underline{X} \times \underline{W}^*$ remains an assumption in the generality of problem (P_q) . It will be verified in a second step for the optimal stabilization problems in the following sections.

Assumption H7. For $(\beta_1, \beta_2) \in \hat{V} \subset \underline{X}^* \times W$, the solution $(x_{(\beta_1, \beta_2)}, \lambda_{(\beta_1, \beta_2)})$ to $\mathcal{T} \begin{pmatrix} x \\ \lambda \end{pmatrix} = \begin{pmatrix} \beta_1 \\ \beta_2 \end{pmatrix}$ lies in $\underline{X} \times \underline{W}^*$. Moreover there exists a constant $k > 0$ such that

$$\left\| x_{(\beta_1, \beta_2)} - x_{(\hat{\beta}_1, \hat{\beta}_2)} \right\|_{\underline{X}} + \left\| \lambda_{(\beta_1, \beta_2)} - \lambda_{(\hat{\beta}_1, \hat{\beta}_2)} \right\|_{\underline{W}^*} \leq k \left[\left\| (\beta_1, \beta_2) - (\hat{\beta}_1, \hat{\beta}_2) \right\|_{\underline{X}^* \times W} + \left\| x_{(\beta_1, \beta_2)} - x_{(\hat{\beta}_1, \hat{\beta}_2)} \right\|_{\underline{X}} \right]$$

for all $(\beta_1, \beta_2) \in \hat{V}$, $(\hat{\beta}_1, \hat{\beta}_2) \in \hat{V}$.

This condition is used after the existence of $x_\beta = x_{(\beta_1, \beta_2)}$ was already established. Note that for $(\beta_1, \beta_2)^T = 0$ we have $(x_{(0,0)}, \lambda_{(0,0)}) = (x_0, \lambda_0)$ and hence (H7) in particular implies that $(x_0, \lambda_0) \in \underline{X} \times \underline{W}^*$. We arrive at the announced stability result.

Theorem 2.1. Assume that (H1)-(H7) hold at a local solution x_0 of (P_{q_0}) . Then there exist a neighborhood $U = U(x_0, \lambda_0) \subset \underline{X} \times \underline{W}^*$, a neighborhood $N = N(q_0) \subset P$, and a constant μ such that for all $q \in N$ there exists a unique $(x(q), \lambda(q)) \in U$ satisfying

$$0 \in \begin{cases} \mathcal{L}'(x(q), q, \lambda(q)) + \underline{\partial \mathbf{I}_C}(x(q)), & \text{in } \underline{X}^*, \\ e(x(q), q), & \text{in } W, \end{cases} \quad (2.15)$$

and

$$\|(x(q_1), \lambda(q_1)) - (x(q_2), \lambda(q_2))\|_{\underline{X} \times \underline{W}^*} \leq \mu \|q_1 - q_2\|_P, \quad \forall q_1, q_2 \in N. \quad (2.16)$$

In addition there exists a nontrivial neighborhood $\tilde{N} \subset N$ of q_0 such that $x(q)$ is a local solution of (P_q) for $q \in \tilde{N}$.

For the proof we shall employ the following lemma in which $A \in \mathcal{L}(X, X^*)$ and $E \in \mathcal{L}(X, W)$ denote generic operators. For the sake of completeness we also include its proof.

Lemma 2.1. Let $(\tilde{a}, \tilde{b}) \in X^* \times W$, assume that $A \in \mathcal{L}(X, X^*)$ is self-adjoint and satisfies (H3), and that the set $S(\tilde{b}) = \{x \in C : Ex = \tilde{b}\}$ is nonempty. Then the problem

$$\begin{cases} \min_{x \in C} \tilde{J}(x) = \min_{x \in C} \frac{1}{2} \langle Ax, x \rangle_{X^*, X} + \langle \tilde{a}, x \rangle_{X^*, X}, \\ Ex = \tilde{b}, \end{cases} \quad (2.17)$$

admits a unique solution $x = x(\tilde{a}, \tilde{b})$ satisfying

$$\langle Ax + \tilde{a}, v - x \rangle_{X^*, X} \geq 0, \quad \text{for all } v \in S(\tilde{b}). \quad (2.18)$$

If moreover the regular point condition $0 \in \text{int } E(C - x(\tilde{a}, \tilde{b}))$ holds, then there exists $\lambda = \lambda(\tilde{a}, \tilde{b}) \in W^*$ such that

$$0 \in \left\{ \begin{pmatrix} A & E^* \\ E & 0 \end{pmatrix} \begin{pmatrix} x \\ \lambda \end{pmatrix} + \begin{pmatrix} \tilde{a} \\ -\tilde{b} \end{pmatrix} + \begin{pmatrix} \partial \mathbf{I}_C(x) \\ 0 \end{pmatrix} \right\}. \quad (2.19)$$

Proof. Since C is a closed and convex, $S(\tilde{b})$ is closed and convex. By assumption $S(\tilde{b})$ is nonempty. Hence there exists an $x \in C$ such that $Ex = \tilde{b}$. Note that each such x can be uniquely decomposed as $x = w + y$, with $y \in \ker E$, $w \in \ker E^\perp$ and $Ew = \tilde{b}$. By (H3) the functional \tilde{J} is bounded from below and coercive on $S(\tilde{b})$. Hence there exists a minimizing sequence $\{x_n\}$ in $S(\tilde{b})$ such that $\lim_{n \rightarrow \infty} \tilde{J}(x_n) = \inf_{x \in S(\tilde{b})} \tilde{J}(x)$. Each x_n can be decomposed as $x_n = w + y_n$, with $y_n \in \ker E$.

By (H3) the sequences $\{y_n\}_{n=1}^\infty$ and hence $\{x_n\}_{n=1}^\infty$ are bounded. Thus there exists a subsequence $\{x_{n_k}\}$ with weak limit $x = x(\tilde{a}, \tilde{b})$ in $S(\tilde{b})$. Since \tilde{J} weakly lower semi-continuous, we have that $\tilde{J}(x) \leq \liminf_{k \rightarrow \infty} \tilde{J}(x_{n_k})$ and x minimizes \tilde{J} over $S(\tilde{b})$. This further implies that $\langle Ax + \tilde{a}, v - x \rangle_{X^*, X} \geq 0$ for all $v \in S(\tilde{b})$. Uniqueness of x follows from (H3).

The regular point condition implies the existence of a multiplier $\lambda = \lambda(\tilde{a}, \tilde{b}) \in W^*$ such that (2.19) holds. See e.g. [IK, Theorem 1.6] \square

Proof of the Theorem 2.1.

- (i) The proof of the first assertion of the Theorem 2.1 is based on the implicit function theorem of Dontchev for generalized equations, see [Don, Theorem 2.4, Remark 2.5]. We introduce the mapping $\underline{\mathcal{F}} : \underline{X} \times P \times \underline{W}^* \rightarrow \underline{X}^* \times W$ given by

$$\underline{\mathcal{F}}(x, q, \lambda) = \begin{pmatrix} \underline{\mathcal{L}}'(x, q, \lambda) \\ e(x, q) \end{pmatrix},$$

and observe that Assumption (H6) implies that for all (x, q_1, λ) , and $(x, q_2, \lambda) \in \tilde{U}_1 \times \tilde{U}_2 \times \underline{W}^*$

$$\|\underline{\mathcal{F}}(x, q_1, \lambda) - \underline{\mathcal{F}}(x, q_2, \lambda)\|_{\underline{X}^* \times W} \leq \nu \left(1 + \|\lambda\|_{\underline{W}^*}\right) \|q_1 - q_2\|_P. \quad (2.20)$$

By (H1) and (H5), and using the integral mean value theorem it can be argued that

$$\begin{pmatrix} x \\ \lambda \end{pmatrix} \rightarrow \begin{pmatrix} A & E^* \\ E & 0 \end{pmatrix} \begin{pmatrix} x \\ \lambda \end{pmatrix} + \begin{pmatrix} f'(x_0) - Ax_0 \\ -Ex_0 \end{pmatrix}$$

strongly approximates $\underline{\mathcal{F}}$ at (x_0, q_0, λ_0) , in the sense of Dontchev, [Don]. In the next two steps the strong regularity condition for $\underline{\mathcal{T}}$ will be verified.

- (ii) (Existence). Let, at first, $(\beta_1, \beta_2) \in X^* \times W$ and consider $\mathcal{T} \begin{pmatrix} x \\ \lambda \end{pmatrix} = \begin{pmatrix} \beta_1 \\ \beta_2 \end{pmatrix}$ which is equivalent to,

$$0 \in \begin{pmatrix} a \\ -b \end{pmatrix} + \begin{pmatrix} A & E^* \\ E & 0 \end{pmatrix} \begin{pmatrix} x \\ \lambda \end{pmatrix} + \begin{pmatrix} \partial \mathbf{I}_C(x) \\ 0 \end{pmatrix}, \quad (2.21)$$

with $a = f'(x_0) - Ax_0 - \beta_1$, $b = Ex_0 + \beta_2$, and A, E defined in (2.4), (2.5). To solve (2.21) we consider,

$$\begin{cases} \min_{x \in C} \frac{1}{2} \langle Ax, x \rangle_{X^*, X} + \langle a, x \rangle_{X^*, X}, \\ Ex = b. \end{cases} \quad (2.22)$$

This corresponds to (2.17) with $\tilde{a} = a$, $\tilde{b} = b$ and feasible set $S(\beta_2) = \{x \in C : Ex = b\}$. Clearly $x_0 \in S(0) = \{x \in C : Ex = Ex_0\}$. By (H2) and [IK, Theorem I.2.8], there exists a neighborhood of the origin $\tilde{V} \subset X^* \times W$ such that $S(\beta_2)$ is not empty for all $(\beta_1, \beta_2) \in \tilde{V}$. Thus by Lemma 2.1 there exists a unique solution $x = x(\beta_1, \beta_2)$ to (2.22) for each $(\beta_1, \beta_2) \in \tilde{V}$. By [IK, Theorem I.2.11, I.2.12, I.2.15], possibly after reducing \tilde{V} , these solutions depend Hölder continuously on $(\beta_1, \beta_2) \in \tilde{V} \subset X^* \times W$, with exponent $\frac{1}{2}$. The regular point condition for the solution $x(\beta_1, \beta_2)$ is

$$0 \in \text{int } E(C - x(\beta_1, \beta_2)) = \text{int } E(C - x_0) - \beta_2,$$

which is satisfied due to **(H2)**, possibly after again shrinking \tilde{V} . Hence there exists a Lagrange multiplier $\lambda = \lambda(\beta_1, \beta_2)$ associated to $Ex = b$, and **(2.21)** admits a unique solution $(x(\beta_1, \beta_2), \lambda(\beta_1, \beta_2))$ since it is the first order optimality condition for **(2.17)**.

(iii) (Uniqueness and Lipschitz continuity) Let $(\beta_1, \beta_2) \in \tilde{V}$ and $(\hat{\beta}_1, \hat{\beta}_2) \in \tilde{V}$ with corresponding solutions $(x, \lambda) \in X \times W^*$ and $(\hat{x}, \hat{\lambda}) \in X \times W^*$. This implies that

$$\begin{cases} \langle a + Ax + E^*\lambda, c - x \rangle_{X^*, X} \geq 0, \quad \forall c \in C, \\ Ex = b, \quad \text{with } a = f'(x_0) - Ax_0 - \beta_1, \quad b = Ex_0 + \beta_2, \end{cases} \quad (2.23)$$

and

$$\begin{cases} \langle \hat{a} + A\hat{x} + E^*\hat{\lambda}, c - \hat{x} \rangle_{X^*, X} \geq 0, \quad \forall c \in C, \\ E\hat{x} = \hat{b}, \quad \text{with } \hat{a} = f'(x_0) - Ax_0 - \hat{\beta}_1, \quad \hat{b} = Ex_0 + \hat{\beta}_2. \end{cases} \quad (2.24)$$

By the first equations in **(2.23)** and **(2.24)** we obtain that

$$\langle a + Ax + E^*\lambda, \hat{x} - x \rangle_{X^*, X} \geq 0, \quad \langle \hat{a} + A\hat{x} + E^*\hat{\lambda}, x - \hat{x} \rangle_{X^*, X} \geq 0, \quad x, \hat{x} \in C. \quad (2.25)$$

Combining these inequalities, we have that

$$\langle a - \hat{a} + A(x - \hat{x}) + E^*(\lambda - \hat{\lambda}), x - \hat{x} \rangle_{X^*, X} \leq 0. \quad (2.26)$$

The second equalities in **(2.23)** and **(2.24)** imply that

$$E(x - \hat{x}) = b - \hat{b}. \quad (2.27)$$

Let us set

$$\delta x = \hat{x} - x, \quad \delta \lambda = \hat{\lambda} - \lambda, \quad \delta a = \hat{a} - a, \quad \delta b = \hat{b} - b.$$

Then $\delta\beta_1 = -(\hat{\beta}_1 - \beta_1)$ and $\delta\beta_2 = \hat{\beta}_2 - \beta_2$, and **(2.26)**, **(2.27)** result in

$$\langle \delta x, A\delta x \rangle_{X, X^*} + \langle \delta \lambda, E\delta x \rangle_{W^*, W} - \langle \delta\beta_1, \delta x \rangle_{X^*, X} \leq 0, \quad (2.28)$$

and

$$E\delta x = \delta b. \quad (2.29)$$

By **(H2)** the operator E is surjective. Hence by the closed range theorem expressing $\delta x = \delta v + \delta w \in \ker E + \text{range } E^*$ implies that $E\delta x = E\delta w = \delta b$. Again by the closed range theorem there exists $k_1 > 0$:

$$\|\delta w\|_X \leq k_1 \|\delta\beta_2\|_W. \quad (2.30)$$

From the first equation in **(2.21)** we have

$$Ax + E^*\lambda - Ax_0 + f'(x_0) - \beta_1 \in -\partial\mathbf{I}_C(x).$$

Next we restrict the perturbation parameters to satisfy $(\beta_1, \beta_2) \in (\underline{X}^* \times W) \cap \tilde{V}$. Due to Assumptions **(H4)** and **(H7)** we have $(x, \lambda) \in \underline{X} \times \underline{W}^*$,

$$\underline{Ax} + \underline{E}^*\lambda - \underline{Ax}_0 + \underline{f}'(x_0) - \beta_1 \in \underline{X}^*$$

and hence

$$\underline{Ax} + \underline{E}^*\lambda - \underline{Ax}_0 + \underline{f}'(x_0) - \beta_1 \in -\underline{\partial}\mathbf{I}_C(x).$$

The analogous equation holds with (x, λ, β_1) replaced by $(\hat{x}, \hat{\lambda}, \hat{\beta}_1)$.

By (H3), (2.28) and Assumption (H7) we find

$$\begin{aligned}
\kappa \|\delta v\|^2 &\leq \langle \delta v, A\delta v \rangle_{X, X^*} = \langle \delta x, A\delta x \rangle_{X, X^*} - 2\langle \delta v, A\delta w \rangle_{X, X^*} - \langle \delta w, A\delta w \rangle_{X, X^*} \\
&\leq -\langle \delta \lambda, E\delta x \rangle_{W^*, W} + \langle \delta \beta_1, \delta x \rangle_{X, X^*} - 2\langle \delta v, A\delta w \rangle_{X, X^*} - \langle \delta w, A\delta w \rangle_{X, X^*} \\
&\leq \tilde{k} \|\delta \lambda\|_{\underline{W}^*} \|\delta \beta_2\|_W + \|\delta \beta_1\|_{X^*} \|\delta x\|_X + \|A\| \|\delta w\|_X (2\|\delta v\|_X + \|\delta w\|_X) \\
&\leq \tilde{k}k(\|\delta w\|_X + \|\delta v\|_X + \|\delta \beta\|_{\underline{X}^* \times W}) \|\delta \beta_2\|_W + (\|\delta \beta_1\|_{X^*} + 2\|A\| \|\delta w\|_X)(\|\delta v\|_X + \|\delta w\|_X),
\end{aligned} \tag{2.31}$$

where \tilde{k} denotes the embedding constant of \underline{W}^* into W^* . Using (2.30) and rearranging terms there exists a constant $k_2 > 0$ such that

$$\|\delta v\|_X \leq k_2 \left(\|\delta \beta_1\|_{\underline{X}^*} + \|\delta \beta_2\|_W \right). \tag{2.32}$$

Applying (2.30) again this implies the existence of k_3 such that

$$\|\delta x\|_X \leq k_3 \left(\|\delta \beta_1\|_{\underline{X}^*} + \|\delta \beta_2\|_W \right) \text{ for all } (\beta_1, \beta_2) \in (\underline{X}^* \times W) \cap \tilde{V}. \tag{2.33}$$

Another application of (H7) and (2.33) imply the existence of a constant k_4 and a neighborhood \hat{V} of the origin in $\underline{X}^* \times W$ such that the desired Lipschitz stability estimate for $(\mathcal{T})^{-1}$

$$\|\delta x\|_{\underline{X}} + \|\delta \lambda\|_{\underline{W}^*} \leq k_4 \left(\|\delta \beta_1\|_{\underline{X}^*} + \|\delta \beta_2\|_W \right) \text{ for all } (\beta_1, \beta_2) \in \hat{V} \subset \underline{X}^* \times W \tag{2.34}$$

holds.

(iv) As a consequence of the previous two steps \mathcal{T} is strongly regular at (x_0, q_0, λ_0) . Together with step (i), Dontchev's theorem is applicable [Don, Theorem 2.4, Remark 2.5], and (2.15), and (2.16) follow.

(v) (Local solution to (P_q)) Now we show that there exists a neighborhood \tilde{N} of q_0 such that for $q \in \tilde{N}$ the second order sufficient optimality condition is satisfied at $x(q)$, so that $x(q)$ is a local solution of (P_q) by eg. [IK, Theorem 2.12, p42]. Due to (H3) and regularity of f, e we obtain

$$\mathcal{L}''(x(q), q, \lambda(q))(h, h) \geq \frac{\kappa}{2} \|h\|^2, \text{ for all } h \in \ker E, \text{ if } q \in N(q_0). \tag{2.35}$$

Let us define $E_q = (e_y(x(q), q))$ for $q \in N(q_0)$. By the surjectivity of E_{q_0} and regularity of e there exists a neighborhood $\tilde{N} \subset N(q_0)$ such that E_q is surjective for all $q \in \tilde{N}$. Here we also use continuity of $q \mapsto e_y(x(q), q)$ from $P \rightarrow W$ at q_0 , which follows from (H1) and the continuity of $q \rightarrow x(q)$ at q_0 . Consequently exist $\delta_0, \gamma > 0$ such that

$$\mathcal{L}''(x(q), q, \lambda(q))(h + z, h + z) \geq \delta_0 \|h + z\|^2, \text{ for all } h \in \ker E, z \in X \tag{2.36}$$

satisfying $\|z\| \leq \gamma \|h\|$ by [IK, Lemma 2.13, p43]. Let us define the orthogonal projection onto $\ker E_q$ given by $P_{\ker E_q} = I - E_q^*(E_q E_q^*)^{-1} E_q$. We choose \tilde{N} so that

$$\|E_q^*(E_q E_q^*)^{-1} E_q - E_{q_0}^*(E_{q_0} E_{q_0}^*)^{-1} E_{q_0}\| \leq \frac{\gamma}{1 + \gamma}$$

for all $q \in \tilde{N}$. For $x \in \ker E_q$, we have $x = h + z$ for $h \in \ker E$, $z \in (\ker E)^\perp$ and $\|x\|^2 = \|h\|^2 + \|z\|^2$. Thus,

$$\|z\| \leq \|E_q^*(E_q E_q^*)^{-1} E_q x - E_{q_0}^*(E_{q_0} E_{q_0}^*)^{-1} E_{q_0} x\| \leq \frac{\gamma}{1+\gamma} (\|h\| + \|z\|)$$

and hence $\|z\| \leq \gamma \|h\|$. From (2.35) this implies

$$\mathcal{L}''(x(q), q, \lambda(q)) \geq \delta_0 \|x\|^2, \text{ for all } x \in \ker E.$$

This concludes the proof. \square

3 Differentiability of value function for optimal stabilization subject to semi-linear parabolic equations.

Here we describe the optimal control problems which we shall analyze and state the main results.

3.1 Notation

Let Ω be an open connected bounded subset of \mathbb{R}^d with a Lipschitz continuous boundary Γ . The associated space-time cylinder is denoted by $Q = \Omega \times (0, \infty)$ and the associated lateral boundary by $\Sigma = \Gamma \times (0, \infty)$. We define the Hilbert spaces

$$Y = L^2(\Omega), \quad V = H_0^1(\Omega), \quad \text{and} \quad U = L^2(0, \infty; \mathcal{U}),$$

where \mathcal{U} is a Hilbert space which will be identified with its dual. Observe that the embedding $V \subset Y$ is dense and compact. Further $V \subset Y \subset V^*$, is a Gelfand triple. Here V^* denotes the topological dual of V with respect to the pivot space Y . For any $T \in (0, \infty)$ we define the space

$$W(0, T) = \left\{ y \in L^2(0, T; V); \frac{dy}{dt} \in L^2(0, T; V^*) \right\},$$

endowed with the norm

$$\|y\|_{W(0, T)} = \left(\|y\|_{L^2(0, T; V)}^2 + \left\| \frac{dy}{dt} \right\|_{L^2(0, T; V^*)}^2 \right)^{1/2}.$$

For $T = \infty$, we write W_∞ and $I = (0, \infty)$. We further set $W_\infty^0 = \{y \in W_\infty : y(0) = 0\}$. We also set

$$W(T, \infty) = \left\{ y \in L^2(T, \infty; V); \frac{dy}{dt} \in L^2(T, \infty; V^*) \right\}.$$

We shall frequently use that W_∞ embeds continuously into $C([0, \infty), Y)$, see e.g. [LM, Theorem 4.2] and that $\lim_{t \rightarrow \infty} y(t) = 0$, for $y \in W_\infty$, see e.g. [CK]. The set of admissible controls U_{ad} is chosen to be

$$U_{ad} \subset \{u \in U : \|u(t)\|_{\mathcal{U}} \leq \eta, \text{ for a.e. } t > 0\} \quad (3.1)$$

where η is a positive constant. We further set $\mathcal{U}_{ad} = \{v \in \mathcal{U} : \|v\|_{\mathcal{U}} \leq \eta\}$ and denote by $\mathbb{P}_{\mathcal{U}_{ad}}$ the projection of \mathcal{U} on \mathcal{U}_{ad} . For this choice of admissible controls the dynamical system can be stabilized for all sufficiently small initial conditions in Y , see Corollary 4.3 and Remark 4.1.

For $\delta > 0$ and $\bar{y} \in Y$, we define the open neighborhoods $B_Y(\delta) = \{y \in Y : \|y\|_Y < \delta\}$, and $B_Y(\bar{y}, \delta) = \{y \in Y : \|y - \bar{y}\|_Y < \delta\}$.

3.2 Problem formulation and assumptions.

We focus on the stabilization problem for an abstract semi-linear parabolic equation formulated as infinite horizon optimal control problem under control constraints:

$$(\mathcal{P}) \quad \mathcal{V}(y_0) = \min_{(y,u) \in W_\infty \times U_{ad}} J(y, u) = \min_{(y,u) \in W_\infty \times U_{ad}} \frac{1}{2} \int_0^\infty \|y(t)\|_Y^2 dt + \frac{\alpha}{2} \int_0^\infty \|u(t)\|_{\mathcal{U}}^2 dt, \quad (3.2a)$$

subject to the semilinear parabolic equation

$$y_t = \mathcal{A}y + \mathcal{F}(y) + Bu \quad \text{in } L^2(I; V^*) \quad (3.2b)$$

$$y(x, 0) = y_0 \quad \text{in } Y. \quad (3.2c)$$

Throughout \mathcal{F} is the substitution operator associated to a mapping $f: \mathbb{R} \rightarrow \mathbb{R}$ so that $(\mathcal{F}y)(t) = f(y(t))$. Sufficient conditions which guarantee the existence of solutions to (3.2b), (3.2c), as well as solutions (\bar{y}, \bar{u}) to (P), for $y_0 \in Y$ sufficiently small, will be given below. We shall also make use of the adjoint equation associated to an optimal state \bar{y} , given by

$$-p_t - \mathcal{A}^*p - \mathcal{F}'(\bar{y})^*p = -\bar{y} \quad \text{in } L^2(I; V^*). \quad (3.2d)$$

Its adjoint state p which will be considered in $L^2(I; V)$ or in W_∞ . The following assumption will be essential.

3.2.1 Assumptions A.

A1 The operator \mathcal{A} with domain $\mathcal{D}(\mathcal{A}) \subset Y$ and range in Y , generates a strongly continuous analytic semigroup $e^{\mathcal{A}t}$ on Y and can be extended to $\mathcal{A} \in \mathcal{L}(V, V^*)$.

A2 $B \in \mathcal{L}(U, Y)$ and there exists a stabilizing feedback operator $K \in \mathcal{L}(Y, U)$ such that the semigroup $e^{(\mathcal{A}-BK)t}$ is exponentially stable on Y .

A3 The nonlinearity $\mathcal{F}: W_\infty \rightarrow L^2(I; V^*)$ is twice continuously Fréchet differentiable, with second Fréchet derivative \mathcal{F}'' bounded on bounded subsets of W_∞ , and $\mathcal{F}(0) = 0$.

A4 $\mathcal{F}: W(0, T) \rightarrow L^1(0, T; \mathcal{H}^*)$ is weak-to-weak continuous for every $T > 0$, for some Hilbert space \mathcal{H} which embeds densely in V .

Note that $(L^1(0, T; \mathcal{H}^*))^* = L^\infty(0, T; \mathcal{H})$, see [Emm, Theorem 7.1.23(iv), p 164]. Moreover, $L^\infty(0, T; \mathcal{H})$ is dense in $L^2(0, T; V)$, see [MS, Lemma A.1, p 2231].

A5 $\mathcal{F}'(\bar{y}) \in \mathcal{L}(L^2(I; V), L^2(I; V^*))$.

Remark 3.1. The requirement that $\mathcal{F}(0) = 0$ in (A2) is consistent with the fact that we focus on the stabilization problem with 0 as steady state for (3.2b). Without loss of generality we further assume that

$$\mathcal{F}'(0) = 0, \quad (3.3)$$

which can always be achieved by making $\mathcal{F}'(0)$ to be perturbation of \mathcal{A} .

Remark 3.2. Let us assume that (A3) holds. Then in view of the fact that \mathcal{F} is a substitution operator we have $[\mathcal{F}'(y)v](t) = f'(y(t))v(t)$ for y and v in W_∞ , and $\mathcal{F}'(y) \in \mathcal{L}(W_\infty, L^2(I; V^*))$. Its adjoint $[\mathcal{F}'(y)^*v](t) = f'(y(t))v(t)$, for $v \in L^2(I; V)$, satisfying $\mathcal{F}'(y)^* \in \mathcal{L}(L^2(I; V), W_\infty^*)$. It has a natural restriction to an operator $\underline{\mathcal{F}'(y)^*} \in \mathcal{L}(W_\infty, L^2(I; V^*))$. With (A3) holding it is differentiable and $[\underline{\mathcal{F}'(y)^*}]'$ is a bilinear form on $W_\infty \times W_\infty$ with values in $L^2(I; V^*)$. - For examples of functions \mathcal{F} which satisfy A4 we refer to see Section 6.

3.2.2 Abstract setup.

Here we relate problem (P) to the abstract problem (P_q), which is used with the following spaces:

$$\begin{aligned} X &= W_\infty \times U, \quad W = L^2(I; V^*) \times Y, \quad P = Y, \quad C = U_{ad}, \quad X^* = W_\infty^* \times U, \quad W^* = L^2(I; V) \times Y, \\ \underline{X} &= W_\infty \times (U \cap C(\bar{I}; U)), \quad \underline{X}^* = L^2(I; V^*) \times (U \cap C(\bar{I}; U)), \quad \underline{W}^* = \widetilde{W}_\infty, \end{aligned} \quad (3.4)$$

where $I = (0, \infty)$, and $\widetilde{W} = \{(\varphi, \varphi(0)) : \varphi \in W_\infty\}$, endowed with the norm of W_∞ . At times we identify \widetilde{W} with W_∞ . We recall that the dual space of $W_\infty = L^2(I; V) \cap W^{1,2}(I; V^*)$ is $W_\infty^* = L^2(I; V^*) + (W^{1,2}(I; V^*))^*$, endowed with the norm $\|z\|_{W_\infty^*} = \inf_{z=z_1+z_2} \|z_1\|_{L^2(I; V^*)} + \|z_2\|_{(W^{1,2}(I; V^*))^*}$, where $z_1 \in L^2(I; V^*)$, $z_2 \in (W^{1,2}(I; V^*))^*$.

To express (P_q) for the present case, we set $x = (y, u) \in W_\infty \times U$, and the parameter q becomes the initial condition $y_0 \in Y$. Further $f : W_\infty \times U \rightarrow \mathbb{R}$ is given by

$$f(y, u) = \frac{1}{2} \int_0^\infty \|y(t)\|_Y^2 dt + \frac{\alpha}{2} \int_0^\infty \|u(t)\|_U^2 dt, \quad (3.5)$$

and $e(x, q) = e(y, u, y_0)$ is

$$e(y, u, y_0) = \begin{pmatrix} y_t - \mathcal{A}y - \mathcal{F}(y) - Bu \\ y(0) - y_0 \end{pmatrix} : W_\infty \times U \times Y \rightarrow L^2(I; V^*) \times Y \quad (3.6)$$

By (A3) the mapping e is Fréchet differentiable with respect to $x = (y, u) \in W_\infty \times U$ and thus for $(y, u, y_0) \in W_\infty \times U \times Y$ we have

$$e'(y, u, y_0)(v, w) = \begin{pmatrix} v_t - \mathcal{A}v - \mathcal{F}'(y)v - Bw \\ v(0, \cdot) \end{pmatrix} : W_\infty \times U \rightarrow L^2(I; V^*) \times Y. \quad (3.7)$$

The Lagrange functional $\mathcal{L} : W_\infty \times U \times Y \times L^2(I; V) \times Y \rightarrow \mathbb{R}$ corresponding to our optimal control problem is given by

$$\mathcal{L}(y, u, y_0, p, p_1) = J(y, u) + \int_0^\infty \langle p, y_t - \mathcal{A}y - \mathcal{F}(y) - Bu \rangle_{V, V^*} dt + (p_1, y(0) - y_0)_Y,$$

where $(p, p_1) \in L^2(I; V) \times Y$ corresponds to the abstract Lagrange multiplier $\lambda \in W^*$.

In the remainder of this subsection we specify the mappings \mathcal{T} and $\underline{\mathcal{T}}$ for problem (P). This will facilitate the proofs of the main results further below.

At first we take a closer look to the adjoint $\tilde{E}^* := e'(y, u, y_0)^* \in \mathcal{L}(L^2(I; V) \times Y, W_\infty^* \times U)$ at a generic element $(y, u, y_0) \in W_\infty \times U \times Y$. It is characterized by the property that for all $(v, w) \in W_\infty \times U$, $(p, p_1) \in L^2(I; V) \times Y$ we have

$$\langle \tilde{E}(v, w), (p, p_1) \rangle_{L^2(I; V^*) \times Y, L^2(I; V) \times Y} = \langle v, \tilde{E}_1^*(p, p_1) \rangle_{W_\infty, W_\infty^*} + (w, \tilde{E}_2^*(p, p_1))_U$$

where

$$\langle v, \tilde{E}_1^*(p, p_1) \rangle_{W_\infty, W_\infty^*} = \langle v_t - \mathcal{A}v - \mathcal{F}'(y)v, p \rangle_{L^2(I; V^*), L^2(I; V)} + (v(0), p_1)_Y,$$

and

$$(w, \tilde{E}_2^*(p, p_1))_U = -(w, B^*p)_U.$$

If for some $\tilde{\beta}_1 \in L^2(I; V^*)$ the pair $(p, p_1) \in L^2(I; V) \times Y$ is a solution to $\tilde{E}_1^*(p, p_1) = \tilde{\beta}_1$ then for all $v \in W_\infty$:

$$\langle v_t - \mathcal{A}v - \mathcal{F}'(y)v, p \rangle_{L^2(I; V^*), L^2(I; V)} + (v(0), p_1)_Y - (w, B^*p)_U = \langle \tilde{\beta}_1, v \rangle_{L^2(I; V^*), L^2(I; V)}. \quad (3.8)$$

Now we assume that $\mathcal{F}'(y)$ is not only an element of $\mathcal{L}(W_\infty, L^2(I; V^*))$ but rather that it can be extended to an operator $\mathcal{F}'(y) \in \mathcal{L}(L^2(I; V), L^2(I; V^*))$. This is guaranteed by (A5) at minimizers \bar{y} . Then (3.8) implies that $p \in W_\infty$, and hence $p_1 \in C(I; Y)$ and $p_1 = p(0)$, see Proposition 4.1. In particular $(p, p_1) = (p, p(0)) \in \widetilde{W}_\infty$, and (3.8) can equivalently be expressed as

$$\langle v, \tilde{E}_1^*(p, p_1) \rangle_{L^2(I; V), L^2(I; V^*)} = \langle v_t - \mathcal{A}v - \mathcal{F}'(y)v, p \rangle_{L^2(I; V^*), L^2(I; V)} = \langle v, \tilde{\beta}_1 \rangle_{L^2(I; V), L^2(I; V^*)}, \quad (3.9)$$

for all $v \in L^2(I; V)$, where we assumed that $\tilde{\beta}_1 \in L^2(I; V^*)$. Conversely, of course, if $p \in \widetilde{W}_\infty$, then $\tilde{E}_1^*(p, p(0)) = -p_t - \mathcal{A}^*p - \mathcal{F}'(y)^*p \in L^2(I; V^*)$.

From now on, let $q_0 = \bar{y}_0$ denote a reference (or nominal) parameter with associated solution $x_0 = (\bar{y}, \bar{u})$. In Proposition 4.1 we shall argue that the regular point condition Assumption (H2) is satisfied and that consequently there exists a Lagrange multiplier (\bar{p}, \bar{p}_1) such that the pair $(x_0, \lambda_0) = (\bar{y}, \bar{u}, \bar{p}, \bar{p}_1)$ satisfies (2.3). Moreover, it will turn out that $\bar{p} \in W_\infty$, $\bar{p}_1 = p(0)$, and that $\bar{u} \in U \cap C(I; \mathcal{U})$. For convenience let us present (2.3) for the present case

$$0 \in \begin{cases} \bar{y} + E_1^*(\bar{p}, \bar{p}(0)), \\ \alpha \bar{u} - B^*\bar{p} + \partial \mathbf{I}_{U_{ad}}(\bar{u}), \\ \bar{y}_t - \mathcal{A}\bar{y} - \mathcal{F}(\bar{y}) - B\bar{u}, \\ \bar{y}(0) - y_0, \end{cases} \quad (3.10)$$

where $E = \begin{pmatrix} E_1 \\ E_2 \end{pmatrix} = e'(\bar{y}, \bar{u}, \bar{y}_0)$. We stress that while the Lagrange multiplier \bar{p} belongs to W_∞ , the operator E_1^* in (3.9) is still considered as an element of $\mathcal{L}(L^2(I; V) \times Y, W_\infty^*)$.

We are now prepared to specify the multivalued operators

$$\mathcal{T} : W_\infty \times U \times L^2(I; V) \times Y \longrightarrow W_\infty^* \times U \times L^2(I; V^*) \times Y, \quad \text{and} \quad (3.11)$$

$$\mathcal{I} : W_\infty \times (U \cap C(I; \mathcal{U})) \times \widetilde{W}_\infty \longrightarrow L^2(I; V^*) \times (U \cap C(I; \mathcal{U})) \times L^2(I; V^*) \times Y \quad (3.12)$$

corresponding to (2.7) and (2.14) by

$$\mathcal{T} \begin{pmatrix} y \\ u \\ p \\ p_1 \end{pmatrix} = \begin{pmatrix} E_1^*(p, p_1) + y - [\mathcal{F}'(\bar{y})^* p]'(y - \bar{y}) \\ \alpha u - B^* p \\ y_t - \mathcal{A}y - Bu - \mathcal{F}'(\bar{y})(y - \bar{y}) - \mathcal{F}(\bar{y}) \\ y(0) - \bar{y}_0 \end{pmatrix} + \begin{pmatrix} 0 \\ \partial \mathbf{I}_{U_{ad}}(u) \\ 0 \\ 0 \end{pmatrix}, \quad (3.13)$$

and

$$\underline{\mathcal{T}} \begin{pmatrix} y \\ \underline{u} \\ \underline{p} \\ \underline{p}(0) \end{pmatrix} = \begin{pmatrix} -\underline{p}_t - \mathcal{A}^* \underline{p} - \mathcal{F}'(\bar{y})^* \underline{p} + y - [\mathcal{F}'(\bar{y})^* \bar{p}]'(y - \bar{y}) \\ \alpha \underline{u} - B^* \underline{p} \\ y_t - \mathcal{A}y - B\underline{u} - \mathcal{F}'(\bar{y})(y - \bar{y}) - \mathcal{F}(\bar{y}) \\ y(0) - \bar{y}_0 \end{pmatrix} + \begin{pmatrix} 0 \\ \partial \mathbf{I}_{U_{ad}}(\underline{u}) \\ 0 \\ 0 \end{pmatrix}. \quad (3.14)$$

where

$$\underline{\partial \mathbf{I}_{U_{ad}}}(\underline{u}) = \{\tilde{u} \in U \cap C(I; \mathcal{U}) : (\tilde{u}(t), v - \underline{u}(t))_{\mathcal{U}} \leq 0, \forall t \in I, v \in B_\eta(0)\}, \quad (3.15)$$

with $B_\eta(0) = \{v \in \mathcal{U}; \|v\|_{\mathcal{U}} \leq \eta\}$. In (3.14), we underline the elements which are taken from different domains when compared to (3.13). The range of the first two coordinates of $\underline{\mathcal{T}}$ is smaller than that of \mathcal{T} . Accordingly we can make use of (3.9) when moving from the first row of (3.13) to the first row of (3.14).

For convenience for the subsequent work we recall that the strong regularity condition introduced below (2.14) requires us to find neighborhoods of 0 and $(\bar{y}, \bar{u}, \bar{p}, \bar{p}(0))$ of the form $\hat{V} \subset L^2(I; V^*) \times (U \cap C(\bar{I}; \mathcal{U})) \times L^2(I; V^*) \times Y$ and $\hat{U} \subset W_\infty \times (U \cap C(\bar{I}; \mathcal{U})) \times W_\infty$, such that for all $\beta = (\beta_1, \beta_2, \beta_3, \beta_4) \in \hat{V}$ the equation

$$\underline{\mathcal{T}}(y, u, \underline{p}, \underline{p}(0))^T = (\beta_1, \beta_2, \beta_3, \beta_4)^T \quad (3.16)$$

admits a unique solution $(y, \underline{u}, \underline{p}, \underline{p}(0)) \in \hat{U}$ depending Lipschitz-continuously on β .

Remark 3.3. We observe that as a consequence of (A3) and Remark 3.2 the operator $\underline{\mathcal{T}}$ is continuous.

Subsequently we shall frequently refrain from the underline-notation since the meaning should be clear from the context.

3.3 Main Theorems.

In this subsection, we present the main theorems of this paper. The first theorem asserts local continuous differentiability of the value function \mathcal{V} w.r.t. y_0 , with y_0 small enough. The second theorem establishes that \mathcal{V} satisfies the HJB equation in the classical sense. The proof of the first theorem is based on Theorem 2.1. It will be given in Section 4 below. For this purpose it will be shown that assumptions A imply (H1)-(H7). Moreover we need to assert the underlying assumption that problem (P) is well-posed. This will lead to a smallness assumption on the initial states y_0 . Consequently it would suffice to assume that (A3) and (A4) only hold locally in the neighborhood of the origin. Concerning (A5) observe that it is not implied by (A3). It is vacuously satisfied for $\bar{y} = 0$, which is the case for $y_0 = 0$, since then $\mathcal{F}'(0) = 0$, see (3.3).

We invoke Theorem 2.1 to assert the Lipschitz continuity of the state, the adjoint state, and the control with respect to the initial condition $y_0 \in Y$ in the neighborhood of a locally optimal solution (\bar{y}, \bar{u}) corresponding to a sufficiently small reference initial state \bar{y}_0 . This will imply the differentiability of the value function associated to local minima. We shall refer to the value function associated to local minima as 'local value function'.

Theorem 3.1. *Let the assumptions (A) hold. Then associated to each local solution $(\bar{y}(y_0), \bar{u}(y_0))$ of (P) there exists a neighborhood of $U(y_0)$ such that the local value function $\mathcal{V} : U(y_0) \subset Y \rightarrow \mathbb{R}$ is continuously differentiable, provided that y_0 is sufficiently close to the origin in Y .*

To obtain a HJB equation we require additionally that $t \rightarrow (\mathcal{F}(\bar{y}))(t)$ is continuous with values in Y for global solutions (\bar{y}, \bar{u}) to (P), with $y_0 \in \mathcal{D}(\mathcal{A})$. In view of the fact that for $y_0 \in V$ we can typically expect that the solutions of semilinear parabolic equations satisfy $y \in L^2(I; \mathcal{D}(\mathcal{A})) \cap W^{1,2}(I; Y) \subset C([0, \infty), V)$ this is not a restrictive assumption beyond that what is already assumed in (A3).

Theorem 3.2. *Let the assumptions (A) hold, and let $(\bar{y}(y_0), \bar{u}(y_0))$ denote a global solution of (P), for $y_0 \in \mathcal{D}(\mathcal{A})$ with sufficiently small norm in Y . Assume that there exists $T_{y_0} > 0$ such that $\mathcal{F}(\bar{y}) \in C([0, T_{y_0}); Y)$. Then the following Hamilton-Jacobi-Bellman equation holds at y_0 :*

$$\mathcal{V}'(y)(\mathcal{A}y + \mathcal{F}(y)) + \frac{1}{2} \|y\|_Y^2 + \frac{\alpha}{2} \left\| \mathbb{P}_{\mathcal{U}_{ad}} \left(-\frac{1}{\alpha} B^* \mathcal{V}'(y) \right) \right\|_Y^2 + \left\langle B^* \mathcal{V}'(y), \mathbb{P}_{\mathcal{U}_{ad}} \left(-\frac{1}{\alpha} B^* \mathcal{V}'(y) \right) \right\rangle_Y = 0. \quad (3.17)$$

Moreover the optimal feedback law is given by

$$\bar{u}(0) = \mathbb{P}_{\mathcal{U}_{ad}} \left(-\frac{1}{\alpha} B^* \mathcal{V}'(\bar{y}(0)) \right). \quad (3.18)$$

The condition on the smallness of y_0 will be discussed in Remark 4.2 below. Roughly it involves well-posedness of the optimality system and second order sufficient optimality at local solutions. A more detailed, respectively stronger statement of Theorem 3.1 and Theorem 3.2, will be given in Theorem 4.1 and Theorem 5.1 below. The regularity assumptions $\mathcal{F}(\bar{y}) \in C([0, T_{y_0}); Y)$ of Theorem 3.2 will be addressed in Section 6.

4 Proof of Theorem 3.1.

In this section we give the proof for Theorem 3.1. Many of the technical difficulties arise from the fact that we are working with an infinite horizon optimal control problem. In this respect we can profit from techniques which were developed in [BKP3], which, however, do not include the case of constraints on the norm. Throughout we assume that assumptions (A1) - (A4) hold.

4.1 Well-posedness of problem (P).

Here we prove well-posedness for (P) with small initial data. First, we recall two consequences of the assumption that \mathcal{A} is the generator of an analytic semigroup.

Consequence 1. *Since \mathcal{A} generates a strongly continuous analytic semigroup on Y , there exist $\rho \geq 0$ and $\theta > 0$ such that*

$$\langle (\rho I - \mathcal{A})v, v \rangle_{V^*, V} \geq \theta \|v\|_V^2$$

See [BPDM, Part II, Chapter 1, p 115], [Paz, Theorem 4.2, p14].

Consequence 2. For all $y_0 \in Y, f \in L^2(0, T; V^*)$, and $T > 0$, there exists a unique solution $y \in W(0, T)$ to

$$\dot{y} = \mathcal{A}y + f, \quad y(0) = y_0. \quad (4.1)$$

Furthermore, y satisfies

$$\|y\|_{W(0, T)} \leq c(T) \left(\|y_0\|_Y + \|f\|_{L^2(0, T; V^*)} \right) \quad (4.2)$$

for a continuous function c . Assuming that $y \in L^2(0, \infty; Y)$, consider the equation

$$\dot{y} = \underbrace{(\mathcal{A} - \rho I)}_{\mathcal{A}_\rho} y + \underbrace{\rho y + f}_{f_\rho}, \quad y(0) = y_0,$$

where $f_\rho \in L^2(I; V^*)$. Then the operator \mathcal{A}_ρ generates a strongly continuous analytic semigroup on Y which is exponentially stable, see [BPD M , p 115, Theorem II.1.2.12]. It follows that $y \in W_\infty$, that there exists M_ρ such that

$$\|y\|_{W_\infty} \leq M_\rho \left(\|y_0\|_Y + \|f_\rho\|_{L^2(I; V^*)} \right), \quad (4.3)$$

and that y is the unique solution to (4.1) in W_∞ , see [BKP3, Section 2.2].

Lemma 4.1. There exists a constant $C > 0$, such that for all $\delta < (0, 1]$ and for all y_1 and y_2 in W_∞ with $\|y_1\|_{W_\infty} \leq \delta$ and $\|y_2\|_{W_\infty} \leq \delta$, it holds that

$$\|\mathcal{F}(y_1) - \mathcal{F}(y_2)\|_{L^2(I; V^*)} \leq \delta C \|y_1 - y_2\|_{W_\infty}. \quad (4.4)$$

Proof. Let y_1, y_2 be as in the statement of the lemma. Using (A3) and Remark 3.1 we obtain the estimate

$$\begin{aligned} \|\mathcal{F}(y_1) - \mathcal{F}(y_2)\|_{L^2(I; V^*)} &\leq \int_0^1 \|\mathcal{F}'(y_1 + t(y_2 - y_1)) - \mathcal{F}'(0)\|_{\mathcal{L}(W_\infty, L^2(I; V^*))} dt \|y_2 - y_1\|_{W_\infty} \\ &\leq \int_0^1 \int_0^1 \|\mathcal{F}''(s(y_1 + t(y_2 - y_1)))(ty_2 + (1-t)y_1)\|_{\mathcal{L}(W_\infty, L^2(I; V^*))} ds dt \|y_2 - y_1\|_{W_\infty}. \end{aligned}$$

Now the claim follows by assumption (A3). □

Lemma 4.2. Let \mathcal{A}_s be the generator of an exponentially stable analytic semigroup $e^{\mathcal{A}_s t}$ on Y . Let C denote the constant from Lemma 4.1. Then there exists a constant M_s such that for all $y_0 \in Y$ and $f \in L^2(I; V^*)$ with

$$\tilde{\gamma} = \|y_0\|_Y + \|f\|_{L^2(I; V^*)} \leq \frac{1}{4CM_s^2}$$

the system

$$y_t = \mathcal{A}_s y + \mathcal{F}(y) + f, \quad y(0) = y_0 \quad (4.5)$$

has a unique solution $y \in W_\infty$, which satisfies

$$\|y\|_{W_\infty} \leq 2M_s \tilde{\gamma}.$$

With Lemma 4.1 holding, this lemma can be verified in the same manner as [BKP3, Lemma 5, p 6]. In the following corollary we shall use Lemma 4.2 with $\mathcal{A}_s = \mathcal{A} - BK$, and the constant corresponding to M_s will be denoted by M_K . Further $\|\mathcal{I}\|$ denotes the norm of the embedding constant of W_∞ into $C(I; Y)$, $\|i\|$ is the norm of the embedding V into Y , and we recall the constant η from (3.1).

Corollary 4.3. *For all $y_0 \in Y$ with*

$$\|y_0\|_Y \leq \min \left\{ \frac{1}{4CM_K^2}, \frac{\eta}{2M_K \|K\|_{\mathcal{L}(Y)} \|\mathcal{I}\|} \right\}$$

there exists a control $u \in U_{ad}$ such that the system

$$y_t = \mathcal{A}y + \mathcal{F}(y) + Bu, \quad y(0) = y_0 \quad (4.6)$$

has a unique solution $y \in W_\infty$ satisfying

$$\|y\|_{W_\infty} \leq 2M_K \|y_0\|_Y \quad \text{and} \quad \|u\|_U \leq \|K\|_{\mathcal{L}(Y, U)} \|\mathcal{I}\| \|y\|_{W_\infty} \leq 2M_K \|y_0\|_Y \|K\|_{\mathcal{L}(Y, U)} \|\mathcal{I}\|. \quad (4.7)$$

Proof. By Assumption (A2), there exists K such that $\mathcal{A} - BK$ generates an exponentially stable analytic semigroup on Y . Taking $u = -Ky$, equation (4.6) becomes

$$y_t = (\mathcal{A} - BK)y + \mathcal{F}(y), \quad y(0) = y_0. \quad (4.8)$$

Then by Lemma 4.2 with $\tilde{\gamma} = \|y_0\|_Y$ there exists M_K such that (4.8) has a solution $y \in W_\infty$ satisfying

$$\|y\|_{W_\infty} \leq 2M_K \|y_0\|_Y,$$

and thus the first inequality in (4.7) holds. For every $t \in I$ we have

$$\|u\|_U = \|Ky\|_U \leq \|K\|_{\mathcal{L}(Y, U)} \|y\|_Y \leq \|K\|_{\mathcal{L}(Y, U)} \|\mathcal{I}\| \|y\|_{W_\infty} \leq 2M_K \|y_0\|_Y \|K\|_{\mathcal{L}(Y, U)} \|\mathcal{I}\|, \quad (4.9)$$

and thus the second inequality in (4.7) holds. We still need to assert that $u \in U_{ad}$. This follows from the second smallness condition on $\|y_0\|_Y$ and (4.9). \square

Remark 4.1. In the above proof stabilization was achieved by the feedback control $u = -Ky$. For this u to be admissible it is needed that U_{ad} has nonempty interior. The upper bound η could be allowed to be time dependent as long as it satisfies $\inf_{t \geq 0} |\eta(t)| > 0$.

Corollary 4.4. *Let $y_0 \in Y$ and let $u \in U_{ad}$ be such that the system*

$$y_t = \mathcal{A}y + \mathcal{F}(y) + Bu, \quad y(0) = y_0 \quad (4.10)$$

has a unique solution $y \in L^2(I; Y)$. If

$$\gamma := \|y_0\|_Y + \|\rho y + Bu\|_{L^2(I; V^*)} \leq \min \left\{ \frac{1}{4CM_\rho^2}, \frac{\eta}{2M_\rho \|K\|_{\mathcal{L}(Y)} \|\mathcal{I}\|} \right\},$$

then $y \in W_\infty$ and it holds that

$$\|y\|_{W_\infty} \leq 2M_\rho \gamma.$$

Proof. Since $y \in L^2(I; Y)$, we can apply Lemma 4.2 to the equivalent system

$$y_t = (\mathcal{A} - \rho I)y + \mathcal{F}(y) + \tilde{f},$$

where $\tilde{f} = \rho y + Bu$. This proves the assertion. \square

Lemma 4.5. *There exists $\delta_1 > 0$ such that for all $y_0 \in B_Y(\delta_1)$, problem (P) possesses a solution $(\bar{y}, \bar{u}) \in W_\infty \times U_{ad}$. Moreover, there exists a constant $M > 0$ independent of y_0 such that*

$$\max \{ \|\bar{y}\|_{W_\infty}, \|\bar{u}\|_U \} \leq M \|y_0\|_Y \quad (4.11)$$

Proof. The proof of this lemma follows with analogous argumentation as provided in [BKP3, Lemma 8]. Let us choose, $\delta_1 \leq \min \left\{ \frac{1}{4CM_K^2}, \frac{\eta}{2M_K \|K\|_{\mathcal{L}(Y, U)}} \right\}$, where C as in Lemma 4.1 and M_K denotes the constant from the Corollary 4.3. We obtain that for each $y_0 \in B_Y(\delta_1)$, there exists a control $u \in U_{ad}$ with associated state y satisfying

$$\max \{ \|u\|_U, \|y\|_{W_\infty} \} \leq \tilde{M} \|y_0\|_Y, \quad (4.12)$$

where $\tilde{M} = 2M_K \max(1, \|i\| \|K\|_{\mathcal{L}(Y, U)})$. We can thus consider a minimizing sequence $(y_n, u_n)_{n \in \mathbb{N}} \in W_\infty \times U_{ad}$ with $J(y_n, u_n) \leq \frac{1}{2} M^2 \|y_0\|_Y^2 (1 + \alpha)$. For all $n \in \mathbb{N}$ that

$$\|y_n\|_{L^2(I; Y)} \leq \tilde{M} \|y_0\|_Y \sqrt{1 + \alpha} \quad \text{and} \quad \|u_n\|_{L^2(I; U)} \leq \tilde{M} \|y_0\|_Y \sqrt{\frac{1 + \alpha}{\alpha}}. \quad (4.13)$$

We set $\eta(\alpha, \tilde{M}) = \left[1 + \tilde{M} \|i\| \sqrt{1 + \alpha} \left(\rho + \frac{\|B\|_{\mathcal{L}(U, Y)}}{\sqrt{\alpha}} \right) \right]$. Then we have $\|y_0\| + \|\rho y_n + Bu_n\|_{L^2(I; V^*)} \leq \eta(\alpha, \tilde{M}) \|y_0\|_Y$. After further reduction of δ_1 , we obtain with M_ρ from Corollary 4.4:

$$\|y_0\| + \|\rho y_n + Bu_n\|_{L^2(I; V^*)} \leq \frac{1}{4CM_\rho^2}.$$

It follows from this corollary that the sequence $\{y_n\}_{n \in \mathbb{N}}$ is bounded in W_∞ with

$$\sup_{n \in \mathbb{N}} \|y_n\|_{W_\infty} \leq 2M_\rho (1 + \eta(\alpha, \tilde{M})) \|y_0\|_Y. \quad (4.14)$$

Extracting if necessary a subsequence, there exists $(\bar{y}, \bar{u}) \in W_\infty \times U$ such that $(y_n, u_n) \rightharpoonup (\bar{y}, \bar{u}) \in W_\infty \times U$, and (\bar{y}, \bar{u}) satisfies (4.12).

Let us prove that (\bar{y}, \bar{u}) is feasible and optimal. Since U_{ad} is weakly sequentially closed and $u_n \in U_{ad}$, we find $\bar{u} \in U_{ad}$. For each fixed $T > 0$ and arbitrary $z \in L^\infty(0; T; \mathcal{H}) \subset L^2(0, T; V)$, see (A4), we have for all $n \in \mathbb{N}$ that

$$\int_0^T \langle \dot{y}_n(t), z(t) \rangle_{V^*, V} dt = \int_0^T \langle \mathcal{A}y_n(t) + \mathcal{F}(y_n(t)) + Bu_n(t), z(t) \rangle_{V^*, V} dt. \quad (4.15)$$

Since $\dot{y}_n \rightharpoonup \dot{y}$ in $L^2(0, T; V^*)$, we can pass to the limit in the l.h.s. of the above equality. Moreover, since $\mathcal{A}y_n \rightharpoonup \mathcal{A}\bar{y}$ in $L^2(0, T; V^*)$,

$$\int_0^T \langle \mathcal{A}y_n(t), z(t) \rangle_{V^*, V} dt \xrightarrow{n \rightarrow \infty} \int_0^T \langle \mathcal{A}\bar{y}(t), z(t) \rangle_{V^*, V} dt.$$

Analogously, we obtain that

$$\int_0^T \langle Bu_n(t), z(t) \rangle_{V^*, V} dt \xrightarrow{n \rightarrow \infty} \int_0^T \langle B\bar{u}(t), z(t) \rangle_{V^*, V} dt.$$

If moreover $z \in L^\infty(0, T; \mathcal{H}) \subset L^2(0, T; V)$, we use (A4) to assert

$$\int_0^T \langle \mathcal{F}(y_n(t)) - \mathcal{F}(\bar{y}(t)), z(t) \rangle_{V^*, V} dt = \int_0^T \langle \mathcal{F}(y_n(t)) - \mathcal{F}(\bar{y}(t)), z(t) \rangle_{\mathcal{H}^*, \mathcal{H}} dt \xrightarrow{n \rightarrow \infty} 0.$$

Thus we have for all $z \in L^\infty(0, T; \mathcal{H})$

$$\int_0^T \langle \dot{y}(t) - \mathcal{A}y(t) - Bu(t), z(t) \rangle_{V^*, V} dt = \int_0^T \langle \mathcal{F}(y(t)), z(t) \rangle_{V^*, V} dt. \quad (4.16)$$

Since $\dot{y} - \mathcal{A}y - Bu \in L^2(0, T; V^*)$ and $L^\infty(0, T; \mathcal{H})$ is dense in $L^2(0, T; V)$ we conclude that (4.16) holds for all $z \in L^2(0, T; V)$ and $T > 0$. This yields $e(\bar{y}, \bar{u}) = (0, 0)$, and thus (\bar{y}, \bar{u}) is feasible. By weak lower semicontinuity of norms it follows that $J(\bar{y}, \bar{u}) \leq \liminf_{n \rightarrow \infty} J(\bar{y}_n, \bar{u}_n)$, which proves the optimality of (\bar{y}, \bar{u}) , and (4.11) follows from (4.13). \square

For the derivation of the optimality system for (P), we need the following lemma which is taken from [BKP1, Lemma 2.5].

Lemma 4.6. *Let $G \in \mathcal{L}(W_\infty, L^2(I; V^*))$ such that $\|G\| < \frac{1}{M_K}$, where $\|G\|$ denotes the operator norm of G . Then for all $f \in L^2(I; V^*)$ and $y_0 \in Y$, there exists a unique solution to the problem:*

$$y_t = (\mathcal{A} - BK)y(t) + (Gy)(t) + f(t), \quad y = y_0.$$

Moreover,

$$\|y\|_{W_\infty} \leq \frac{M_K}{1 - M_K \|G\|} \left(\|f\|_{L^2(I; V^*)} + \|y_0\|_Y \right).$$

We close this section by deriving the optimality conditions for (P).

Proposition 4.1. *Let the assumptions (A1) - (A4) hold. Then there exists $\delta_2 \in (0, \delta_1]$ such that each local solution (\bar{y}, \bar{u}) with $y_0 \in B_Y(\delta_2)$ is a regular point, i.e. (2.3) is satisfied, and there exists an adjoint state $(\bar{p}, \bar{p}_1) \in L^2(I; V) \times Y$ satisfying*

$$\langle v_t - \mathcal{A}v - \mathcal{F}'(\bar{y})v, \bar{p} \rangle_{L^2(I; V^*), L^2(I; V)} + (v(0), \bar{p}_1)_Y + (\bar{y}, \bar{p})_{L^2(I; V)} = 0, \quad \text{for all } v \in W_\infty, \quad (4.17)$$

$$\langle \alpha \bar{u} - B^* \bar{p}, u - \bar{u} \rangle_U \geq 0, \quad \text{for all } u \in U_{ad}. \quad (4.18)$$

If the assumption (A5) is satisfied, then

$$-\bar{p}_t - \mathcal{A}^* \bar{p} - \mathcal{F}'(\bar{y})^* \bar{p} = -\bar{y} \quad \text{in } L^2(I; V^*),$$

and hence $\bar{p} \in W_\infty$ and

$$\lim_{t \rightarrow \infty} \bar{p}(t) = 0. \quad (4.19)$$

Moreover, there exists $\widetilde{M} > 0$, independent of $y_0 \in B_Y(\delta_2)$, such that

$$\|\bar{p}\|_{W_\infty} \leq \widetilde{M} \|y_0\|_Y, \quad \text{and } u \in C(\bar{I}, \mathcal{U}). \quad (4.20)$$

Proof. To verify the regular point condition, we evaluate e defined in (3.6) at (\bar{y}, \bar{u}, y_0) . To check the claim on the range of $e'(\bar{y}, \bar{u}, y_0)$ we consider for arbitrary $(r, s) \in L^2(I, V^*) \times Y$ the equation

$$z_t - \mathcal{A}z - \mathcal{F}'(\bar{y})z - B(w - \bar{u}) = r, \quad z(0) = s, \quad (4.21)$$

for unknowns $(z, w) \in W_\infty \times U_{ad}$. By taking $w = -Kz \in U$ we obtain

$$z_t - (\mathcal{A} - BK)z - \mathcal{F}'(\bar{y})z + B\bar{u} = r, \quad z(0) = s.$$

We apply Lemma 4.6 to this equation with $G = -\mathcal{F}'(\bar{y})$ and $f = r - B\bar{u}$. By Lemma 4.5 and (3.3) in Remark 3.1 there exists $\delta_2 \in (0, \delta_1]$ such that $\|\mathcal{F}'(\bar{y})\|_{\mathcal{L}(W_\infty, L^2(I; V^*))} \leq \frac{1}{2}M_K$. Consequently by Lemma 4.6 there exists \widetilde{M} such that

$$\begin{aligned} \|z\|_{W_\infty} &\leq \widetilde{M}(\|r\|_{L^2(I; V^*)} + \|s\|_Y + \|B\|_{\mathcal{L}(U, Y)} \|\bar{u}\|_U) \\ &\leq \widetilde{M}(\|r\|_{L^2(I; V^*)} + \|s\|_Y + \|B\|_{\mathcal{L}(U, Y)} M \|y_0\|_Y), \end{aligned} \quad (4.22)$$

with M as in (4.11). We shall need to check whether $w = -Kz$ is feasible, which will be the case if $w(t) \leq \eta$ for a.e. $t \in I$. Indeed we have

$$\|w(t)\|_Y \leq \|K\|_{\mathcal{L}(Y, U)} \|z(t)\|_Y \leq \|K\|_{\mathcal{L}(Y, U)} \|\mathcal{I}\| \widetilde{M}(\|r\|_{L^2(I; V^*)} + \|s\|_Y + \|B\|_{\mathcal{L}(U, Y)} M \|y_0\|_Y).$$

Consequently, possibly after further reducing δ_2 , and choosing $\tilde{\delta} > 0$ sufficiently small we have

$$\|w\|_{L^\infty(I; Y)} \leq \eta \text{ for all } y_0 \in B_Y(\delta_2) \text{ and all } (r, s) \text{ satisfying } \|(r, s)\|_{L^2(I; V^*) \times Y} \leq \tilde{\delta}. \quad (4.23)$$

Consequently the regular point condition is satisfied. Hence there exists a multiplier $\lambda = (p, p_1) \in L^2(I; V) \times Y$ satisfying,

$$\langle \mathcal{L}_y(\bar{y}, \bar{u}, y_0, \bar{p}, \bar{p}_1), v \rangle_{L^2(I; V^*), L^2(I; V)} = 0, \quad \langle \mathcal{L}_u(\bar{y}, \bar{u}, y_0, \bar{p}, \bar{p}_1), u - \bar{u} \rangle_U \geq 0, \quad \forall u \in U_{ad} \quad (4.24)$$

where

$$\mathcal{L}(y, u, y_0, p, p_1) = J(y, u) + \int_0^\infty \langle p, y_t - \mathcal{A}y - \mathcal{F}(y) - Bu \rangle_{V, V^*} dt + \langle p_1, y(0) - y_0 \rangle_Y.$$

This implies that (4.17) holds.

Now, if we impose the additional assumption (A5), we have $\mathcal{F}'(\bar{y})^* \bar{p} \in L^2(I; V^*)$. Thus $-\mathcal{A}^* \bar{p} - \mathcal{F}'(\bar{y})^* \bar{p} + \bar{y} \in L^2(I; V^*)$ and the previous identity implies that $\bar{p} \in W_\infty$. Thus we derive

$$-\bar{p}_t - \mathcal{A}^* \bar{p} - \mathcal{F}'(\bar{y})^* \bar{p} = -\bar{y} \text{ in } L^2(I; V^*) \text{ and } \lim_{t \rightarrow \infty} \bar{p}(t) = 0,$$

and (4.17)-(4.19). Testing the first identity in (4.24) with $v \in L^2(I; V)$ we also have $\bar{p}_1 = \bar{p}(0)$, which is well-defined since $\bar{p} \in W_\infty \subset C(I; Y)$. The second identity in (4.24) gives (4.18). It remains to estimate $\bar{p} \in W_\infty$.

Let $r \in L^2(I; V^*)$ with $\|r\|_{L^2(I; V^*)} \leq \tilde{\delta}$ and consider

$$z_t - \mathcal{A}z - \mathcal{F}'(\bar{y})z - B(w - \bar{u}) = -r, \quad z(0) = 0. \quad (4.25)$$

Arguing as in (4.21)-(4.22) there exists a solution to (4.25) with $w = -Kz$ such that

$$\|z\|_{W_\infty} \leq \widetilde{M}(\widetilde{\delta} + \|B\|_{\mathcal{L}(\mathcal{U}, Y)} M \|y_0\|_Y) \leq \widetilde{M}(\widetilde{\delta} + \|B\|_{\mathcal{L}(\mathcal{U}, Y)} M \delta_2) =: C_1. \quad (4.26)$$

From (4.23) we have that $\|w\|_{L^\infty(I, \mathcal{U})} \leq \eta$. Let us now observe that

$$\begin{aligned} \langle \bar{p}, r \rangle_{L^2(I, V), L^2(I, V^*)} &= \langle \bar{p}, -z_t + \mathcal{A}z + \mathcal{F}'(\bar{y})z \rangle_{L^2(I, V), L^2(I, V^*)} + \langle \bar{p}, B(w - \bar{u}) \rangle_{L^2(I; Y)} \\ &= \langle \bar{p}_t + \mathcal{A}^* \bar{p} + \mathcal{F}'(\bar{y})^* \bar{p}, z \rangle + \langle B^* \bar{p}, w - \bar{u} \rangle_U, \end{aligned}$$

where we have used that $z(0) = 0$ and $\lim_{t \rightarrow \infty} \bar{p}(t) = 0$, since $\bar{p} \in W_\infty$. We next estimate using (4.17), (4.19) and (4.26)

$$\langle \bar{p}, r \rangle_{L^2(I, V), L^2(I, V^*)} \leq \|\bar{y}\|_{L^2(I, V^*)} \|z\|_{L^2(I, V)} + \alpha \langle \bar{u}, w - \bar{u} \rangle_U \leq \left(\|\bar{y}\|_{L^2(I, V^*)} + \alpha \|\bar{u}\|_U \right) (C_1 + \eta + \|\bar{u}\|_U).$$

By (4.11), this implies the existence of a constant C_2 such that

$$\sup_{\|r\|_{L^2(I, V^*)} \leq \widetilde{\delta}} \langle \bar{p}, r \rangle_{L^2(I, V), L^2(I, V^*)} \leq C_2 \|y_0\|_Y$$

and thus

$$\|\bar{p}\|_{L^2(I, V)} \leq \frac{C_2}{\widetilde{\delta}} \|y_0\|_Y, \quad \text{for all } y_0 \in B_Y(\delta_2). \quad (4.27)$$

Now we estimate, again using (A5)

$$\|\bar{p}_t\|_{L^2(I; V^*)} \leq \|\mathcal{A}^* \bar{p} + \mathcal{F}'(\bar{y})^* \bar{p} - \bar{y}\|_{L^2(I; V^*)} \leq C_3 \|\bar{p}\|_{L^2(I, V)} + C_4 \|\bar{p}\|_{L^2(I, V)} + \|\bar{y}\|_{L^2(I; V^*)}.$$

By (4.11) and (4.27) we obtain $\|\bar{p}_t\|_{L^2(I; V^*)} \leq C_5 \|y_0\|_Y$. Combining this estimate with (4.27) yields (4.20). Finally, by (4.18) we find $\bar{u}(t) = \mathbb{P}_{\mathcal{U}_{ad}} \left(\frac{1}{\alpha} B^* \bar{p}(t) \right)$. Since $\bar{p} \in C(\bar{I}; Y)$ and $B^* \in \mathcal{L}(Y, U)$ this implies that $u \in C(\bar{I}; \mathcal{U})$. \square

4.2 Verification of (H1)-(H6).

In this section we specialize the previously proved abstract results in Section 2 to the semilinear parabolic setting. We start with the following lemma which shows that assumptions A imply (H1)-(H6).

Lemma 4.7. *Consider problem (P) with assumptions (A1)-(A4) holding. Then (H1)-(H4), (H6) are satisfied for (P) uniformly for all $y_0 \in B_Y(\delta_2)$ for some $\delta_2 \in (0, \delta_2]$. If moreover, (A5) holds, then (H5) holds as well.*

Proof. Throughout $y_0 \in B_Y(\delta_2)$, (\bar{y}, \bar{u}) denotes a local solution to (P), and $(p, p_1) \in L^2(I; V^*) \times Y$ the associated Lagrange multiplier.

- (i) Verification of (H1): The initial condition y_0 is our nominal reference parameter q . Lemma 4.5 guarantees the existence of a local solution $(\bar{y}, \bar{u}) \sim x_0$ to (P) $\sim (P_{q_0})$. Clearly f defined in (3.5) satisfies the required regularity assumptions. Moreover e satisfies the regularity assumptions as a consequence of (A3).

(ii) Verification of **(H2)**: Proposition 4.1 implies that (\bar{y}, \bar{u}) is a regular point.

(iii) Verification of **(H3)**: The second derivative of e is given by

$$e''(\bar{y}, \bar{u}, y_0)((v_1, w_1), (v_2, w_2)) = \begin{pmatrix} \mathcal{F}''(\bar{y})(v_1, v_2) \\ 0 \end{pmatrix}, \quad \forall v_1, v_2 \in W_\infty, \forall w_1, w_2 \in U. \quad (4.28)$$

For the second derivative of \mathcal{L} w.r.t. (y, u) , we find

$$\begin{aligned} \mathcal{L}''(\bar{y}, \bar{u}, y_0, \bar{p}, \bar{p}_1)((v_1, w_1), (v_2, w_2)) &= \int_0^\infty (v_1, v_2)_Y dt + \alpha \int_0^\infty (w_1, w_2)_Y dt \\ &\quad + \int_0^\infty \langle \bar{p}, \mathcal{F}''(\bar{y})(v_1, v_2) \rangle_{V, V^*} dt. \end{aligned} \quad (4.29)$$

By **(A3)** for \mathcal{F}'' and Lemma 4.5, there exists \widetilde{M}_1 such that

$$\int_0^\infty \langle \bar{p}, \mathcal{F}''(\bar{y})(v, v) \rangle_{V, V^*} dt \leq \widetilde{M}_1 \|\bar{p}\|_{L^2(I; V)} \|v\|_{W_\infty}^2, \quad \forall v \in W_\infty, \quad (4.30)$$

for each solution (\bar{y}, \bar{u}) of **(P)** with $y_0 \in B_Y(\delta_2)$. Then we obtain

$$\mathcal{L}''(\bar{y}, \bar{u}, y_0, \bar{p}, \bar{p}_1)((v, w), (v, w)) \geq \int_0^\infty \|v\|_Y^2 dt + \alpha \int_0^\infty \|w\|_U^2 dt - \widetilde{M}_1 \|\bar{p}\|_{L^2(I; V)} \|v\|_{W_\infty}^2. \quad (4.31)$$

Now let $0 \neq (v, w) \in \ker E \subset W_\infty \times U_{ad}$, where E as defined in (3.7) is evaluated at (\bar{y}, \bar{u}) . Then,

$$v_t - \mathcal{A}v - \mathcal{F}'(\bar{y})v - Bw = 0, \quad v(0) = 0.$$

Next choose $\rho > 0$, such that the semigroup generated by $(\mathcal{A} - \rho I)$ is exponentially stable. This is possible due to **(A1)**. We equivalently write the system in the previous equation as,

$$v_t - (\mathcal{A} - \rho I)v - \mathcal{F}'(\bar{y})v - \rho v - Bw = 0, \quad v(0) = 0.$$

Now, we invoke Lemma 4.6 with $\mathcal{A} - BK$ replaced by $\mathcal{A} - \rho I$, $G = \mathcal{F}'(\bar{y})$, and $f(t) = \rho v(t) + Bw(t)$, and the role of the constant M_K will now be assumed by a parameter M_ρ . By selecting $\widetilde{\delta}_2 \in (0, \delta_2]$ such that $\|\bar{y}\|_{W_\infty}$ sufficiently small, we can guarantee that $\|\mathcal{F}'(\bar{y})\|_{\mathcal{L}(W_\infty; L^2(I; V^*))} \leq 1/2M_\rho$, see (4.11) and (3.3) in Remark 3.1. Then the following estimate holds,

$$\|v\|_{W_\infty} \leq 2M_\rho \|v + Bw\|_{L^2(I; V^*)}.$$

This implies that

$$\|v\|_{W_\infty}^2 \leq \widetilde{M}_2 (\|v\|_{L^2(I; Y)}^2 + \|w\|_{L^2(I; Y)}^2). \quad (4.32)$$

for a constant \widetilde{M}_2 depending on $M_\rho, \|B\|$, and the embedding of Y into V^* . These preliminaries allow the following lower bound on \mathcal{L}'' :

$$\begin{aligned} \mathcal{L}''(\bar{y}, \bar{u}, y_0, \bar{p}, \bar{p}_1)((v, w), (v, w)) &\geq \int_0^\infty \|v\|_Y^2 dt + \alpha \int_0^\infty \|w\|_Y^2 dt - \widetilde{M}_1 \|\bar{p}\|_{L^2(I; V)} \|v\|_{W_\infty}^2 \\ \text{by (4.32)} &\geq \int_0^\infty \|v\|_Y^2 + \alpha \int_0^\infty \|w\|_Y^2 - \widetilde{M}_1 \widetilde{M}_2 \|\bar{p}\|_{L^2(I; V)} \left[\|v\|_{L^2(I; Y)}^2 + \|w\|_{L^2(I; Y)}^2 \right] \end{aligned}$$

$$\begin{aligned}
&= \left(1 - \widetilde{M}_1 \widetilde{M}_2 \|\bar{p}\|_{L^2(I;V)}\right) \|v\|_{L^2(I;Y)}^2 + \left(\alpha - \widetilde{M}_1 \widetilde{M}_2 \|\bar{p}\|_{L^2(I;V)}\right) \|w\|_{L^2(I;Y)}^2 \\
&\geq \tilde{\gamma} \left[\|v\|_{L^2(I;Y)}^2 + \|w\|_{L^2(I;Y)}^2 \right]
\end{aligned} \tag{4.33}$$

where $\tilde{\gamma} = \min \left\{ 1 - \widetilde{M}_1 \widetilde{M}_2 \|\bar{p}\|_{L^2(I;V)}, \alpha - \widetilde{M}_1 \widetilde{M}_2 \|\bar{p}\|_{L^2(I;V)} \right\}$. By possible further reduction of $\tilde{\delta}_2$ it can be guaranteed that $\tilde{\gamma} > 0$, see (4.27). Then by (4.32), we obtain,

$$\begin{aligned}
\mathcal{L}''(\bar{y}, \bar{u}, y_0, \bar{p}, \bar{p}_1)((v, w), (v, w)) &\geq \frac{\tilde{\gamma}}{2} \left[\|v\|_{L^2(I;Y)}^2 + \|w\|_{L^2(I;Y)}^2 \right] + \frac{\tilde{\gamma}}{2\widetilde{M}_2} \|v\|_{W_\infty}^2, \\
&\geq \frac{\tilde{\gamma}}{2\widetilde{M}_2} \|v\|_{W_\infty}^2 + \frac{\tilde{\gamma}}{2} \|w\|_{L^2(I;Y)}^2.
\end{aligned}$$

By selecting $\bar{\gamma} = \min \left\{ \frac{\tilde{\gamma}}{2\widetilde{M}_2}, \frac{\tilde{\gamma}}{2} \right\}$, we obtain the positive definiteness of \mathcal{L}'' , i.e.

$$\mathcal{L}''(\bar{y}, \bar{u}, y_0, \bar{p}, \bar{p}_1)((v, w), (v, w)) \geq \bar{\gamma} \|(v, w)\|_{W_\infty \times U}^2, \quad y_0 \in B_Y(\tilde{\delta}_2), \quad (v, w) \in \ker E. \tag{4.34}$$

Thus (H3) is satisfied.

(iv) Verification of (H4): It can easily be checked that $f'(y, u)$ can be extended to an element in $\underline{X}^* = L^2(I; V^*) \times (U \cap C(\bar{I}; \mathcal{U}))$ for each $(y, u) \in X = W_\infty \times U$. We refer to Remark 3.2 to show that the restriction of $e'(y, u, y_0)^*$ to \underline{W}^* satisfies $\underline{e}'(y, u, y_0)^* \in \mathcal{L}(\underline{W}^*, \underline{X}^*) = \mathcal{L}(W_\infty \times Y, L^2(I; V^*) \times (U \cap C(\bar{I}; \mathcal{U})))$.

(v) Verification of (H6): This is trivially satisfied.

Thus we have proved that assumptions (A1)-(A4) imply (H1)-(H4), and (H6) for all $y_0 \in B_Y(\tilde{\delta}_2)$.

(vi) Verification of (H5): Here we use (A5) and have $(p, p_1) = (p, p(0)) \in \widetilde{W}_\infty$. Observe that $\underline{\mathcal{L}}' : W_\infty \times (U \cap C(\bar{I}; \mathcal{U})) \times Y \times W_\infty \rightarrow L^2(I; V^*) \times (U \cap C(\bar{I}; \mathcal{U}))$ evaluated at $(\bar{y}, \bar{u}, y_0, p)$ is given by

$$\underline{\mathcal{L}}'(\bar{y}, \bar{u}, y_0, \bar{p}) = \begin{pmatrix} \bar{y} + \frac{e'(\bar{y}, \bar{u}, y_0)^* \bar{p}}{\alpha \bar{u} - B^* \bar{p}} \\ \alpha \bar{u} - B^* \bar{p} \end{pmatrix} = \begin{pmatrix} \bar{y} - \bar{p}_t - \mathcal{A}^* \bar{p} - \mathcal{F}'(\bar{y})^* \bar{p} \\ \alpha \bar{u} - B^* \bar{p} \end{pmatrix}.$$

Further for $(v, w) \in W_\infty \times U$ we have

$$(\underline{\mathcal{L}}')'(\bar{y}, \bar{u}, y_0, \bar{p})(v, w) = \begin{pmatrix} v - [\mathcal{F}'(\bar{y})^*]'(\bar{p}, v) \\ \alpha w \end{pmatrix} \in L^2(I; V^*) \times (U \cap C(\bar{I}; \mathcal{U})).$$

By (A3) and Remark 3.2 we have that $\underline{\mathcal{L}}'$ and $(\underline{\mathcal{L}}')'$ are continuous as mappings from $W_\infty \times (U \cap C(\bar{I}; \mathcal{U})) \times Y \times W_\infty$ to $L^2(I; V^*) \times (U \cap C(\bar{I}; \mathcal{U}))$, respectively to $\mathcal{L}(W_\infty \times (U \cap C(\bar{I}; \mathcal{U})); L^2(I; V^*) \times (U \cap C(\bar{I}; \mathcal{U})))$.

This proves the lemma. \square

Remark 4.2. Let us summarize our findings so far. There exists $\tilde{\delta}_2$ such that for each $y_0 \in B_Y(\tilde{\delta}_2)$ problem (P) possesses a solution $(\bar{y}, \bar{u}) \in W_\infty \times (U \cap C(\bar{I}; \mathcal{U}))$, with an adjoint $\bar{p} \in \widetilde{W}_\infty$. Further (A1)-(A5) imply (H1)-(H6) for (P) with $y_0 \in B_Y(\tilde{\delta}_2)$. As a consequence for each $y_0 \in B_Y(\tilde{\delta}_2)$ and each associated local solution (\bar{y}, \bar{u}) there exists a neighborhood \hat{V} of the origin in

$\mathcal{Y} := L^2(I; V^*) \times (U \cap C(\bar{I}; \mathcal{U})) \times L^2(I; V^*) \times Y$ such that for each $\beta \in \hat{V}$ there exists a unique solution $(y_{(\beta)}, u_{(\beta)}, p_{(\beta)}, p_{1(\beta)}) \in W_\infty \times U \times L^2(I; V) \times Y$ to $\mathcal{T}(y, u, p, p_1) = \beta$, see step (ii) of the proof of Theorem 2.1. – To verify the remaining assumption (H7) we need to argue that $(u_{(\beta)}, p_{(\beta)}) \in (U \cap C(\bar{I}; \mathcal{U})) \times W_\infty$ and that $\beta \mapsto (y_{(\beta)}, u_{(\beta)}, p_{(\beta)})$ is Lipschitz continuous from $\hat{V} \subset \mathcal{Y}$ to $W_\infty \times (U \cap C(\bar{I}; \mathcal{U})) \times W_\infty$.

Remark 4.3. Here we remark on the smallness assumption on y_0 expressed by δ_2 , respectively $\tilde{\delta}_2$. The condition $y_0 \in B_Y(\delta_2)$ guarantees the well-posedness of (P), existence and boundedness of adjoint states as expressed in Proposition 4.1. The additional condition $y_0 \in B_Y(\tilde{\delta}_2)$ implies that the second order optimality condition (H3) is satisfied, for each local solution associated to an initial condition $y_0 \in B_Y(\delta_2)$. In the following we formulate the results for all $y_0 \in B_Y(\tilde{\delta}_2)$. Alternatively we could narrow down the claims to neighborhoods of single local solutions (\bar{y}, \bar{u}) with $y_0 \in B_Y(\delta_2)$ and additionally assuming that the second order condition is satisfied at (\bar{y}, \bar{u}) . Concerning the second order condition itself, in some publications, see e.g. [Gri], it is required to hold only for elements $x = (y, u) \in \ker E$ and $u = u_1 - u_2$, with u_1, u_2 in U_{ad} . By a scaling argument it can easily be seen that this condition is equivalent to the one we use.

4.3 Verification of (H7) and Lipschitz stability of the linearized problem.

Throughout the remainder, we assume that (A1)-(A5) are satisfied and that $y_0 \in B_Y(\tilde{\delta}_2)$ so that Proposition 4.1 and Lemma 4.7 are applicable. In the following, the triple (y, u, p) refers to the solution $\mathcal{T}(y, u, p, p_1) = \beta$. Throughout without loss of generality, we also assume that \hat{V} is bounded.

Lemma 4.8. *Let assumptions (A) hold and let (\bar{y}, \bar{u}) , and \bar{p} denote a local solution and associated adjoint state to (P) corresponding to an initial datum $y_0 \in B_Y(\tilde{\delta}_2)$. Then the mapping $\beta \mapsto p_{(\beta)}$ is continuous from \hat{V} to W_∞ .*

Proof.

Step 1: For $\beta \in \hat{V}$, with \hat{V} as in Remark 4.2, let $(y_{(\beta)}, u_{(\beta)}, p_{(\beta)}, p_{1(\beta)})$ be the solution to $\mathcal{T}(y, u, p, p_1) = \beta$. As a consequence of (A5) it is also a solution to $\underline{\mathcal{T}}(y, u, p, p(0)) = \beta$ with $p \in W_\infty$. Thus the first two equations of this latter equality can be expressed as

$$-p_t - \mathcal{A}^* p - \mathcal{F}'(\bar{y})^* p + y - [\mathcal{F}'(\bar{y})^* \bar{p}]'(y - \bar{y}) = \beta_1 \quad \text{in } L^2(I; V^*), \quad (4.35a)$$

$$\langle \alpha u - B^* p - \beta_2, w - u \rangle_U \geq 0 \quad \text{for all } w \in U_{ad}, \quad (4.35b)$$

where we dropped the dependence of $(y_{(\beta)}, u_{(\beta)}, p_{(\beta)})$ on β . Since $p \in \widetilde{W}_\infty \subset C(\bar{I}; Y)$ inequality (4.35b) implies that $u \in C(\bar{I}; \mathcal{U})$.

Step 2: (Boundedness of $\{p_{(\beta)} : \beta \in \hat{V}\}$). Since \hat{V} is assumed to be bounded, the discussion in Remark 4.2 shows that there exists a constant $M > 0$ such that

$$\|y_{(\beta)}\|_{W_\infty} + \|u_{(\beta)}\|_U \leq M \quad \text{for all } \beta \in \hat{V}.$$

To argue the boundedness of $p_{(\beta)}$, we use similar techniques as in the proof of Proposition 4.1. In the following $\tilde{\delta}, z$ and C_1 are taken as in the proof of that proposition.

For arbitrary $r \in R = \left\{ r \in L^2(I; V^*) : \|r\|_{L^2(I; V^*)} \leq \tilde{\delta} \right\}$, z denotes the solution to

$$z_t - \mathcal{A}z - \mathcal{F}'(\bar{y})z - B(w - u) = -r, \quad z(0) = 0, \quad w = -Kz. \quad (4.36)$$

We know that $\|z\|_{W_\infty} \leq C_1$ and $\|w\|_{L^\infty(I;\mathcal{U})} \leq \eta$ independently of $r \in R$. Due to (4.35a), we have

$$\begin{aligned} \langle p, r \rangle_{L^2(I;V), L^2(I;V^*)} &= \langle p, -z_t + \mathcal{A}z + \mathcal{F}'(\bar{y})z \rangle_{L^2(I;V), L^2(I;V^*)} + \langle B^*p, w - u \rangle_U, \\ &= \langle y - [\mathcal{F}'(\bar{y})^* \bar{p}]'(y - \bar{y}) - \beta_1, z \rangle_{L^2(I;V^*), L^2(I;V)} + \langle \alpha u - \beta_2, w - u \rangle_U, \end{aligned} \quad (4.37)$$

where we used (4.35b) and the feasibility of $w \in U_{ad}$. Consequently

$$\begin{aligned} \langle p, r \rangle_{L^2(I;V), L^2(I;V^*)} &\leq \|z\|_{L^2(I;V)} \left(\|y\|_{L^2(I;V^*)} + \|\beta_1\|_{L^2(I;V^*)} + \|[\mathcal{F}'(\bar{y})^* \bar{p}]'(y - \bar{y})\|_{L^2(I;V^*)} \right) \\ &\quad + (\alpha \|u\|_U + \|\beta_2\|_U) \|w - u\|_U. \end{aligned}$$

The right hand side is uniformly bounded for β in the bounded set \hat{V} and w.r.t. $r \in R$. Hence taking the supremum w.r.t. $r \in R$ we verified that $\left\{ \|p(\beta)\|_{L^2(I;V^*)} : \beta \in \hat{V} \right\}$ is bounded. Boundedness of $\left\{ \|p(\beta)\|_{W_\infty} : \beta \in \hat{V} \right\}$ follows from (4.35a).

Step 3: (Continuity of $p(\beta)$ in W_∞). Let $\{\beta_n\}$ be a convergent sequence in \hat{V} with limit β . Since $\left\{ \|p(\beta)\|_{W_\infty} : n \in \mathbb{N} \right\}$ is bounded, there exists a subsequence $\{\beta_{n_k}\}$ such that $p(\beta_{n_k}) \rightharpoonup \tilde{p}$ weakly in W_∞ and strongly $L^2(I;Y)$. Here we need the compactness of W_∞ in $L^2(I;Y)$, see e.g. [Emm, Satz 8.1.12, pg 213]. Passing to the limit in the variational form of

$$-\partial_t p(\beta_{n_k}) - \mathcal{A}^* p(\beta_{n_k}) - \mathcal{F}'(\bar{y})^* p(\beta_{n_k}) + y - [\mathcal{F}'(\bar{y})^* \bar{p}]'(y(\beta_{n_k}) - \bar{y}) = (\beta_{n_k})_1,$$

we obtain

$$-\partial_t \tilde{p} - \mathcal{A}^* \tilde{p} - \mathcal{F}'(\bar{y})^* \tilde{p} + y(\beta) - [\mathcal{F}'(\bar{y})^* \bar{p}]'(y(\beta) - \bar{y}) = (\beta)_1. \quad (4.38)$$

Since the solution to this equation is unique we have $p(\beta_{n_k}) \rightharpoonup p(\beta)$ weakly in W_∞ . To obtain strong convergence we set $\delta\beta = \beta_{n_k} - \beta$, $\delta p = p(\beta_{n_k}) - p(\beta)$, $\delta y = y(\beta_{n_k}) - y(\beta)$. From (4.35a) we derive that

$$-\partial_t(\delta p) - \mathcal{A}^*(\delta p) - \mathcal{F}'(\bar{y})^*(\delta p) + (I - [\mathcal{F}'(\bar{y})^* \bar{p}]')(\delta y) = (\delta\beta)_1. \quad (4.39)$$

Consider again (4.36) with $r \in R$. Then we obtain

$$\langle \delta p, r \rangle_{L^2(I;V), L^2(I;V^*)} = \langle (I - [\mathcal{F}'(\bar{y})^* \bar{p}]')(\delta y) - (\delta\beta)_1, z \rangle_{L^2(I;V^*), L^2(I;V)} + \langle \delta p, BKz - \bar{u} \rangle_U.$$

This implies that for some $C_2 > 0$,

$$\sup_{r \in R} \langle \delta p, r \rangle_{L^2(I;V), L^2(I;V^*)} \leq C_2 \left(\|\delta y\|_{W_\infty} + \|\delta p\|_{L^2(I;Y)} + \|(\delta\beta)_1\|_{L^2(I;V^*)} \right). \quad (4.40)$$

Since $\|\delta y\|_{W_\infty} \rightarrow 0$, $\|\delta p\|_{L^2(I;Y)} \rightarrow 0$, $\|(\delta\beta)_1\|_{L^2(I;V^*)} \rightarrow 0$ for $n \rightarrow \infty$ this implies that $\|\delta p\|_{L^2(I;V)} \rightarrow 0$. Together with (4.39) this implies that $\lim_{n \rightarrow \infty} \|\delta p\|_{W_\infty} = 0$. \square

Proposition 4.2. *Let assumptions (A) hold and let (\bar{y}, \bar{u}) , and \bar{p} denote a local solution and associated adjoint state state to (P) corresponding to an initial condition $\bar{y}_0 \in B_Y(\bar{\delta}_2)$. Then there exists $\varepsilon > 0$ and $C > 0$ such that for all $\hat{\beta}$ and $\beta \in \hat{V} \cap B_Y(\varepsilon)$*

$$\begin{aligned} \left\| \hat{p}(\hat{\beta}) - p(\beta) \right\|_{W_\infty} + \left\| u(\hat{\beta}) - u(\beta) \right\|_{C(\bar{I}, \mathcal{U})} \\ \leq C \left(\left\| \hat{y}(\hat{\beta}) - y(\beta) \right\|_{W_\infty} + \left\| u(\hat{\beta}) - u(\beta) \right\|_U + \left\| \hat{\beta} - \beta \right\|_Y \right) \end{aligned} \quad (4.41)$$

holds.

Proof. Since $\bar{p} \in W_\infty$ and $\lim_{t \rightarrow \infty} \bar{p}(t) = 0$ in Y , there exists $T > 0$ such that

$$\frac{1}{\alpha} \|B^* \bar{p}(t)\|_Y \leq \frac{\eta}{2}, \quad \forall t > T.$$

Since $p(0) = \bar{p}$ and since by Lemma 4.8, $\beta \in \mathcal{Y} \mapsto p(\beta) \in W_\infty \subset C(\bar{I}; Y)$ is continuous, there exists $\varepsilon > 0$ such that

$$\frac{1}{\alpha} \|B^* p(\beta)(t) + \beta_2(t)\|_Y \leq \frac{\eta}{4}, \quad \forall t \geq T, \quad \forall \beta \in \hat{V} \cap B_{\mathcal{Y}}(\varepsilon).$$

Consequently the constraints are inactive for these parameter values, i.e. we have

$$u_{(\beta)}(t) = \frac{1}{\alpha} [B^* p(\beta)(t) + \beta_2(t)], \quad \|u_{(\beta)}(t)\|_Y \leq \eta, \quad \forall t \geq T, \quad \forall \beta \in \hat{V} \cap B_{\mathcal{Y}}(\varepsilon). \quad (4.42)$$

We next treat separately the cases $[0, T)$ and $[T, \infty)$. We consider first the case $[T, \infty)$ and set $(y, u, p) = (y_{(\beta)}, u_{(\beta)}, p_{(\beta)})$, and $(\hat{y}, \hat{u}, \hat{p}) = (\hat{y}_{(\hat{\beta})}, \hat{u}_{(\hat{\beta})}, \hat{p}_{(\hat{\beta})})$. We shall use that

$$\|\hat{p} - p\|_{L^2(T, \infty; V)} = \sup_{\|r\|_{L^2(T, \infty; V^*)} \leq 1} \int_T^\infty \langle \hat{p}(t) - p(t), r(t) \rangle_{V, V^*} dt.$$

Let $z \in W(T, \infty)$ be such that,

$$z_t - (\mathcal{A} - BK)z - \mathcal{F}'(\bar{y})z = r, \quad z(T) = 0,$$

From Lemma 4.6, see also the proof of Proposition 4.1, we know that there exists a constant $C_1 > 0$ such that $\|z\|_{W(T, \infty)} \leq C_1 \|r\|_{L^2(T, \infty; V^*)}$. Then we can estimate

$$\begin{aligned} \|\hat{p} - p\|_{L^2(T, \infty; V)} &= \sup_{\|r\|_{L^2(T, \infty; V^*)} \leq 1} \int_T^\infty \langle \hat{p} - p, r \rangle_{V, V^*} dt \\ &\leq \sup_{\|r\| \leq 1} \int_T^\infty -\langle (\hat{p}_t - p_t) + \mathcal{A}^*(\hat{p} - p) + \mathcal{F}'(\bar{y})^*(\hat{p} - p), z \rangle_{V^*, V} dt \\ &\quad + \sup_{\|r\| \leq 1} \int_T^\infty \langle B^*(\hat{p} - p), Kz \rangle_{V, V^*} dt. \end{aligned}$$

In the following, C_i denote constants independent of $\hat{\beta}$ and $\beta \in \hat{V} \cap B_{\mathcal{Y}}(\varepsilon)$. From (4.35a) and (4.42) we obtain, for $C_2 > 0$,

$$\begin{aligned} \|\hat{p} - p\|_{L^2(T, \infty; V)} &\leq C_2 \sup_{\|r\| \leq 1} \int_T^\infty \left[\|\hat{y} - y\|_{V^*} + \|[\mathcal{F}'(\bar{y})^* \bar{p}]'(\hat{y} - y)\|_{V^*} + \|\hat{\beta}_1 - \beta_1\|_{V^*} \right. \\ &\quad \left. + \|\hat{\beta}_2 - \beta_2\|_{U \cap C(I; U)} + \alpha \|B^*\| \|K\| \|\hat{u} - u\| \right] \|z\|_V dt. \end{aligned}$$

From (A3) recall that $\|[\mathcal{F}'(\bar{y})^* \bar{p}]'(\hat{y} - y)\|_{L^2(I; V^*)} \leq C_3 \|\hat{y} - y\|_{W_\infty}$. This gives the following estimate for $C_4 > 0$,

$$\|\hat{p} - p\|_{L^2(T, \infty; V)} \leq C_4 \left(\|\hat{y} - y\|_{W_\infty} + \|\hat{u} - u\|_U + \|\hat{\beta} - \beta\|_{\mathcal{Y}} \right). \quad (4.43)$$

By (4.35a) we have $(\hat{p}_t - p_t) \in L^2(T, \infty; V^*)$. Then we obtain $\hat{p} - p \in W(T, \infty)$. Then there exists $C_5 > 0$ independent of $\hat{\beta}$ and $\beta \in \hat{V} \cap B_Y(\varepsilon)$ such that,

$$\|\hat{p} - p\|_{W(T, \infty)} \leq C_5 \left(\|\hat{y} - y\|_{W_\infty} + \|\hat{u} - u\|_U + \|\hat{\beta} - \beta\|_Y \right). \quad (4.44)$$

By the embedding $W(T, \infty) \subset C(T, \infty; Y)$, there exists a constant $C_6 > 0$:

$$\|\hat{p} - p\|_{C([T, \infty); Y)} \leq C_6 \left(\|\hat{y} - y\|_{W_\infty} + \|\hat{u} - u\|_U + \|\hat{\beta} - \beta\|_Y \right). \quad (4.45)$$

Similarly, we estimate on $[0, T]$:

$$\|\hat{p} - p\|_{L^2(0, T; V)} = \sup_{\|r\|_{L^2(0, T; V^*)} \leq 1} \int_0^T \langle \hat{p} - p, r \rangle_{V, V^*} dt. \quad (4.46)$$

Choose z as

$$z_t - (\mathcal{A}z + \mathcal{F}'(\bar{y})z) = r, \quad z(0) = 0,$$

Then there exists $C_7 > 0$ such that $\|z\|_{W(0, T)} \leq C_7 \|r\|_{L^2(0, T; V^*)}$ by Lemma 4.6. Note that C_7 depends on T , but T is fixed. We obtain the following estimate,

$$\begin{aligned} \|\hat{p} - p\|_{L^2(0, T; V)} &\leq \sup_{\|r\| \leq 1} \int_0^T -\langle (\hat{p}_t - p_t) + \mathcal{A}^*(\hat{p} - p) + \mathcal{F}'(\bar{y})^*(\hat{p} - p), z \rangle_{V^*, V} dt \\ &\quad + \|\hat{p}(T) - p(T)\|_Y \|z(T)\|_Y. \end{aligned}$$

Then by a similar computation to that for the $t \in [T, \infty)$ case, we obtain,

$$\|\hat{p} - p\|_{L^2(0, T; V)} \leq C_8 \left(\|\hat{y} - y\|_{W_\infty} + \|\hat{\beta}_1 - \beta_1\|_{L^2(I; V^*)} \right) + \|\hat{p}(T) - p(T)\|_Y \|z(T)\|_Y. \quad (4.47)$$

By (4.45) with $\|z(T)\|_Y \leq C_9$, we obtain

$$\|\hat{p}(T) - p(T)\|_Y \|z(T)\|_Y \leq C_7 C_9 \left(\|\hat{y} - y\|_{W_\infty} + \|\hat{u} - u\|_U + \|\hat{\beta} - \beta\|_Y \right).$$

Combining this estimate with (4.43) and (4.47), we obtain for some $C_{10} > 0$

$$\|\hat{p}_{(\hat{\beta})} - p_{(\beta)}\|_{W_\infty} \leq C_{10} \left(\|\hat{y}_{(\hat{\beta})} - y_{(\beta)}\|_{W_\infty} + \|\hat{u}_{(\hat{\beta})} - u_{(\beta)}\|_U + \|\hat{\beta} - \beta\|_Y \right). \quad (4.48)$$

We also have

$$u_{(\beta)} = \mathbb{P}_{\mathcal{U}_{ad}} \left[-\frac{1}{\alpha} (B^* p_{(\beta)} + \beta_2) \right] \in U \cap C(\bar{I}; \mathcal{U}),$$

and thus

$$\begin{aligned} \|\hat{u}_{(\hat{\beta})}(t) - u_{(\beta)}(t)\|_{\mathcal{U}} &\leq \left\| \mathbb{P}_{\mathcal{U}_{ad}} \left[-\frac{1}{\alpha} (B^* \hat{p}_{(\hat{\beta})}(t) + \hat{\beta}_2(t)) \right] - \mathbb{P}_{\mathcal{U}_{ad}} \left[-\frac{1}{\alpha} (B^* p_{(\beta)}(t) + \beta_2(t)) \right] \right\|_{\mathcal{U}} \\ &\leq \frac{1}{\alpha} \left(\|B^*\| \|\hat{p}_{(\hat{\beta})}(t) - p_{(\beta)}(t)\|_Y + \|\hat{\beta}_2(t) - \beta_2(t)\|_{\mathcal{U}} \right). \end{aligned}$$

This yields

$$\|\hat{u}_{(\hat{\beta})} - u_{(\beta)}\|_{C(\bar{I}; \mathcal{U})} \leq C_{11} \left(\|\hat{p}_{(\hat{\beta})} - p_{(\beta)}\|_{W_\infty} + \|\hat{\beta}_2 - \beta_2\|_{C(\bar{I}; \mathcal{U})} \right), \quad (4.49)$$

and (4.41) follows. \square

Combining Remark 4.2, Step (iii) of the proof of Theorem 2.1, and (4.41) there exists a constant L such that

$$\left\| \hat{y}(\hat{\beta}) - y(\beta) \right\|_{W_\infty} + \left\| \hat{p}(\hat{\beta}) - p(\beta) \right\|_{W_\infty} + \left\| \hat{u}(\hat{\beta}) - u(\beta) \right\|_{U \cap C(\bar{I}; Y)} \leq L \left\| \hat{\beta} - \beta \right\|_{\mathcal{Y}} \quad (4.50)$$

for all $\hat{\beta}$ and $\beta \in \hat{V} \cap B_{\mathcal{Y}}(\varepsilon)$. Thus the verification of (H1)–(H7) is concluded. Here and in the following the p_1 coordinate of the adjoint state coincides with $p(0)$. Therefore it is not indicated.

We now obtain the following corollary to Theorem 2.1.

Corollary 4.9. *Let assumptions (A) hold and let (\bar{y}, \bar{u}) be a local solution of (P) corresponding to an initial datum $\bar{y}_0 \in B_Y(\delta_2)$. Then there exist $\delta_3 > 0$, a neighborhood $\hat{U} = \hat{U}(\bar{y}, \bar{u}, p) \subset W_\infty \times (U \cap C(\bar{I}; \mathcal{U})) \times W_\infty$, and a constant $\mu > 0$ such that for each $y_0 \in B_Y(\bar{y}_0, \delta_3)$ there exists a unique $(y(y_0), u(y_0), p(y_0)) \in \hat{U}$ satisfying the first order condition, and*

$$\|(y(\hat{y}_0), u(\hat{y}_0), p(\hat{y}_0)) - (y(\tilde{y}_0), u(\tilde{y}_0), p(\tilde{y}_0))\|_{W_\infty \times (U \cap C(\bar{I}; \mathcal{U})) \times W_\infty} \leq \mu \|\hat{y}_0 - \tilde{y}_0\|_Y, \quad (4.51)$$

for all $\hat{y}_0, \tilde{y}_0 \in B_Y(\bar{y}_0, \delta_3)$, and $(y(y_0), u(y_0))$ is a local solution of (P).

Next we obtain one of the main results of this paper, the Fréchet differentiability of the local value function associated to (P). By referring to a local value function we pay attention to the fact that for some $y_0 \in B_Y(\delta_2)$, problem (P) may not admit a unique solution. But since due to the second order optimality condition local solutions are locally unique under small perturbations of y_0 , there is a well-defined local value function. We continue to use the notation for \hat{U} and $B_Y(\bar{y}_0, \delta_3)$ of Corollary 4.9.

Theorem 4.1. *(Sensitivity of Cost) Let assumptions (A) hold and let (\bar{y}, \bar{u}) be a local solution of (P) corresponding to an initial datum $\bar{y}_0 \in B_Y(\delta_2)$. Then for each $y_0 \in B_Y(\bar{y}_0, \delta_3)$ the local value function \mathcal{V} associated to (P) is Fréchet differentiable with derivative given by*

$$\mathcal{V}'(y_0) = -p(0; y_0). \quad (4.52)$$

Proof. Let $\bar{y}_0 \in B_Y(\delta_2)$, $y_0 \in B_Y(\bar{y}_0, \delta_3)$, and choose δy_0 sufficiently small so that $y_0 + \delta y_0 \in B_Y(\bar{y}_0, \delta_3)$ as well. Following Corollary 4.9 let $(\tilde{y}(y_0 + s(\delta y_0)), \tilde{u}(y_0 + s(\delta y_0)), \tilde{p}(y_0 + s(\delta y_0))) \in \hat{U}$ for $s \in [0, 1]$ be solutions of the optimality system with $(\tilde{y}(y_0 + s(\delta y_0)), \tilde{u}(y_0 + s(\delta y_0)))$ local solutions to (P). We obtain

$$\begin{aligned} \mathcal{V}(y_0 + s(\delta y_0)) - \mathcal{V}(y_0) &= \left(\frac{1}{2} \|\tilde{y}\|_{L^2(I; Y)}^2 + \frac{\alpha}{2} \|\tilde{u}\|_U^2 \right) - \left(\frac{1}{2} \|y\|_{L^2(I; Y)}^2 + \frac{\alpha}{2} \|u\|_U^2 \right) \\ &= \langle y, \tilde{y} - y \rangle_{L^2(I; Y)} + \alpha \langle u, \tilde{u} - u \rangle_U + \frac{1}{2} \|\tilde{y} - y\|_{L^2(I; Y)}^2 + \frac{\alpha}{2} \|\tilde{u} - u\|_U^2. \end{aligned} \quad (4.53)$$

Observe the identity

$$\begin{aligned} \langle y, \tilde{y} - y \rangle_{L^2(I; Y)} + \alpha \langle u, \tilde{u} - u \rangle_U &= -(p(0), s(\delta y_0))_Y - \langle (\tilde{y}_t - y_t) - \mathcal{A}(\tilde{y} - y) - \mathcal{F}'(y)(\tilde{y} - y), p \rangle + \alpha \langle u, \tilde{u} - u \rangle_U \\ &= -(p(0), s(\delta y_0))_Y - \langle \mathcal{F}(\tilde{y}) - \mathcal{F}(y) - \mathcal{F}'(y)(\tilde{y} - y), p \rangle_{L^2(I; V^*), L^2(I; V)} + \langle \alpha u - B^* p, \tilde{u} - u \rangle_U, \end{aligned}$$

where $p = p(y_0)$. Now we have for $\mathcal{V}(y_0 + s(\delta y_0)) - \mathcal{V}(y_0)$,

$$\mathcal{V}(y_0 + s(\delta y_0)) - \mathcal{V}(y_0) = -(p(0), s(\delta y_0))_Y + \langle \mathcal{F}(y) - \mathcal{F}(\tilde{y}) + \mathcal{F}'(y)(\tilde{y} - y), p \rangle_{L^2(I; V^*), L^2(I; V)}$$

$$+ \langle \alpha u - B^* p, \tilde{u} - u \rangle_U + \frac{1}{2} \|\tilde{y} - y\|_{L^2(I; Y)}^2 + \frac{\alpha}{2} \|\tilde{u} - u\|_U^2. \quad (4.54)$$

Since $p \in L^2(I; V)$, $\|\tilde{y} - y\|_{W_\infty} = O(s)$, and by the continuous Fréchet differentiability of \mathcal{F}' due to (A3) we have

$$|\langle \mathcal{F}'(\tilde{y}) - \mathcal{F}'(y) + \mathcal{F}'(y)(\tilde{y} - y), p \rangle_{L^2(I; V^*), L^2(I; V)}| = o(s) \quad (4.55)$$

Let $s_n \rightarrow 0$ be an arbitrary convergent sequence. By Corollary 4.9 we have that

$$\|\tilde{u}(y_0 + s_n(\delta y_0)) - u(y_0)\|_U \leq \mu s_n(\delta y_0)$$

for all s_n sufficiently small. Hence there exists a subsequence, denoted by the same notation and some \dot{u} such that

$$s_n^{-1} (\tilde{u}(y_0 + s_n(\delta y_0)) - u(y_0)) \rightharpoonup \dot{u} \text{ weakly in } U.$$

Using (4.18), we have

$$\lim_{n \rightarrow \infty} s_n^{-1} \langle \alpha u - B^* p, \tilde{u} - u \rangle_U = \langle \alpha u - B^* p, \dot{u} \rangle_U \geq 0.$$

Analogously

$$\lim_{n \rightarrow \infty} s_n^{-1} \langle \alpha \tilde{u} - B^* p, u - \tilde{u} \rangle_U = \langle \alpha u - B^* p, \dot{u} \rangle_U \leq 0.$$

and hence $\langle \alpha u - B^* p, \dot{u} \rangle_U = 0$. Since the sequence $\{s_n\}$ is arbitrary, we obtain

$$\langle \alpha u - B^* p, \tilde{u} - u \rangle_U = o(s). \quad (4.56)$$

Corollary 4.9 yields,

$$\|\tilde{y}(y_0 + s_n(\delta y_0)) - y(y_0)\|_{L^2(I; Y)}^2 + \alpha \|\tilde{u}(y_0 + s_n(\delta y_0)) - u(y_0)\|_{L^2(I; Y)}^2 = o(s_n). \quad (4.57)$$

Combining (4.55), (4.56), and (4.57) we obtain

$$\lim_{s \rightarrow 0^+} s^{-1} (\mathcal{V}(y_0 + s(\delta y_0)) - \mathcal{V}(y_0)) = -(p(0), (\delta y_0))_Y. \quad (4.58)$$

This implies the Gateaux differentiability. Since $y_0 \rightarrow p(y_0)$ is continuous from $B_Y(\bar{y}_0; \delta_3)$ to $C(\bar{I}, Y)$ the mapping $y_0 \rightarrow \mathcal{V}(y_0)$ is Fréchet differentiable in $B_Y(\bar{y}_0; \delta_3)$. \square

Remark 4.4 (Sensitivity w.r.t. other parameters). We have developed a technique to verify the continuous differentiability of the local value function \mathcal{V} pertaining to a semilinear parabolic equation on infinite time horizon subject to control constraints with respect to small initial data $y_0 \in Y$. Thus the parameter q in (P_q) is the initial condition y_0 . The reason to focus on this case is due to feedback control. Without much additional effort the sensitivity analysis of the value function could be carried out with respect to other parameters as for instance additive noise on the right hand side of the state equation. The papers cited in the introduction, see e.g. [GHH], [GV], consider such situations for the finite horizon case.

5 Proof of Theorem 3.2: Derivation of the HJB Equation.

Utilizing the results established so far we now verify that the (global) value function \mathcal{V} (i.e. the value function associated to global minima) is a solution to a Hamilton–Jacobi–Bellman equation. The initial conditions will be chosen from the neighborhood Y_0 of the origin in Y so that the assertions of Theorem 4.1 and Corollary 4.9 are available. It will be convenient to recall the dynamic programming principle for the infinite time horizon problem: let y_0 be an initial condition for which a solution to (\mathcal{P}) exists. Then for all $\tau > 0$, we have

$$\mathcal{V}(y_0) = \inf_{u \in L^2(0, \tau; \mathcal{U}_{ad})} \int_0^\tau \ell(S(u, y_0; t), u(t)) dt + \mathcal{V}(S(u, y_0; \tau)), \quad (5.1)$$

where $\ell(y, u) = \frac{1}{2} \|y\|_Y^2 + \frac{\alpha}{2} \|u\|_{\mathcal{U}}^2$, and $S(u, y_0; t)$ denotes the solution to (3.2b), (3.2c) on $(0, \tau]$.

For convenience we restate Theorem 3.2. Utilizing the notation that we have already established we can now slightly ease the assumption on the regularity of $\mathcal{F}(\bar{y})$.

Theorem 5.1. *Let assumptions (A) hold and let (\bar{y}, \bar{u}) be a global solution of (\mathcal{P}) corresponding to an initial datum $\bar{y}_0 \in B_Y(\delta_2)$. Let Y_0 denote the subset of initial conditions in $B_Y(\bar{y}_0, \delta_3)$ which allow global solutions in \hat{U} , and assume that for each $y_0 \in \mathcal{D}(\mathcal{A}) \cap Y_0$ there exists $T_{y_0} > 0$ such that $\mathcal{F}(\bar{y}) \in C([0, T_{y_0}); Y)$. Then the following Hamilton–Jacobi–Bellman equation holds at y_0 :*

$$\mathcal{V}'(y)(\mathcal{A}y + \mathcal{F}(y)) + \frac{1}{2} \|y\|_Y^2 + \frac{\alpha}{2} \left\| \mathbb{P}_{\mathcal{U}_{ad}} \left(-\frac{1}{\alpha} B^* \mathcal{V}'(y) \right) \right\|_Y^2 + \left\langle B^* \mathcal{V}'(y), \mathbb{P}_{\mathcal{U}_{ad}} \left(-\frac{1}{\alpha} B^* \mathcal{V}'(y) \right) \right\rangle_Y = 0. \quad (5.2)$$

If for the optimal trajectory $\bar{y}(t) \in B_Y(\bar{y}_0, \delta_3) \cap \mathcal{D}(\mathcal{A})$ for a.a. $t \in (0, \infty)$ and $T_{y_0} = \infty$, then (5.2) holds at a.a. $t \in (0, \infty)$ and

$$\bar{u}(t) = \mathbb{P}_{\mathcal{U}_{ad}} \left(-\frac{1}{\alpha} B^* \mathcal{V}'(\bar{y}(t)) \right). \quad (5.3)$$

Proof. The proof is similar to that of [BKP1, Proposition 10]. For the sake of completeness and since it also requires some changes we provide it here. Choose and fix some $y_0 \in \mathcal{D}(\mathcal{A}) \cap Y_0$. Then the existence of a (globally) optimal pair $(\hat{y}, \hat{u}) \in W_\infty \times U_{ad}$ to (\mathcal{P}) and of an associated adjoint state $\hat{p} \in W_\infty$ with $(\hat{y}, \hat{u}, \hat{p}) \in \hat{U}$ are guaranteed, see Corollary 4.9. In particular we have that $\hat{u}(t) = \mathbb{P}_{\mathcal{U}_{ad}} \left(\frac{1}{\alpha} B^* \hat{p}(t) \right)$, and since $\hat{p} \in C([0, \infty); Y)$ we have that $\hat{u} \in C([0, \infty); Y)$. Let u_0 denote the limit of \hat{u} as time t tends to 0. Since $\hat{y} \in C([0, \infty); Y)$ and since $B_Y(y_0, \delta_3)$ is open there exists $\tau_{y_0} > 0$ such that $\hat{y}(t) \in B_Y(y_0, \delta_3)$, for all $t \in [0, \tau_{y_0})$.

Step 1: Let us first prove that

$$\mathcal{V}'(y_0)(\mathcal{A}y_0 + \mathcal{F}(y_0) + Bu_0) + \ell(y_0, u_0) = 0. \quad (5.4)$$

For this purpose we invoke the dynamic programming principle: We have

$$\frac{1}{\tau} \int_0^\tau \ell(\hat{y}(s), \hat{u}(s)) ds + \frac{1}{\tau} (\mathcal{V}(\hat{y}(\tau)) - \mathcal{V}(y_0)) = 0, \quad (5.5)$$

where we choose $\tau \in (0, \min(T_{y_0}, \tau_{y_0}))$. By continuity of \hat{y} and \hat{u} at time 0, the first term converges to $\ell(y_0, u_0)$ as $\tau \rightarrow 0$. To take $\tau \rightarrow 0$ in the second term we first consider

$$\frac{1}{\tau}(\hat{y}(\tau) - y_0) = \frac{1}{\tau}(e^{A\tau}y_0 - y_0) + \frac{1}{\tau} \int_0^\tau e^{A(\tau-s)} [\mathcal{F}(\hat{y}(s)) + B\hat{u}(s)] ds. \quad (5.6)$$

Using the facts that $y_0 \in \mathcal{D}(\mathcal{A})$, that the terms in square brackets are continuous with values in Y , and that \mathcal{A} generates a strongly continuous semigroup on Y , we can pass to the limit in (5.6) to obtain that

$$\lim_{\tau \rightarrow 0^+} \frac{1}{\tau}(\hat{y}(\tau) - y_0) = \mathcal{A}y_0 + \mathcal{F}(y_0) + Bu_0 \text{ in } Y. \quad (5.7)$$

Now we return to the second term in (5.5) which we express as

$$\frac{1}{\tau}(\mathcal{V}(\hat{y}(\tau)) - \mathcal{V}(y_0)) = \int_0^1 \mathcal{V}'(y_0 + s(\hat{y}(\tau) - y_0)) ds \frac{1}{\tau}(\hat{y}(\tau) - y_0). \quad (5.8)$$

Using (5.7) and since $y \rightarrow \mathcal{V}'(y)$ is continuously differentiable at y_0 , we can pass to the limit in (5.8) to obtain

$$\lim_{\tau \rightarrow 0^+} \frac{1}{\tau}(\mathcal{V}(\hat{y}(\tau)) - \mathcal{V}(y_0)) = \mathcal{V}'(y_0)(\mathcal{A}y_0 + \mathcal{F}(y_0) + Bu_0). \quad (5.9)$$

Now we can pass to the limit in (5.5) and obtain (5.4).

Step 2: For $u \in \mathcal{U}_{ad}$ we define $\tilde{u} \in U_{ad}$ by,

$$\tilde{u}(\tau, x) = \begin{cases} u & \text{for } \tau \in (0, 1) \\ 0 & \text{for } \tau \in [1, \infty) \end{cases}$$

and define $\tilde{y} = S(y_0, \tilde{u})$ as the solution to (3.2b), (3.2c). Then $\tilde{y}(t) \in B_Y(\bar{y}_0, \delta_3)$, for all t sufficiently small, and by (5.1) we have,

$$\frac{1}{\tau} \int_0^\tau \ell(\tilde{y}(s), u(s)) ds + \frac{1}{\tau}(\mathcal{V}(\tilde{y}(\tau)) - \mathcal{V}(y_0)) \geq 0,$$

for all τ sufficiently small. We pass to the limit $\tau \rightarrow 0^+$ with the same arguments as in Step 1 and obtain

$$\mathcal{V}'(y_0)(\mathcal{A}y_0 + \mathcal{F}(y_0) + Bu) + \ell(y_0, u) \geq 0. \quad (5.10)$$

This inequality becomes an equality if $u = u_0$, and thus the quadratic function on the left hand side of (5.10) reaches its minimum 0 at $u = u_0$. This implies that

$$u_0 = \mathbb{P}_{\mathcal{U}_{ad}} \left(-\frac{1}{\alpha} B^* \mathcal{V}'(y_0) \right).$$

Inserting this expression into (5.4) we obtain

$$\mathcal{V}'(y_0)(\mathcal{A}y_0 + \mathcal{F}(y_0)) + \frac{1}{2} \|y_0\|_Y^2 + \frac{\alpha}{2} \left\| \mathbb{P}_{\mathcal{U}_{ad}} \left(-\frac{1}{\alpha} B^* \mathcal{V}'(y_0) \right) \right\|_Y^2 + \left\langle B^* \mathcal{V}'(y_0), \mathbb{P}_{\mathcal{U}_{ad}} \left(-\frac{1}{\alpha} B^* \mathcal{V}'(y_0) \right) \right\rangle_Y = 0. \quad (5.11)$$

Under the additional assumptions on the trajectory, (5.3) follows. \square

6 Some Applications

In this section we discuss the applicability of the framework in two specific cases. It should be noted that even for linear state equations, the sensitivity result for the constraint infinite horizon optimal control problem may be new.

6.1 Fisher's Equation

We consider the optimal stabilization problem for the Fisher equation in an open connected bounded domain Ω in \mathbb{R}^d , $d \in \{1, 2, 3, 4\}$, with Lipschitzian boundary $\Gamma = \partial\Omega$:

$$(\mathcal{P}_{Fis}) \quad \mathcal{V}(y_0) = \min_{(y, u) \in W_\infty \times U_{ad}} \frac{1}{2} \int_0^\infty \|y\|_Y^2 dt + \frac{\alpha}{2} \int_0^\infty \|u\|_{\mathcal{U}}^2 dt, \quad (6.1a)$$

subject to

$$\begin{cases} y_t = \Delta y + y(1 - y) + Bu & \text{in } Q = (0, \infty) \times \Omega & (6.1b) \\ y = 0 & \text{on } \Sigma = (0, \infty) \times \Gamma & (6.1c) \\ y(0) = y_0 & \text{in } \Omega. & (6.1d) \end{cases}$$

where \mathcal{U} and U_{ad} are as in Section 3.1, $B \in \mathcal{L}(\mathcal{U}, Y)$, with $Y = L^2(\Omega)$ and $V = H_0^1(\Omega)$. To further cast this problem in the framework of Section 3, we define the operator

$$\mathcal{A}y = (\Delta + \mathbf{I})y \quad \text{and} \quad y|_\Gamma = 0, \quad \mathcal{D}(\mathcal{A}) = H^2(\Omega) \cap V.$$

Clearly \mathcal{A} has an extension as operator $\mathcal{A} \in \mathcal{L}(V, V^*)$. Moreover it generates an analytic semigroup on Y . Thus (A1) holds. For $\mathcal{U} = Y$ and $B = \mathbf{I}$, condition (A2) is trivially satisfied. Feedback stabilization by finite dimensional controllers was analyzed in [Tri], for example.

It can readily be checked that the nonlinearity $\mathcal{F}(y) = -y^2$ is twice continuously differentiable as mapping $\mathcal{F} : W_\infty \rightarrow L^2(I; V^*)$. The first and second derivatives of \mathcal{F} are given by,

$$\mathcal{F}'(y)v_1 = 2yv_1, \quad \mathcal{F}''(y)(v_1, v_2) = 2(v_1, v_2), \quad \text{for } y, v_1, v_2 \in W_\infty.$$

Since the second derivative is independent of y , its boundedness is automatic. For the sake of illustration we verify the boundedness of the bilinear form of the second derivative on $W_\infty \times W_\infty$. For this purpose, for arbitrary $y \in W_\infty, v_1, v_2 \in W_\infty, \phi \in L^2(I; V)$ we estimate

$$\begin{aligned} \int_0^\infty \langle \mathcal{F}''(y)(v_1, v_2), \phi \rangle_{V^*, V} dt &\leq 2 \int_0^\infty \int_\Omega v_1 v_2 \phi \, dx dt \leq \int_0^\infty \|v_1\|_{L^2(\Omega)} \|v_2\|_{L^4(\Omega)} \|\phi\|_{L^4(\Omega)} \, dt, \\ &\leq C_1 \|v_1\|_{W_\infty} \int_0^\infty \|v_2\|_V \|\phi\|_V \, dt \leq C_2 \|v_1\|_{W_\infty} \|v_2\|_{L^2(I; V)} \|\phi\|_{L^2(I; V)}, \\ &\leq C_3 \|v_1\|_{W_\infty} \|v_2\|_{W_\infty} \|\phi\|_{L^2(I; V)}, \end{aligned} \quad (6.2)$$

where C_i are embedding constants, independent of $y \in W_\infty, v \in W_\infty, \phi \in L^2(I; V)$. We use that V embeds continuously into $L^4(\Omega)$ in dimension up to 4. This implies that $\|\mathcal{F}''(y)(v_1, v_2)\|_{L^2(I; V^*)} \leq C_3 \|v_1\|_{W_\infty} \|v_2\|_{W_\infty}$. Finally we have $\mathcal{F}(0) = \mathcal{F}'(0) = 0$ and thus (A3) and (3.3) are satisfied.

Turning to (A4) we show that $\mathcal{F}(y) : W(0, T) \rightarrow L^1(0, T; V^*)$ is continuous for every $T > 0$. We consider the sequence $y_n \rightharpoonup \hat{y}$ in W_∞ and let $z \in L^\infty(0, T; V)$ be given. Then we estimate

$$\begin{aligned} \int_0^T \langle \mathcal{F}(y_n) - \mathcal{F}(\hat{y}), z \rangle_{V^*, V} dt &= \int_0^T \langle y_n^2 - \hat{y}^2, z \rangle_{V^*, V} = \int_0^T \int_\Omega (y_n - \hat{y})(y_n + \hat{y})z \, dx dt \\ &\leq C_4 \int_0^T \|y_n - \hat{y}\|_Y \|y_n + \hat{y}\|_{L^4(\Omega)} \|z\|_{L^4(\Omega)} \, dt \\ &\leq C_4 \|y_n - \hat{y}\|_{L^2(0, T; Y)} \left[\|y_n\|_{L^2(0, T; V)} + \|\hat{y}\|_{L^2(0, T; V)} \right] \|z\|_{L^\infty(0, T; V)}. \end{aligned}$$

Since V is compactly embedded in Y , we obtain by the Aubin Lions lemma that $\|y_n - \hat{y}\|_{L^2(0, T; Y)} \rightarrow 0$ for $n \rightarrow \infty$. This implies

$$\int_0^T \langle \mathcal{F}(y_n) - \mathcal{F}(\hat{y}), z \rangle_{V^*, V} dt \xrightarrow{n \rightarrow \infty} 0,$$

and (A4) follows. It is simple to check that $\mathcal{F}'(\bar{y}) = 2\bar{y} \in \mathcal{L}(L^2(I; V), L^2(I; V^*))$ and thus (A5) holds as well.

We turn to the assumption $\mathcal{F}(\bar{y}) \in C([0, T_{y_0}); Y)$, for $y_0 \in \mathcal{D}(\mathcal{A})$ and some T_{y_0} , arising in Theorem 3.2 for $y_0 \in \mathcal{D}(\mathcal{A})$. Utilizing the fact that V embeds continuously into $L^4(\Omega)$ in dimension $d \leq 4$ and $\bar{y} \in L^2(I; V)$, we have $\mathcal{F}(\bar{y}) \in L^2(I; Y)$. Hence parabolic regularity theory implies that $\bar{y} \in C([0, \infty); V)$ for $y_0 \in V$, and $\mathcal{F}(\bar{y}) \in C([0, \infty); Y)$ follows.

Remark 6.1. The specificity of this example rests in the fact that the second derivative is independent of the point where it is taken. Other nontrivial cases of analogous structure are reaction diffusion systems with bilinear coupling, see [Gri] where the finite horizon case was treated. Even the case of the Navier Stokes equations falls in this category. Sensitivity for the infinite horizon problems was treated by independent techniques in [BKP3].

6.2 Nonlinearities induced by functions with globally Lipschitz continuous second derivative.

Consider the system (P) with \mathcal{A} associated to a strongly elliptic second order operator with domain $H^2(\Omega) \cap H_0^1(\Omega)$, so that (A1)-(A2) are satisfied. Let $\mathcal{F} : W_\infty \rightarrow L(I; V^*)$ be the Nemytskii operator associated to a mapping $f : \mathbb{R} \rightarrow \mathbb{R}$ which is assumed to be $C^2(\mathbb{R})$ with first and second derivatives globally Lipschitz continuous, and second derivative globally bounded. The regularity assumption $\mathcal{F}(\bar{y}) \in C([0, T_{y_0}); Y)$ for $y_0 \in V = H_0^1(\Omega)$ is satisfied by parabolic regularity theory. We discuss assumption (A3)-(A5) for such an \mathcal{F} , and show that they are satisfied for dimensions $d \in \{1, 2\}$. For the finite horizon problem it will turn out that $d = 3$ is also admissible. By direct calculation it can be checked that \mathcal{F} is continuously Fréchet differentiable for $d \in \{1, 2, 3\}$. We leave this part to the reader and immediately turn to the second derivative.

We proceed by considering the general dimension d to highlight, how the restrictions on the dimension arise. Thus let $d \in \mathbb{N}$ with $d > 1$. The case $d = 1$ can be treated with minor modifications from those in the following steps.

6.2.1 Second derivative of $\mathcal{F}(y)$.

For $y, h_1, h_2 \in W_\infty$ the relevant expression is given by

$$\begin{aligned} & \left\| \mathcal{F}'(y + h_2)h_1 - \mathcal{F}'(y)h_1 - \mathcal{F}''(y)(h_1, h_2) \right\|_{L^2(I; V^*)} \\ &= \sup_{\|\varphi\|_{L^2(I; V)} \leq 1} \langle \mathcal{F}'(y + h_2)h_1 - \mathcal{F}'(y)h_1 - \mathcal{F}''(y)(h_1, h_2), \varphi \rangle_{L^2(I; V^*), L^2(I; V)} \\ &= \sup_{\|\varphi\|_{L^2(I; V)} \leq 1} \int_0^\infty \int_\Omega (\mathfrak{f}'(y(t, x) + h_2(t, x)) - \mathfrak{f}'(y(t, x)) - \mathfrak{f}''(y(t, x))h_2(t, x))h_1(t, x)\varphi(t, x) dx dt \\ &= \sup_{\|\varphi\|_{L^2(I; V)} \leq 1} \int_0^\infty \int_\Omega g(t, x) h_2(t, x)h_1(t, x)\varphi(t, x) dx dt, \end{aligned}$$

where $g(t, x) = \int_0^1 (\mathfrak{f}''(y(t, x) + sh_2(t, x)) - \mathfrak{f}''(y(t, x))) ds$. Note that g is bounded on $I \times \Omega$ and $g \in W_\infty$. Here we use that \mathfrak{f}'' is globally Lipschitz continuous and that $h_1 \in W_\infty$. Henceforth we let $r \in (1, \frac{2d}{d-2}]$ so that $W^{1,2}(\Omega) \subset L^r(\Omega)$ continuously. Let r' denote the conjugate of r so that $r' \in [\frac{2d}{d+2}, \infty)$ for $d > 2$ and $r' \in (1, \infty)$ for $d = 2$. We further choose $\rho > 1, \sigma > 2$ such that $\frac{1}{\rho} + \frac{2}{\sigma} = 1$. Then we estimate

$$\begin{aligned} \left| \int_0^\infty \int_\Omega gh_1h_2\varphi dx dt \right| &\leq \int_0^\infty \left(\int_\Omega |gh_1h_2|^{r'} dx \right)^{1/r'} \|\varphi(t)\|_{L^r(\Omega)} dt \\ &\leq \left(\int_0^\infty \left(\int_\Omega |gh_1h_2|^{r'} dx \right)^{2/r'} dt \right)^{1/2} \left(\int_0^\infty \|\varphi\|_{L^r(\Omega)}^2 dt \right)^{1/2}. \end{aligned}$$

This further implies that

$$\left| \int_0^\infty \int_\Omega gh_1h_2\varphi dx dt \right| \leq C_0 \left[\int_0^\infty \|g(t)\|_{L^{r'\rho}(\Omega)}^2 \|h_1(t)\|_{L^{r'\sigma}(\Omega)}^2 \|h_2(t)\|_{L^{r'\sigma}(\Omega)}^2 dt \right]^{1/2} \|\varphi\|_{L^2(I; V)} \leftarrow (a). \quad (6.3)$$

Here and below $C_i, i = 0, 1, 2, \dots$ denote constant which are independent of y, φ, h_1, h_2 . We next recall Gagliardo's inequality [BF, p 173]:

$$\|u\|_{L^q(\Omega)} \leq \|u\|_{L^2(\Omega)}^{1-d/q-d/2} \|u\|_{W^{1,2}(\Omega)}^{d/2+2/q}, \text{ for all } q > 2, \text{ and } u \in W^{1,2}(\Omega) \equiv V,$$

where $q \in [2, 2^*]$ and

$$q^* \begin{cases} \in [2, \infty] \text{ for } d = 1, \\ \in [2, \infty) \text{ for } d = 2, \\ \in [2, \frac{2d}{d-2}], \text{ for } d > 2. \end{cases}$$

In the above estimate we take, $q = r'\sigma$. We obtain

$$1 + \frac{d}{q} - \frac{d}{2} = \frac{(2-d)r'\sigma + 2d}{2r'\sigma}, \quad \frac{d}{2} - \frac{d}{q} = \frac{d(r'\sigma - 2)}{2r'\sigma}, \text{ also } r'\sigma > 2, (2-d)r'\sigma + 2d > 0.$$

We estimate (6.3), (and check the conditions on the ranges of the parameters below)

$$\begin{aligned}
& \sup_{\|\varphi\|_{L^2(I;V)} \leq 1} (a) \tag{6.4} \\
&= C_1 \left(\int_0^\infty \|g(t)\|_{L^{r'\rho}(\Omega)}^2 \left(\|h_1(t)\|_{L^2(\Omega)} \|h_2(t)\|_{L^2(\Omega)} \right)^{\frac{(2-d)r'\sigma+2d}{r'\sigma}} \left(\|h_1(t)\|_V \|h_2(t)\|_V \right)^{\frac{d(r'\sigma-2)}{r'\sigma}} \right)^{1/2} dt \\
&\leq C_2 \left(\|h_1\|_{W_\infty} \|h_2\|_{W_\infty} \right)^{\frac{(2-d)r'\sigma+2d}{2r'\sigma}} \left(\int_0^\infty \|g(t)\|_{L^{r'\rho}(\Omega)}^2 \left(\|h_1(t)\|_V \|h_2(t)\|_V \right)^{\frac{d(r'\sigma-2)}{r'\sigma}} \right)^{1/2} dt \leftarrow (b)
\end{aligned}$$

We set $\frac{d(r'\sigma-2)}{r'\sigma} = \frac{2}{3}$. This yields,

$$\begin{aligned}
(b) &\leq C_3 \left(\|h_1\|_{W_\infty} \|h_2\|_{W_\infty} \right)^{2/3} \left(\|h_1\|_{W_\infty} \|h_2\|_{W_\infty} \right)^{1/3} \left(\int_0^\infty \|g(t)\|_{L^{r'\rho}(\Omega)}^6 \right)^{1/6} dt \\
&= C_4 \|h_1\|_{W_\infty} \|h_2\|_{W_\infty} \left(\int_0^\infty \left(\int_\Omega |g(t)|^{r'\rho} dx \right)^{6/r'\rho} dt \right)^{1/6}. \tag{6.5}
\end{aligned}$$

Now we check the conditions on the parameter r, σ, d , and r', σ . Since $\frac{d(r'\sigma-2)}{r'\sigma} = \frac{2}{3}$, together with the conditions on r' and σ these parameters need to satisfy

$$r'\sigma = \frac{6d}{3d-2}, \quad r' \in \left[\frac{2d}{d+2}, \infty \right), \quad \sigma \in (2, \infty) \quad r'\sigma \in [2, 2^*], \tag{6.6}$$

and $r' > 1$ if $d = 2$. The last condition above holds without restricting the dimension d . From the first three relations we infer that necessarily $\frac{6d}{3d-2} = r'\sigma > \frac{4d}{d+2}$ which is only possible for $d \leq 3$.

Let us focus on $d = 2$. Then the choice of parameters $r = 6, r' = 6/5, \sigma = 5/2, \rho = 5$ satisfies all the above requirements and it is convenient to further estimate (6.5). In fact we obtain

$$\begin{aligned}
& \left\| \mathcal{F}'(y + h_2)h_1 - \mathcal{F}'(y)h_1 - \mathcal{F}''(y)(h_1, h_2) \right\|_{L^2(I;V^*)} \\
&\leq C_5 \|h_1\|_{W_\infty} \|h_2\|_{W_\infty} \left(\int_0^\infty \left(\int_\Omega |g(t, x)|^6 dx \right) dt \right)^{1/6} \\
&\leq C_6 \|h_1\|_{W_\infty} \|h_2\|_{W_\infty} \left(\int_0^\infty \left(\int_\Omega |g(t, x)|^2 dx \right) dt \right)^{1/6}
\end{aligned}$$

for all $y, h_1, h_2 \in W_\infty$. Here we use the boundedness of g . By Lebesgue's bounded convergence theorem the last factor converges to 0 for $\|h_2\|_{W_\infty} \rightarrow 0$ and hence the fact that \mathcal{F} is twice differentiable is verified. The continuity of the second derivative follows with the above estimates and again by the Lebesgue theorem.

Next we consider $d = 3$. In this case an analogous procedure is not possible, since the relations (6.6) and $r'\rho \leq 6$ cannot be fulfilled simultaneously. In fact, $r'\sigma = 18/7, r' \geq 6/5$, and thus necessarily $\sigma \in (2, 15/7]$. The condition $r'\rho \leq 6$ is equivalent to $12 \leq \sigma(6 - r') = 18/7r'(6 - r')$, which in turn is equivalent to $17r' \leq 18$, which contradicts $r' \geq 6/5$.

Thus we fix parameters r and σ such that (6.6) are satisfied for $d = 3$, as for instance $r' = 6/5, \sigma = 15/7$, which implies that $\rho = 15$ and $r'\rho = 18$. Then for the *finite horizon problem* we can estimate by Hölder's inequality with $\eta = r'\rho/6$:

$$\begin{aligned} (b) &\leq C_7 \|h_1\|_{W_\infty} \|h_2\|_{W_\infty} \left(\int_0^T \left(\int_\Omega |g(t,x)|^2 dx \right)^{6/r'\rho} dt \right)^{1/6} \\ &\leq C_8 \|h_1\|_{W_\infty} \|h_2\|_{W_\infty} \left(\int_0^T \left(\int_\Omega |g(t,x)|^2 dx \right) dt \right)^{1/6\eta} T^{1/6\eta'}. \end{aligned}$$

From here we can proceed as in the case $d = 2$ to assert the continuous second Fréchet differentiability of \mathcal{F} in $d = 3$ for the finite horizon case.

6.2.2 Assumptions (A4) and (A5).

In order to verify (A4), we show that $\mathcal{F}(y) : W(0, T) \rightarrow L^1(0, T; V^*)$ is continuous for every $T > 0$. We consider the sequence $y_n \rightarrow \hat{y}$ in $W(0, T)$ and let $z \in L^\infty(0, T; V)$ be given. Then we estimate

$$\int_0^T \langle \mathcal{F}(y_n) - \mathcal{F}(\hat{y}), z \rangle_{V, V^*} dt = \int_0^T \int_\Omega (\mathfrak{f}(y_n) - \mathfrak{f}(\hat{y}))z \, dx dt \leq C \|y_n - \hat{y}\|_{L^2(0, T; Y)} \|z\|_{L^2(0, T; Y)}.$$

Then by the compactness of V in Y , we obtain (A4).

Now we verify (A5). We recall Remark 3.1, and proceed as in (6.2) for $y \in W_\infty, \varphi \in L^2(I; V)$,

$$\begin{aligned} \|\mathcal{F}'(y)^* p\|_{L^2(I; V^*)} &= \|(\mathcal{F}'(y)^* - \mathcal{F}'(0)^*)p\|_{L^2(I; V^*)} = \sup_{\|\varphi\|_{L^2(I; V)} \leq 1} \int_0^\infty \int_\Omega \langle (\mathcal{F}'(y)^* - \mathcal{F}'(0)^*)p, \varphi \rangle_{V^*, V} \\ &= \sup_{\|\varphi\|_{L^2(I; V)} \leq 1} \int_0^\infty \int_\Omega (\mathfrak{f}'(y) - \mathfrak{f}'(0))p\varphi \, dx dt \leq C \|y\|_{W_\infty} \|p\|_{L^2(I; V)}. \end{aligned}$$

This shows $\mathcal{F}'(y)^*$ satisfies (A5).

6.3 Cubic nonlinearity y^3 in one dimension ($\Omega \subset \mathbb{R}$).

We can also consider the optimal stabilization problem with cubic nonlinearity, i.e. $\mathcal{F}(y) = y^3$ in one dimension. This is a special monotone case of the Schlögl model of theoretical chemistry.

$$(\mathcal{P}_{Sch}) \quad \mathcal{V}(y_0) = \min_{(y, u) \in W_\infty \times U_{ad}} \frac{1}{2} \int_0^\infty \|y\|_Y^2 dt + \frac{\alpha}{2} \int_0^\infty \|u\|_{\mathcal{U}}^2 dt, \quad (6.7a)$$

subject to

$$\begin{cases} y_t = \Delta y + y^3 + Bu & \text{in } Q = (0, \infty) \times \Omega & (6.7b) \\ y = 0 & \text{on } \Sigma = (0, \infty) \times \Gamma & (6.7c) \\ y(0) = y_0 & \text{in } \Omega. & (6.7d) \end{cases}$$

In this model, one can easily verify assumption (A1) is satisfied by taking $\mathcal{A}y = \Delta y, y|_\Gamma = 0$, and $\mathcal{D}(\mathcal{A}) = H^2(\Omega) \cap V$. Clearly \mathcal{A} can be extended to $\mathcal{A} \in \mathcal{L}(V, V^*)$. Moreover \mathcal{A} generates an analytic semigroup on Y which is uniformly stable. Assumption (A2) is satisfied under the

same argumentation as in Fisher's equation. Differentiability assumption (A3), and continuity assumption (A4) are satisfied along similar computations as in subsections 6.2.1, 6.2.2. For (A5) we require that $y_0 \in V$. Indeed in this case for $\bar{y} \in W_\infty$ by Gagliardo's inequality

$$\int_0^\infty \int_\Omega |\bar{y}^3|^2 dx dt = \int_0^\infty \|\bar{y}\|_{L^6(\Omega)}^6 dt \leq \int_0^\infty \|\bar{y}\|_{L^2(\Omega)}^4 \|\bar{y}\|_V^2 dt \leq C \|\bar{y}\|_{W_\infty}^4 \int_0^\infty \|\bar{y}\|_V^2 dt \leq C \|\bar{y}\|_{W_\infty}^6.$$

Thus $\bar{y}^3 \in L^2(I; Y)$ and parabolic regularity theory implies that $\bar{y} \in C(I; V)$ if $y_0 \in V$. We estimate for $h, \varphi \in L^2(I; V)$, suppressing the arguments (t, x) ,

$$\begin{aligned} \left| \int_0^\infty \int_\Omega \mathcal{F}'(\bar{y}) h \varphi dx dt \right| &\leq \left| \int_0^\infty \int_\Omega \bar{y}^2 h \varphi dx dt \right| \leq \int_0^\infty \|\bar{y}\|_{L^4(\Omega)}^2 \|h\|_{L^4(\Omega)} \|\varphi\|_{L^4(\Omega)} dt, \\ &\leq C \|\bar{y}\|_{C(I; V)}^2 \|h\|_{L^2(I; V)} \|\varphi\|_{L^2(I; V)} \end{aligned}$$

which implies (A5). Moreover we have $\mathcal{F}(\bar{y}) \in C([0, T_{y_0}); Y)$, since $V \subset C(\bar{\Omega})$ in dimension 1, and thus the extra regularity demanded in Theorem 3.2 is satisfied.

References

- [BPDM] A. Bensoussan, G. Da Prato, M. Delfour, S. Mitter Representation and Control of Infinite Dimensional Systems, *Birkhäuser Boston Basel Berlin*, 2007.
- [BM] M. Bergounioux, N. Merabet, *Sensitivity analysis for optimal control problems governed by semilinear parabolic equations*, *Control and Cybernetics*, vol: 28(1999), No 3.
- [BF] F. Boyer, P. Fabrie, *Mathematical Tools for the Study of the Incompressible Navier-Stokes Equations and Related Models*, *Springer-Verlag New York*, 2013, pp 526.
- [BKP1] T. Breiten, K. Kunisch, L. Pfeiffer, *Infinite-horizon bilinear optimal control problems: Sensitivity analysis and polynomial feedback laws*, *SIAM Journal on Control and Optimization*, 56 (2018), pp. 3184-3214.
- [BKP2] T. Breiten, K. Kunisch, L. Pfeiffer, *Taylor Expansion of the value Function associated with a bilinear optimal control problem*, *Annales de l'Institut Henri Poincaré C, Analyse Non Linéaire*, (2019).
- [BKP3] T. Breiten, K. Kunisch, L. Pfeiffer, *Feedback Stabilization of the Two-Dimensional Navier-Stokes Equations by Value Function Approximation*, *Appl. Math. Optim.* 80, 599–641 (2019).
- [CF] P. Cannarsa, H. Frankowska, *Local regularity of the value function in optimal control*, *Systems & Control Letters*, Vol 62, Issue 9, 2013, pp 791-794, ISSN 0167-6911, <https://doi.org/10.1016/j.sysconle.2013.06.001>.
- [Cas] E. Casas, *Pontryagin's Principle for State-Constrained Boundary Control Problems of Semilinear Parabolic Equations*, *SIAM J. Control Optim.*, 35(4), 1297-1327. (31 pages), 1997. <https://doi.org/10.1137/S0363012995283637>
- [CK] E. Casas, K. Kunisch, *Stabilization by Sparse Controls for a Class of Semilinear Parabolic Equations*, *SIAM J. Control Optim.*, 55(1): 512-532, 2017.

- [Don] A. L. Dontchev, *Implicit function theorems for generalized equations*, *Mathematical Programming* 70, 91–106 (1995). <https://doi.org/10.1007/BF01585930>.
- [Emm] E. Emmrich, *Gewöhnliche und Operator Differentialgleichungen*, Vieweg, Wiesbaden, 2004.
- [Goe] R. Goebel, *Convex Optimal Control Problems with Smooth Hamiltonians*, *SIAM J. Control Optim.*, 43(5), 1787-1811 (2005). (25 pages). <https://doi.org/10.1137/S0363012902411581>.
- [Gri] R. Griesse, *Parametric sensitivity analysis in optimal control of reaction diffusion system - Part I: Solution Differentiability*, *Numerical Functional Analysis and Optimization*, 25(1-2):93-117, 2004.
- [GHH] R. Griesse, M. Hintermüller, Michael Hinze, *Differential stability of control-constrained optimal control problems for the Navier-Stokes equations*, *Numerical Functional Analysis and Optimization*, 26:7-8, 829-850, 2005.
- [GV] R. Griesse, S. Volkwein, *Parametric Sensitivity Analysis for Optimal Boundary Control of a 3D Reaction-Diffusion System*, *Large-Scale Nonlinear Optimization. Nonconvex Optimization and Its Applications*, vol 83. (2006) Springer, Boston, MA.
- [IK] K. Ito, K. Kunisch, *Lagrange Multiplier Approach to Variational Problems and Applications*, *Society for Industrial and Applied Mathematics*, Philadelphia, PA, USA, 2008.
- [LM] J.-L. Lions, E. Magenes, *Non-Homogeneous Boundary Value Problems and Applications: Vol. 1*, *Die Grundlehren der mathematischen Wissenschaften* 181, Springer-Verlag, Berlin, 1972.
- [Mal] K. Malanowski, *Sensitivity analysis for parametric optimal control of semilinear parabolic equations*, *Journal of Convex Analysis*, Vol 9(2002), No. 2, 543-569.
- [MT] K. Malanowski, F. Tröltzsch, *Lipschitz Stability of Solutions to Parametric Optimal Control for Parabolic Equations*, *Journal of Analysis and its Applications*, 18(2):469-489, 1999.
- [MS] C. Meyer, L. M. Susu, *Optimal control of nonsmooth, semilinear parabolic equations*, *SIAM J. Control Optim.* Vol. 55(2017), No. 4, pp. 2206-2234.
- [MZ] H. Maurer, J. Zowe, *First and second-order necessary and sufficient optimality conditions for infinite-dimensional programming problems*, *Math. Programming* 16(1979), 98-110.
- [Paz] A. Pazy, *Semigroups of Linear Operators and Applications to Partial Differential Equations*, *Springer-Verlag New York*, 1983.
- [Rob] S. M. Robinson, *Strongly regular generalized equations*, *Math. of Oper. Res.* 5(1980), 43-62.
- [Tri] R. Triggiani, *On the Stabilizability Problem of Banach Spaces*, *J. Math. Anal. Appl.* 55,1975, pp 303-403.
- [Tro1] F. Tröltzsch, *Lipschitz stability of solutions to linear-quadratic parabolic control problems with respect to perturbations*, *Dynamics of Continuous, Discrete and Impulsive Systems Series A Mathematical Analysis*, 7(2):289-306, 2000.

[Tro2] F. Tröltzsch, *Optimal Control of Partial Differential Equations: Theory, Methods and Applications* AMS, ISBN 978-0-8218-4904-0, 2010, pp 399.

[Wac] D. Wachsmuth, *Regularity and stability of optimal controls of nonstationary Navier-Stokes equations*, *Control and Cybernetics*, vol 34(2005), No 2.

Received xxxx 20xx; revised xxxx 20xx.