

Error estimates for the numerical approximation of optimal control problems with non-smooth pointwise-integral control constraints[†]

EDUARDO CASAS[‡]

*Department of Applied Mathematics and Computer Science, University of Cantabria
Avda. de los Castros s/n, 39005 Santander, Spain*

KARL KUNISCH[§]

*Institute for Mathematics and Scientific Computing, University of Graz
Heinrichstrasse 36, A-8010 Graz, Austria
and Johann Radon Institute, Austrian Academy of Sciences, Linz, Austria*

AND

MARIANO MATEOS[¶]

Department of Mathematics, University of Oviedo, 33203 Gijón, Spain

[Received on xxxxx; revised on xxxxx]

The numerical approximation of an optimal control problem governed by a semilinear parabolic equation and constrained by a bound on the spatial L^1 -norm of the control at every instant of time is studied. Spatial discretizations of the controls by piecewise constant and continuous piecewise linear functions are investigated. Under finite element approximations the sparsity properties of the continuous solutions are preserved in a natural way using piecewise constant approximations of the control, but suitable numerical integration of the objective functional and of the constraint must be used to keep the sparsity pattern when using spatial continuous piecewise linear approximations. We also obtain error estimates and finally present some numerical examples.

Keywords: optimal control; semilinear partial differential equations; discontinuous Galerkin approximations; error estimates.

1. Introduction

In this paper, we study the numerical approximation of the optimal control problem

$$(P) \quad \inf_{u \in U_{ad}} J(u) := \frac{1}{2} \int_Q (y_u(x, t) - y_d(x, t))^2 dx dt + \frac{\kappa}{2} \int_Q u(x, t)^2 dx dt,$$

[†]The first and third authors were supported by Spanish Ministerio de Economía, Industria y Competitividad under research project MTM2017-83185-P. The second was supported by the ERC advanced grant 668998 (OCLOC) under the EU's H2020 research program.

[‡]Corresponding author. Email: eduardo.casas@unican.es

[§]Email: karl.kunisch@uni-graz.at

[¶]Email: mmateos@uniovi.es

where $\kappa > 0$, $Q = \Omega \times (0, T)$, with $\Omega \subset \mathbb{R}^n$, $n = 2$ or 3 , a convex polygonal/polyhedral domain with boundary Γ and $0 < T < +\infty$ is fixed,

$$U_{ad} = \{u \in L^\infty(Q) : \|u(t)\|_{L^1(\Omega)} \leq \gamma \text{ for a.a. } t \in (0, T)\}$$

with $0 < \gamma < +\infty$. Further y_u is the solution of the semilinear parabolic equation

$$\begin{cases} \frac{\partial y}{\partial t} + Ay + a(x, t, y) = u & \text{in } Q = \Omega \times (0, T), \\ y = 0 \text{ on } \Sigma = \Gamma \times (0, T), \quad y(0) = y_0 \text{ in } \Omega. \end{cases} \quad (1.1)$$

with

$$Ay = - \sum_{i,j=1}^n \partial_{x_j}(a_{ij}(x)\partial_{x_i}y).$$

Precise assumptions on the operator A and the nonlinearity a are given below.

This problem was studied in Casas & Kunisch (2021), where the authors proved existence of a solution, and obtained first and second order optimality conditions. As it is emphasized in that paper, there are two special difficulties in the study of (P). The first one is given by the fact that, in order to be able to deal with strong non-linear terms such as $a(x, t, y) = a_0(x, t) \exp(y)$ with $a_0 \in L^\infty(Q)$, the framework for the control space cannot be $L^2(Q)$, but should be $L^q(Q)$ with q large enough. This implies that the usual techniques to prove existence of a solution fail, rather, a truncation argument on a is used for this purpose. The second difficulty is the non-differentiability of the constraint. First order optimality conditions are obtained using the convexity of U_{ad} . Second order optimality conditions require a careful setting of the cone of critical directions in order to obtain sufficient conditions with a minimal gap with respect to the necessary ones. With the aid of first order optimality conditions, sparsity properties of the optimal control are derived.

There are numerous references regarding the numerical analysis of problems governed by partial differential equations. Not trying to be exhaustive, and considering only distributed optimal control problems governed by parabolic equations, we can cite Meidner & Vexler (2011) (linear equation, no constraints), Chrysafinos & Karatzas (2012) (semilinear equation, but only dimension 2 and not strong non-linear terms, no constraints), Akman *et al.* (2014) (discontinuous elements for linear convection-diffusion), Leykekhman & Vexler (2013); Gong *et al.* (2014); von Daniels *et al.* (2015) (linear, pointwise control-constraints), Huang *et al.* (2018), (space-time spectral discretization), Casas *et al.* (2017b, 2018) (semilinear, sparsity-promoting term in the functional, no constraints), Casas *et al.* (2019) (semilinear, pointwise control-constraints, no Thikonov regularization), Christof & Vexler (2021) (linear, state constraints), Hoppe & Neitzel (2020) (quasilinear, pointwise state constraints).

The only reference that we have been able to find with a pointwise constraint in time on the norm of the control is Gunzburger & Manservigi (1999). In that reference, the authors impose the differentiable constraint $\|u(\cdot, t)\|_{L^2(\Omega)}^2 \leq 1$. However, they do not address the obtention of error estimates for the discrete problems.

Our objectives in this paper are to discretize (P) in such a way that the sparsity properties are preserved to prove convergence of the discrete solutions to the solutions of the continuous problem, and to obtain error estimates. To discretize the state equation, we use a discontinuous Galerkin scheme, computationally equivalent to the implicit Euler method. For the discretization in space of the state and the adjoint state, continuous piecewise linear finite elements are used, while for the control we study both piecewise constant and continuous piecewise linear approximations. The use of piecewise constant elements leads in a natural way to sparsity properties of the discrete optimal control consistent with

those obtained for the continuous problem, but a straightforward discretization of (P) using continuous piecewise linear space-approximations of the control, may result in a loss of the sparsity properties due to the use of a mass matrix. To overcome this difficulty, we discretize the norm in $L^p(\Omega)$, $p = 1, 2$ with the help of the lumped mass matrix and use Carstensen's quasi interpolation operator. A similar technique for problems with sparsity-promoting terms in the functional was used in Casas *et al.* (2012) for a problem governed by a semilinear elliptic equation and in Casas *et al.* (2018) for a problem governed by a semilinear parabolic equation; this technique is also found in the thesis by Pieper (2015) and in Rösch & Wachsmuth (2017).

The plan of this paper is as follows. In Section 2 we recall some results from Casas & Kunisch (2021) concerned with the continuous problem. In Section 3 the problem is discretized and the sparsity properties of the discrete solution are established. In Section 4 we prove convergence and obtain error estimates. Finally, in Section 5, numerical examples are presented to illustrate the results obtained in the paper.

2. Assumptions and preliminary results

We make the following assumptions along this paper.

Assumption 1- $\Omega \subset \mathbb{R}^n$, $n = 2$ or 3 , is a convex polygonal/polyhedral domain, and $0 < T < \infty$ is fixed. Γ denotes the boundary of Ω . The coefficients of the operator A satisfy: a_{ij} are Lipschitz functions in $\bar{\Omega}$ for every $1 \leq i, j \leq n$, and

$$\Lambda_A |\xi|^2 \leq \sum_{i,j=1}^n a_{ij}(x) \xi_i \xi_j \quad \forall \xi \in \mathbb{R}^n \quad \text{and for a.a. } x \in \Omega \quad (2.1)$$

for some $\Lambda_A > 0$. For the initial state we suppose that $y_0 \in H_0^1(\Omega) \cap C^{0,\alpha}(\bar{\Omega})$, where $C^{0,\alpha}(\bar{\Omega})$ denotes the space of α -Hölder continuous functions in $\bar{\Omega}$ with $\alpha \in (0, 1]$.

Assumption 2- We assume that $a : Q \times \mathbb{R} \rightarrow \mathbb{R}$ is a Carathéodory function of class C^2 with respect to the last variable satisfying the following properties:

$$\exists C_a \in \mathbb{R} : \frac{\partial a}{\partial y}(x, t, y) \geq C_a \quad \forall y \in \mathbb{R}, \quad (2.2)$$

$$a(\cdot, \cdot, 0) \in L^{\hat{r}}(0, T; L^2(\Omega)), \quad \text{with } \hat{r} > \frac{4}{4-n}, \quad (2.3)$$

$$\forall M > 0 \exists C_{a,M} > 0 : \left| \frac{\partial^j a}{\partial y^j}(x, t, y) \right| \leq C_{a,M} \quad \forall |y| \leq M \quad \text{and } j = 1, 2, \quad (2.4)$$

$$\forall \rho > 0 \text{ and } \forall M > 0 \exists \varepsilon > 0 \text{ such that} \\ \left| \frac{\partial^2 a}{\partial y^2}(x, t, y_1) - \frac{\partial^2 a}{\partial y^2}(x, t, y_2) \right| < \rho \quad \forall |y_1|, |y_2| \leq M \text{ with } |y_1 - y_2| < \varepsilon, \quad (2.5)$$

for almost all $(x, t) \in Q$.

Assumption 3- In the control problem (P), we assume that $\kappa > 0$, $\gamma > 0$, and $y_d \in L^{\hat{r}}(0, T; L^2(\Omega))$.

As usual we denote $H^{2,1}(Q) = L^2(0, T; H^2(\Omega) \cap H_0^1(\Omega)) \cap H^1(0, T; L^2(\Omega))$. Then, we have the following result.

THEOREM 2.1 Under under Assumptions 1 and 2, for every $u \in L^r(0, T; L^p(\Omega))$ with $\frac{1}{r} + \frac{n}{2p} < 1$ and $r, p \geq 2$ there exists a unique solution $y_u \in C^{0,\beta}(\bar{Q}) \cap H^{2,1}(Q)$ of (1.1) with $\beta \in (0, \alpha)$. Moreover, the

following estimate holds

$$\begin{cases} \|y_u\|_{C^{0,\beta}(\bar{Q})} + \|y_u\|_{H^{2,1}(Q)} \leq \eta(\|u\|_{L^r(0,T;L^p(\Omega))} + M_{\hat{r},0}), \\ \|y_u\|_{L^\infty(0,T;L^2(\Omega))} + \|y_u\|_{L^2(0,T;H_0^1(\Omega))} \leq C(\|u\|_{L^2(Q)} + M_{2,0}) \end{cases} \quad (2.6)$$

for a constant C and a monotone non-decreasing function $\eta : [0, \infty) \rightarrow [0, \infty)$ with $\eta(0) = 0$ independent of u , and

$$\begin{aligned} M_{\hat{r},0} &= \|a(\cdot, \cdot, 0)\|_{L^{\hat{r}}(0,T;L^2(\Omega))} + \|y_0\|_{C^{0,\alpha}(\bar{\Omega})} + \|y_0\|_{H_0^1(\Omega)}, \\ M_{2,0} &= \|a(\cdot, \cdot, 0)\|_{L^2(Q)} + \|y_0\|_{L^2(\Omega)}. \end{aligned}$$

The existence of a unique solution of (1.1) in the space $L^2(0, T; H_0^1(\Omega)) \cap L^\infty(Q)$ as well as the estimates in $L^\infty(0, T; L^2(\Omega))$ and $L^2(0, T; H_0^1(\Omega))$ were proved in Casas & Kunisch (2021). The $H^{2,1}(Q)$ regularity is a well known consequence of the convexity of Ω and the $H_0^1(\Omega)$ regularity of y_0 . The reader is referred to Ladyzhenskaya *et al.* (1988, Chap. III-§10) or Di Benedetto (1986) for the $C^{0,\beta}(\bar{Q})$ regularity.

Taking $p = 2$ and $r \in (\frac{4}{4-n}, \infty]$ we have that $\frac{1}{r} + \frac{n}{4} < 1$ and $r > 2$. Then, from Theorem 2.1 we deduce that the mapping $G : L^r(0, T; L^2(\Omega)) \rightarrow H^{2,1}(Q) \cap L^\infty(Q)$ given by $G(u) = y_u$ is well defined. Further, we have the following differentiability properties.

THEOREM 2.2 The mapping G is of class C^2 . For $u, v, v_1, v_2 \in L^r(0, T; L^2(\Omega))$ the derivatives $z_v = G'(u)v$ and $z_{v_1, v_2} = G''(u)(v_1, v_2)$ are the solutions of the equations

$$\begin{cases} \frac{\partial z_v}{\partial t} + Az_v + \frac{\partial a}{\partial y}(x, t, y_u)z_v = v \text{ in } Q, \\ z_v = 0 \text{ on } \Sigma, \quad z_v(0) = 0 \text{ in } \Omega, \end{cases} \quad (2.7)$$

$$\begin{cases} \frac{\partial z_{v_1, v_2}}{\partial t} + Az_{v_1, v_2} + \frac{\partial a}{\partial y}(x, t, y_u)z_{v_1, v_2} + \frac{\partial^2 a}{\partial y^2}(x, t, y_u)z_{v_1}z_{v_2} = 0 \text{ in } Q, \\ z_{v_1, v_2} = 0 \text{ on } \Sigma, \quad z_{v_1, v_2}(0) = 0 \text{ in } \Omega \end{cases} \quad (2.8)$$

where $z_{v_i} = G'(u)v_i$, $i = 1, 2$.

This theorem was proved in Casas & Kunisch (2021) with a change in the range of G , namely $G : L^r(0, T; L^2(\Omega)) \rightarrow L^2(0, T; H_0^1(\Omega)) \cap H^1(0, T; H^{-1}(\Omega)) \cap L^\infty(Q)$. The proof given there can be adapted using the extra regularity of the data of the state equation and Theorem 2.1.

Theorem 2.2 along with the chain rule leads to the following differentiability properties of the cost functional J .

COROLLARY 2.1 If $r > \frac{4}{4-n}$, then $J : L^r(0, T; L^2(\Omega)) \rightarrow \mathbb{R}$ is of class C^2 and its derivatives are given by the expressions

$$J'(u)v = \int_Q (\varphi + \kappa u)v \, dx \, dt, \quad (2.9)$$

$$J''(u)(v_1, v_2) = \int_Q \left[\left(1 - \frac{\partial^2 a}{\partial y^2}(x, t, y_u)\varphi\right)z_{v_1}z_{v_2} + \kappa v_1 v_2 \right] \, dx \, dt, \quad (2.10)$$

where $z_{v_i} = G'(u)v_i$, $i = 1, 2$, and $\varphi \in C^{0,\beta}(\bar{Q}) \cap H^{2,1}(Q)$ is the solution of the adjoint state equation

$$\begin{cases} -\frac{\partial \varphi}{\partial t} + A^* \varphi + \frac{\partial a}{\partial y}(x, t, y_u) \varphi = y_u - y_d & \text{in } Q, \\ \varphi = 0 \text{ on } \Sigma, \quad \varphi(T) = 0 & \text{in } \Omega, \end{cases} \quad (2.11)$$

with $A^* \varphi = -\sum_{i,j=1}^n \partial_{x_j}(a_{ji}(x) \partial_{x_i} \varphi)$ the adjoint operator of A

Concerning the control problem (P), the following theorem and corollaries follow from Casas & Kunisch (2021).

THEOREM 2.3 There exists at least one solution of (P). Moreover, for every local minimizer \bar{u} in the $L^r(0, T; L^2(\Omega))$ sense with $r > \frac{4}{4-n}$, there exist $\bar{y} \in H^{2,1}(Q) \cap C^{0,\theta}(\bar{Q})$, $\bar{\varphi} \in C(\bar{Q}) \cap H^{2,1}(Q)$, and $\bar{\mu} \in L^\infty(Q)$ such that

$$\begin{cases} \frac{\partial \bar{y}}{\partial t} + A\bar{y} + a(x, t, \bar{y}) = \bar{u} & \text{in } Q, \\ \bar{y} = 0 \text{ on } \Sigma, \quad \bar{y}(0) = y_0 & \text{in } \Omega, \end{cases} \quad (2.12)$$

$$\begin{cases} -\frac{\partial \bar{\varphi}}{\partial t} + A^* \bar{\varphi} + \frac{\partial a}{\partial y}(x, t, \bar{y}) \bar{\varphi} = \bar{y} - y_d & \text{in } Q, \\ \bar{\varphi} = 0 \text{ on } \Sigma, \quad \bar{\varphi}(T) = 0 & \text{in } \Omega, \end{cases} \quad (2.13)$$

$$\int_Q \bar{\mu}(u - \bar{u}) \, dx \, dt \leq 0 \quad \forall u \in U_{ad}, \quad (2.14)$$

$$\bar{\varphi} + \kappa \bar{u} + \bar{\mu} = 0. \quad (2.15)$$

Let us denote by $\text{Proj}_{B_\gamma} : L^2(\Omega) \rightarrow B_\gamma \cap L^2(\Omega)$ the $L^2(\Omega)$ projection, where $B_\gamma = \{v \in L^1(\Omega) : \|v\|_{L^1(\Omega)} \leq \gamma\}$.

COROLLARY 2.2 Let \bar{u} , $\bar{\varphi}$, and $\bar{\mu}$ satisfy (2.12)–(2.15) and assume that $\bar{u} \in U_{ad}$. Then, the following properties hold

$$\int_\Omega \bar{\mu}(t)(v - \bar{u}(t)) \, dx \leq 0 \quad \forall v \in B_\gamma \text{ and for a.a. } t \in (0, T), \quad (2.16)$$

$$\bar{u}(t) = \text{Proj}_{B_\gamma} \left(-\frac{1}{\kappa} \bar{\varphi}(t) \right) \text{ for a.a. } t \in (0, T), \quad (2.17)$$

$$\begin{cases} \bar{u}(x, t) \bar{\mu}(x, t) = |\bar{u}(x, t)| |\bar{\mu}(x, t)| \text{ for a.a. } (x, t) \in Q, \\ \text{if } \|\bar{u}(t)\|_{L^1(\Omega)} < \gamma \text{ then } \bar{\mu}(t) \equiv 0 \text{ in } \Omega \text{ a.e. in } (0, T), \\ \text{if } \|\bar{u}(t)\|_{L^1(\Omega)} = \gamma \text{ and } \bar{\mu}(t) \not\equiv 0 \text{ in } \Omega, \\ \text{then } \text{supp}(\bar{u}(t)) \subset \{x \in \Omega : |\bar{\mu}(x, t)| = \|\bar{\mu}(t)\|_{L^\infty(\Omega)}\}. \end{cases} \quad (2.18)$$

COROLLARY 2.3 Let $\bar{u} \in U_{ad} \cap L^\infty(\Omega)$ satisfy (2.15) and (2.18). Then, the following identities are

fulfilled

$$\begin{aligned}\bar{u}(x,t) &= -\frac{1}{\kappa} \operatorname{sign}(\bar{\varphi}(x,t)) (|\bar{\varphi}(x,t)| - \|\bar{\mu}(t)\|_{L^\infty(\Omega)})^+ \\ &= -\frac{1}{\kappa} \left\{ \left[\bar{\varphi}(x,t) + \|\bar{\mu}(t)\|_{L^\infty(\Omega)} \right]^- + \left[\bar{\varphi}(x,t) - \|\bar{\mu}(t)\|_{L^\infty(\Omega)} \right]^+ \right\}.\end{aligned}\quad (2.19)$$

Moreover, the regularity $\bar{u} \in H^1(Q)$ and $\bar{\mu} \in H^1(Q)$ hold.

We finish this section by considering the second order optimality conditions. To this end we introduce some notation. We consider the Lipschitz continuous and convex mapping $j : L^1(\Omega) \rightarrow \mathbb{R}$ defined by $j(v) = \|v\|_{L^1(\Omega)}$. Its directional derivative is given by the expression

$$j'(u;v) = \int_{\Omega_u^+} v(x) \, dx - \int_{\Omega_u^-} v(x) \, dx + \int_{\Omega_u^0} |v(x)| \, dx \quad \forall u, v \in L^1(\Omega), \quad (2.20)$$

where

$$\Omega_u^+ = \{x \in \Omega : u(x) > 0\}, \quad \Omega_u^- = \{x \in \Omega : u(x) < 0\} \text{ and } \Omega_u^0 = \Omega \setminus (\Omega_u^+ \cup \Omega_u^-).$$

Given an element $\bar{u} \in U_{ad}$ satisfying the first order optimality conditions (2.12)–(2.15), set

$$I_\gamma = \{t \in (0, T) : j(\bar{u}(t)) = \gamma\} \quad \text{and} \quad I_\gamma^+ = \{t \in I_\gamma : \bar{\mu}(t) \neq 0 \text{ in } \Omega\}.$$

Now, we define the cone of critical directions associated with \bar{u}

$$C_{\bar{u}} = \left\{ v \in L^2(Q) : J'(\bar{u})v = 0 \text{ and } j'(\bar{u}(t); v(t)) \begin{cases} = 0 & \text{if } t \in I_\gamma^+, \\ \leq 0 & \text{if } t \in I_\gamma \setminus I_\gamma^+, \end{cases} \right\}.$$

Then, we have the following theorem, whose proof can be found in Casas & Kunisch (2021).

THEOREM 2.4 Let \bar{u} be a local solution of (P) in the $L^r(0, T; L^2(\Omega))$ sense with $r > \frac{4}{4-n}$. Then, the inequality $J''(\bar{u})v^2 \geq 0$ holds for all $v \in C_{\bar{u}}$. Reciprocally, if $\bar{u} \in U_{ad}$ satisfies the first order optimality conditions and the second order condition $J''(\bar{u})v^2 > 0 \, \forall v \in C_{\bar{u}} \setminus \{0\}$, then there exist $\delta > 0$ and $\varepsilon > 0$ such that

$$J(\bar{u}) + \frac{\delta}{2} \|u - \bar{u}\|_{L^2(Q)}^2 \leq J(u) \quad \forall u \in U_{ad} \cap B_\varepsilon(\bar{u}), \quad (2.21)$$

where $B_\varepsilon(\bar{u}) = \{u \in L^r(0, T; L^2(\Omega)) : \|u - \bar{u}\|_{L^r(0, T; L^2(\Omega))} \leq \varepsilon\}$.

Given $s > 0$ we define the extended cone

$$C_{\bar{u}}^s = \left\{ v \in L^2(Q) : |J'(\bar{u})v| \leq s \|v\|_{L^2(Q)} \text{ and } \begin{cases} |j'(\bar{u}(t); v(t))| \leq s \|v\|_{L^2(Q)} & \text{if } t \in I_\gamma^+, \\ j'(\bar{u}(t); v(t)) \leq s \|v\|_{L^2(Q)} & \text{if } t \in I_\gamma \setminus I_\gamma^+, \end{cases} \right\}.$$

Then, we have the following result.

THEOREM 2.5 Let $\bar{u} \in U_{ad}$ satisfy the first order optimality conditions (2.12)–(2.15) and the second order condition $J''(\bar{u})v^2 > 0 \, \forall v \in C_{\bar{u}} \setminus \{0\}$. Then, for every $r \in (\frac{4}{4-n}, \infty]$ there exist strictly positive numbers ε, s, λ such that

$$J''(u)v^2 \geq \lambda \|v\|_{L^2(Q)}^2 \quad \forall v \in C_{\bar{u}}^s \text{ and } \forall u \in B_\varepsilon(\bar{u}), \quad (2.22)$$

where $B_\varepsilon(\bar{u})$ denotes the $L^r(0, T; L^2(\Omega))$ closed ball.

3. Numerical approximation

In this section we study the numerical discretization of (P) by discontinuous Galerkin finite element methods. To this end we consider a quasi-uniform family of triangulations $\{\mathbb{K}_h\}_{h>0}$ of $\bar{\Omega}$, cf. Brenner & Scott (2008, Definition (4.4.13)), and a quasi-uniform family of partitions of size τ of $[0, T]$, $0 = t_0 < t_1 < \dots < t_{N_\tau} = T$. We will denote by N_h and $N_{I,h}$ the number of nodes and interior nodes of the triangulation \mathbb{K}_h , $I_j = (t_{j-1}, t_j)$, $\tau_j = t_j - t_{j-1}$, $\tau = \max_{1 \leq j \leq N_\tau} \tau_j$, and $\sigma = (h, \tau)$. Following Meidner & Vexler (2018) we make the following assumptions.

Assumption 4- The next properties hold

$$\begin{aligned} &\exists \theta_1, \theta_2 > 0 \text{ such that } \tau_j \geq \theta_1 \tau^{\theta_2} \quad \forall j = 1, \dots, N_\tau, \\ &\exists \rho > 0 \text{ such that } \tau \leq \rho \tau_j \quad \forall j = 1, \dots, N_\tau, \\ &\exists \theta_3, \theta_4 > 0 \text{ and } c_{\Omega, T}, C_{\Omega, T} > 0 \text{ such that } c_{\Omega, T} h^{\theta_3} \leq \tau \leq C_{\Omega, T} h^{\theta_4}, \\ &\tau |C_a| < 1, \text{ where } C_a \text{ satisfies (2.2),} \end{aligned}$$

where the constants are independent of τ and h . Observe that θ_3 and θ_4 can be arbitrarily large and small, respectively. Hence, it is not a strong restriction.

3.1 Approximation of the state equation.

Now we consider the finite dimensional spaces

$$\begin{aligned} Y_h &= \{y_h \in C(\bar{\Omega}) : y_h|_K \in P_1(K) \quad \forall K \in \mathbb{K}_h \text{ and } y_h = 0 \text{ on } \Gamma\}, \\ \mathcal{Y}_\sigma &= \{y_\sigma \in L^2(0, T; Y_h) : y_\sigma|_{I_j} \in Y_h \quad \forall j = 1, \dots, N_\tau\}. \end{aligned}$$

The elements of \mathcal{Y}_σ can be written as

$$y_\sigma = \sum_{j=1}^{N_\tau} y_{h,j} \chi_j = \sum_{j=1}^{N_\tau} \sum_{i=1}^{N_{I,h}} y_{i,j} e_i \chi_j,$$

where $y_{h,j} \in Y_h$ for $j = 1, \dots, N_\tau$, $y_{i,j} \in \mathbb{R}$ for $i = 1, \dots, N_{I,h}$ and $j = 1, \dots, N_\tau$, $\{e_i\}_{i=1}^{N_{I,h}}$ is the nodal basis associated to the interior nodes $\{x_i\}_{i=1}^{N_{I,h}}$ of the triangulation, and χ_j denotes the characteristic function of the interval $I_j = (t_{j-1}, t_j)$.

For every $u \in L^2(Q)$, we define its associated discrete state as the unique element $y_\sigma(u) \in \mathcal{Y}_\sigma$ such that for $j = 1, \dots, N_\tau$

$$\begin{cases} \int_{\Omega} (y_{h,j} - y_{h,j-1}) z_h dx + \tau_j b(y_{h,j}, z_h) + \int_{I_j} \int_{\Omega} a(x, t, y_{h,j}) z_h dx dt = \int_{I_j} \int_{\Omega} u z_h dx dt & \forall z_h \in Y_h, \\ y_{h,0} = P_h y_0, \end{cases} \quad (3.1)$$

where $P_h : L^2(\Omega) \rightarrow Y_h$ denotes the L^2 projection operator, and $b : H^1(\Omega) \times H^1(\Omega) \rightarrow \mathbb{R}$ is the bilinear form

$$b(y, z) = \int_{\Omega} \sum_{i,j=1}^n a_{ij} \partial_{x_i} y \partial_{x_j} z dx \quad \forall y, z \in H^1(\Omega).$$

From a computational point of view, this scheme coincides with the implicit Euler discretization of the system of ordinary differential equations obtained after spatial finite element discretization. The

proof of existence and uniqueness of a solution of (3.1) is standard by using Brouwer's fixed point theorem and the assumption $\tau|C_a| < 1$. Moreover, the system (3.1) realizes an approximation of (1.1) in the following sense.

THEOREM 3.1 Let $u \in L^r(0, T; L^2(\Omega))$ hold with $r > \frac{4}{4-n}$. Under the assumptions 1, 2, and 4, there exist $h_0 > 0$, $\tau_0 > 0$, $\delta_0 > 0$, $C > 0$, and a monotone non-decreasing function $\eta_1 : [0, \infty) \rightarrow [0, \infty)$ independent of u such that for every $\tau < \tau_0$ and $h < h_0$

$$\|y_u - y_\sigma(u)\|_{L^2(Q)} \leq C(\|u\|_{L^r(0, T; L^2(\Omega))} + M_{\hat{r}, 0})(\tau + h^2), \quad (3.2)$$

$$\|y_u - y_\sigma(u)\|_{L^\infty(Q)} \leq \eta_1(\|u\|_{L^r(0, T; L^2(\Omega))} + M_{\hat{r}, 0})|\log h|^3 h^{\delta_0}, \quad (3.3)$$

where $M_{\hat{r}, 0}$ is taken as in Theorem 2.1.

Proof. For the proof of (3.2) the reader is referred to (Meidner & Vexler, 2018, Corollary 6.2). To prove (3.3) we use (Meidner & Vexler, 2018, Theorem 6.5) to deduce the existence of a constant C_1 independent of u such that

$$\|y_u - y_\sigma(u)\|_{L^\infty(Q)} \leq C_1 |\log h| \left(\log \frac{T}{\tau} \right)^2 \|y_u - z_\sigma\|_{L^\infty(Q)} \quad \forall z_\sigma \in \mathcal{Y}_\sigma.$$

Let us select a convenient z_σ . We denote by P_τ the $L^2(0, T)$ projection operator

$$P_\tau w = \sum_{j=1}^{N_\tau} \frac{1}{\tau_j} \int_{I_j} w(t) dt \chi_j \quad \forall w \in L^1(0, T).$$

It is obvious that $\|P_\tau z\|_{L^\infty(Q)} \leq \|z\|_{L^\infty(Q)}$ for every $z \in L^\infty(Q)$.

We also set $\Pi_h : C_0(\bar{\Omega}) \rightarrow Y_h$ the interpolation operator $\Pi_h z = \sum_{i=1}^{N_{I,h}} z(x_i) e_i$. Then, we take $z_\sigma = P_\tau \Pi_h y_u$. From (2.6) we get

$$\begin{aligned} \|y_u - z_\sigma\|_{L^\infty(Q)} &\leq \|y_u - P_\tau y_u\|_{L^\infty(Q)} + \|P_\tau(y_u - \Pi_h y_u)\|_{L^\infty(Q)} \\ &\leq \|y_u - P_\tau y_u\|_{L^\infty(Q)} + \|y_u - \Pi_h y_u\|_{L^\infty(Q)} \\ &\leq (\tau^\beta + (n+1)h^\beta) \|y_u\|_{C^{0,\beta}(\bar{Q})} \leq (\tau^\beta + (n+1)h^\beta) \eta(\|u\|_{L^r(0, T; L^2(\Omega))} + M_{\hat{r}, 0}). \end{aligned}$$

Using the assumption $c_{\Omega, T} h^{\theta_3} \leq \tau \leq C_{\Omega, \tau} h^{\theta_4}$ and taking $\delta_0 = \min\{1, \theta_4\} \beta$ we deduce (3.3). \square

3.2 Approximation of the control problem.

We will consider two different ways to discretize the space of controls:

I - Piecewise constant controls. We introduce the spaces and sets

$$\begin{aligned} U_h &= U_{h,0} = \{u_h \in L^\infty(\Omega) : u_{h|_K} \equiv u_K \in \mathbb{R} \quad \forall K \in \mathbb{K}_h\}, \\ B_{h,\gamma} &= \{u_h = \sum_{K \in \mathbb{K}_h} u_K \chi_K \in U_{h,0} : \sum_{K \in \mathbb{K}_h} |K| |u_K| \leq \gamma\}, \\ \mathbb{U}_\sigma &= \mathbb{U}_{\sigma,0} = \{u_\sigma = \sum_{j=1}^{N_\tau} u_{h,j} \chi_j : u_{h,j} \in U_{h,0} \text{ for } j = 1, \dots, N_\tau\}, \\ \mathbb{U}_{\sigma,ad} &= \{u_\sigma \in \mathbb{U}_{\sigma,0} : u_{h,j} \in B_{h,\gamma} \text{ for } j = 1, \dots, N_\tau\}, \end{aligned}$$

where χ_K and χ_j denote the characteristic functions of the sets K and I_j , respectively. It is immediate to check that $\mathbb{U}_{\sigma,ad} = \mathbb{U}_{\sigma} \cap U_{ad} \subset U_{ad}$.

II - Piecewise linear controls. In this case we take

$$\begin{aligned} U_h &= U_{h,1} = \{u_h \in C(\bar{\Omega}) : u_h|_K \in \mathcal{P}_1(K) \quad \forall K \in \mathbb{K}_h\}, \\ B_{h,\gamma} &= \{u_h = \sum_{i=1}^{N_h} u_i e_i \in U_{h,1} : \sum_{i=1}^{N_h} |u_i| \int_{\Omega} e_i \, dx \leq \gamma\}, \\ \mathbb{U}_{\sigma} &= \mathbb{U}_{\sigma,1} = \{u_{\sigma} = \sum_{j=1}^{N_{\tau}} u_{h,j} \chi_j : u_{h,j} \in U_{h,1} \text{ for } 1 \leq j = 1, \dots, N_{\tau}\}, \\ \mathbb{U}_{\sigma,ad} &= \left\{ u_{\sigma} \in \mathbb{U}_{\sigma,1} : u_{h,j} \in B_{h,\gamma} \text{ for } j = 1, \dots, N_{\tau} \right\}, \end{aligned}$$

where $\mathcal{P}_1(K)$ denotes the space of the polynomials on K of degree ≤ 1 . From the inequality

$$\|u_h\|_{L^1(\Omega)} = \int_{\Omega} \left| \sum_{i=1}^{N_h} u_i e_i \right| \, dx \leq \sum_{i=1}^{N_h} |u_i| \int_{\Omega} e_i \, dx$$

we infer that $\mathbb{U}_{\sigma,ad} \subset U_{ad}$.

We observe that $\mathbb{U}_{\sigma} \subset L^{\infty}(0, T; U_h)$ in both cases and every element $u_{\sigma} \in \mathbb{U}_{\sigma}$ can be written in form

$$u_{\sigma} = \sum_{j=1}^{N_{\tau}} u_{h,j} \chi_j = \begin{cases} \sum_{j=1}^{N_{\tau}} \sum_{K \in \mathbb{K}_h} u_{K,j} \chi_K \chi_j & \text{if } \mathbb{U}_{\sigma} = \mathbb{U}_{\sigma,0}, \\ \sum_{j=1}^{N_{\tau}} \sum_{i=1}^{N_h} u_{i,j} e_i \chi_j & \text{if } \mathbb{U}_{\sigma} = \mathbb{U}_{\sigma,1}. \end{cases}$$

Now, we formulate the discrete control problem

$$(P_{\sigma}) \quad \inf_{u_{\sigma} \in \mathbb{U}_{\sigma,ad}} J_{\sigma}(u_{\sigma}) := \frac{1}{2} \int_Q |y_{\sigma}(u_{\sigma}) - y_d|^2 \, dx \, dt + \frac{\kappa}{2} \|u_{\sigma}\|_{\sigma}^2,$$

where $y_{\sigma}(u_{\sigma})$ is the solution of (3.1) for $u = u_{\sigma}$ and

$$\|u_{\sigma}\|_{\sigma}^2 = \sum_{j=1}^{N_{\tau}} \tau_j \|u_{h,j}\|_h^2$$

with $\|\cdot\|_h$ the norm in U_h defined by

$$\|u_h\|_h = \begin{cases} \left(\sum_{K \in \mathbb{K}_h} |K| u_K^2 \right)^{\frac{1}{2}} & \text{if } U_h = U_{h,0}, \\ \left(\sum_{i=1}^{N_h} \left(\int_{\Omega} e_i(x) \, dx \right) u_i^2 \right)^{\frac{1}{2}} & \text{if } U_h = U_{h,1}. \end{cases}$$

We notice that

$$\|u_h\|_{L^2(\Omega)}^2 = \begin{cases} \sum_{K \in \mathbb{K}_h} \int_K |u_K|^2 dx = \|u_h\|_h^2 & \text{if } U_h = U_{h,0}, \\ \int_{\Omega} \left(\sum_{i=1}^{N_h} u_i e_i(x) \right)^2 dx \leq \int_{\Omega} \left(\sum_{i=1}^{N_h} e_i(x) u_i^2 \right) dx = \|u_h\|_h^2 & \text{if } U_h = U_{h,1}, \end{cases} \quad (3.4)$$

where we have used that $0 \leq e_i(x) \leq 1$ and $\sum_{i=1}^{N_h} e_i(x) = 1$ in Ω .

We also introduce $\|u_{\sigma}\|_{\sigma} = \sqrt{(u_{\sigma}, u_{\sigma})_{\sigma}}$, where the scalar product $(\cdot, \cdot)_{\sigma}$ in \mathbb{U}_{σ} is defined by

$$(u_{\sigma}, v_{\sigma})_{\sigma} = \sum_{j=1}^{N_{\tau}} \tau_j (u_{h,j}, v_{h,j})_h = \begin{cases} (u_{\sigma}, v_{\sigma})_{L^2(Q)} = \sum_{j=1}^{N_{\tau}} \sum_{K \in \mathbb{K}_h} \tau_j |K| u_{K,j} v_{K,j} & \text{if } \mathbb{U}_{\sigma} = \mathbb{U}_{\sigma,0}, \\ \sum_{j=1}^{N_{\tau}} \sum_{i=1}^{N_h} \tau_j \left(\int_{\Omega} e_i(x) dx \right) u_{i,j} v_{i,j} & \text{if } \mathbb{U}_{\sigma} = \mathbb{U}_{\sigma,1}. \end{cases}$$

Due to the compactness of $\mathbb{U}_{\sigma,ad}$ in both definitions and the continuity of J_{σ} , we infer the existence of at least one solution for (P $_{\sigma}$).

Analogously to Corollary 2.1 we have the following differentiability result.

THEOREM 3.2 The functional $J_{\sigma} : \mathbb{U}_{\sigma} \rightarrow \mathbb{R}$ is of class C^2 and its first derivative is given by the expression

$$J'_{\sigma}(u_{\sigma})v_{\sigma} = \int_Q \varphi_{\sigma} v_{\sigma} dx dt + \kappa(u_{\sigma}, v_{\sigma})_{\sigma}, \quad (3.5)$$

where $\varphi_{\sigma}(u_{\sigma}) \in \mathcal{B}_{\sigma}$ is the solution of the adjoint state equation: for $j = N_{\tau}, \dots, 1$

$$\begin{cases} \int_{\Omega} (\varphi_{h,j} - \varphi_{h,j+1}) z_h dx + \tau_j b(z_h, \varphi_{h,j}) + \int_{I_j} \int_{\Omega} \frac{\partial a}{\partial y}(x, t, y_{\sigma}(u_{\sigma})) \varphi_{h,j} z_h dx dt \\ \quad = \int_{I_j} \int_{\Omega} (y_{\sigma}(u_{\sigma}) - y_d) z_h dx dt \quad \forall z_h \in Y_h, \\ \varphi_{h, N_{\tau}+1} = 0. \end{cases} \quad (3.6)$$

Now we compare the continuous and discrete adjoint states.

THEOREM 3.3 Let $u \in L^r(0, T; L^2(\Omega))$ hold with $r > \frac{4}{4-n}$, and let us denote by φ_u and $\varphi_{\sigma}(u)$ the solutions of (2.11) and (3.6) with $y_{\sigma}(u_{\sigma})$ replaced by $y_{\sigma}(u)$. Under the assumptions 1–4, and taking h_0 and τ_0 as in Theorem 3.1, there exists a monotone non-decreasing function $\eta_2 : [0, \infty) \rightarrow \mathbb{R}$ independent of u such that for every $\tau < \tau_0$ and $h < h_0$

$$\|\varphi_u - \varphi_{\sigma}(u)\|_{L^2(Q)} \leq \eta_2(\|u\|_{L^r(0, T; L^2(\Omega))} + M_{\hat{r}, 0})(\tau + h^2), \quad (3.7)$$

$$\|\varphi_u - \varphi_{\sigma}(u)\|_{L^{\infty}(Q)} \leq \eta_2(\|u\|_{L^r(0, T; L^2(\Omega))} + M_{\hat{r}, 0}) |\log h|^3 h^{\delta_0}. \quad (3.8)$$

Proof. Let $\psi_u \in C^{0, \beta}(\bar{Q}) \cap H^{2,1}(Q)$ denote the solution of the adjoint state equation

$$\begin{cases} -\frac{\partial \psi_u}{\partial t} + A^* \psi_u + \frac{\partial a}{\partial y}(x, t, y_{\sigma}(u)) \psi_u = y_{\sigma}(u) - y_d & \text{in } Q, \\ \psi_u = 0 & \text{on } \Sigma, \quad \psi_u(T) = 0 & \text{in } \Omega, \end{cases} \quad (3.9)$$

and set $\varphi_u - \varphi_\sigma(u) = (\varphi_u - \psi_u) + (\psi_u - \varphi_\sigma(u)) = e_u + \xi_u$. Subtracting the equations (2.11) and (3.9) we get

$$\begin{cases} -\frac{\partial e_u}{\partial t} + A^* e_u + \frac{\partial a}{\partial y}(x, t, y_u) e_u = (y_u - y_\sigma(u)) + \left[\frac{\partial a}{\partial y}(x, t, y_\sigma(u)) - \frac{\partial a}{\partial y}(x, t, y_u) \right] \psi_u & \text{in } \mathcal{Q}, \\ e_u = 0 \text{ on } \Sigma, \quad e_u(T) = 0 & \text{in } \Omega. \end{cases} \quad (3.10)$$

Setting $M = \|y_u\|_{L^\infty(\mathcal{Q})} + 1$ and taking h_0 and τ_0 small enough, we infer from (3.3) that $\|y_\sigma(u)\|_{L^\infty(\mathcal{Q})} \leq M$ for every $\sigma = (h, \tau)$ with $h \leq h_0$ and $\tau \leq \tau_0$. Then, from (3.9) it follows with (2.4) that

$$\|\psi_u\|_{L^\infty(\mathcal{Q})} \leq C_1 (\|y_\sigma(u)\|_{L^\infty(\mathcal{Q})} + \|y_d\|_{L^{\hat{r}}(0, T; L^2(\Omega))}) \leq C_1 (M + \|y_d\|_{L^{\hat{r}}(0, T; L^2(\Omega))}). \quad (3.11)$$

From (2.4), (3.3), and the mean value theorem we obtain

$$\left| \frac{\partial a}{\partial y}(x, t, y_\sigma(u)(x, t)) - \frac{\partial a}{\partial y}(x, t, y_u(x, t)) \right| \leq C_{a, M} |y_\sigma(u)(x, t) - y_u(x, t)|. \quad (3.12)$$

From (3.10), (3.11), (3.12), and (3.2), we infer

$$\begin{aligned} \|e_u\|_{L^2(\mathcal{Q})} &\leq C_2 [1 + C_{a, M} C_1 (M + \|y_d\|_{L^{\hat{r}}(0, T; L^2(\Omega))})] \|y_\sigma(u) - y_u\|_{L^2(\mathcal{Q})} \\ &\leq C_M C (\|u\|_{L^{\hat{r}}(0, T; L^2(\Omega))} + M_{\hat{r}, 0}) (\tau + h^2). \end{aligned} \quad (3.13)$$

The constant C_M is a monotone non-decreasing function of M .

Let us estimate ξ_u . Since $\varphi_\sigma(u)$ is the solution of the discretization of the linear equation (3.8), the classical error estimates yield the existence of a constant C_3 such that

$$\|\xi_u\|_{L^2(\mathcal{Q})} \leq C_2 (\tau + h^2) (\|y_\sigma(u)\|_{L^2(\mathcal{Q})} + \|y_d\|_{L^2(\mathcal{Q})}). \quad (3.14)$$

Hence, (3.13) and (3.14) along with (2.6) lead to (3.7).

To prove (3.8) we first modify (3.13) as follows

$$\begin{aligned} \|e_u\|_{L^\infty(\mathcal{Q})} &\leq C_2 [1 + C_{a, M} C_1 (M + \|y_d\|_{L^{\hat{r}}(0, T; L^2(\Omega))})] \|y_\sigma(u) - y_u\|_{L^\infty(\mathcal{Q})} \\ &\leq C_M \eta_1 (\|u\|_{L^{\hat{r}}(0, T; L^2(\Omega))} + M_{\hat{r}, 0}) |\log h|^3 h^{\delta_0}. \end{aligned} \quad (3.15)$$

Finally, using the linearity of the equations satisfied by ψ_u and arguing as for the estimate (3.3) we infer

$$\|\xi_u\|_{L^\infty(\mathcal{Q})} \leq C_3 |\log h|^3 h^{\delta_0}.$$

The last inequality and (3.15) imply (3.8). \square

3.3 First order optimality conditions.

The goal of this subsection is to prove the first order optimality conditions and their consequences.

THEOREM 3.4 Let \bar{u}_σ be a local minimum of (P_σ) . Then there exist $\bar{y}_\sigma, \bar{\varphi}_\sigma \in \mathcal{Y}_\sigma$ and $\bar{\mu}_\sigma \in \mathbb{U}_\sigma$ such

that

$$\left\{ \begin{array}{l} \int_{\Omega} (\bar{y}_{h,j} - \bar{y}_{h,j-1}) z_h dx + \tau_j b(\bar{y}_{h,j}, z_h) + \int_{I_j} \int_{\Omega} a(x, t, \bar{y}_{h,j}) z_h dx dt \\ \qquad \qquad \qquad = \int_{I_j} \int_{\Omega} \bar{u}_{h,j} z_h dx dt \quad \forall z_h \in Y_h \text{ and } \forall j = 1, \dots, N_{\tau}, \\ \bar{y}_{h,0} = P_h y_0, \end{array} \right. \quad (3.16)$$

$$\left\{ \begin{array}{l} \int_{\Omega} (\bar{\varphi}_{h,j} - \bar{\varphi}_{h,j+1}) z_h dx + \tau_j b(z_h, \bar{\varphi}_{h,j}) + \int_{I_j} \int_{\Omega} \frac{\partial a}{\partial y}(x, t, y_{\sigma}(u_{\sigma})) \bar{\varphi}_{h,j} z_h dx dt \\ \qquad \qquad \qquad = \int_{I_j} \int_{\Omega} (\bar{y}_{h,j} - y_d) z_h dx dt \quad \forall z_h \in Y_h \text{ and } \forall j = N_{\tau}, \dots, 1, \\ \bar{\varphi}_{h, N_{\tau}+1} = 0, \end{array} \right. \quad (3.17)$$

$$(\bar{\mu}_{\sigma}, u_{\sigma} - \bar{u}_{\sigma})_{\sigma} \leq 0 \quad \forall u_{\sigma} \in \mathbb{U}_{\sigma, ad}, \quad (3.18)$$

$$\left\{ \begin{array}{l} \frac{1}{|K|} \int_K \bar{\varphi}_{h,j} dx + \kappa \bar{u}_{K,j} + \bar{\mu}_{K,j} = 0 \quad \forall K \in \mathbb{K}_h \text{ and } \forall j = 1, \dots, N_{\tau}, \quad \text{if } \mathbb{U}_{\sigma} = \mathbb{U}_{\sigma,0}, \\ \frac{1}{\int_{\Omega} e_i dx} \int_{\Omega} \bar{\varphi}_{h,j} e_i dx + \kappa \bar{u}_{i,j} + \bar{\mu}_{i,j} = 0 \quad \forall i = 1, \dots, N_h \text{ and } \forall j = 1, \dots, N_{\tau}, \quad \text{if } \mathbb{U}_{\sigma} = \mathbb{U}_{\sigma,1}. \end{array} \right. \quad (3.19)$$

Proof. Taking \bar{y}_{σ} and $\bar{\varphi}_{\sigma}$ as solutions of (3.16) and (3.17), respectively, and using the convexity of $\mathbb{U}_{\sigma, ad}$ we infer with (3.5)

$$\int_Q \bar{\varphi}_{\sigma} (u_{\sigma} - \bar{u}_{\sigma}) dx dt + \kappa (\bar{u}_{\sigma}, u_{\sigma} - \bar{u}_{\sigma})_{\sigma} = J'_{\sigma}(\bar{u}_{\sigma})(u_{\sigma} - \bar{u}_{\sigma}) \geq 0 \quad \forall u_{\sigma} \in \mathbb{U}_{\sigma, ad}. \quad (3.20)$$

Now we distinguish the cases $\mathbb{U}_{\sigma} = \mathbb{U}_{\sigma,0}$ and $\mathbb{U}_{\sigma} = \mathbb{U}_{\sigma,1}$.

Case $\mathbb{U}_{\sigma} = \mathbb{U}_{\sigma,0}$. In this case, (3.20) can be written as follows

$$\sum_{j=1}^{N_{\tau}} \sum_{K \in \mathbb{K}_h} \tau_j \left(\int_K \bar{\varphi}_{h,j} dx + \kappa |K| \bar{u}_{K,j} \right) (u_{K,j} - \bar{u}_{K,j}) \geq 0 \quad \forall u_{\sigma} \in \mathbb{U}_{\sigma, ad}. \quad (3.21)$$

Then, defining

$$\bar{\mu}_{\sigma} = \sum_{j=1}^{N_{\tau}} \sum_{K \in \mathbb{K}_h} \bar{\mu}_{K,j} \chi_K \chi_j \quad \text{with} \quad \bar{\mu}_{K,j} = - \left(\frac{1}{|K|} \int_K \bar{\varphi}_{h,j} dx + \kappa \bar{u}_{K,j} \right),$$

we have that the first identity of (3.19) holds. Inequalities (3.18) are consequence of (3.21):

$$\begin{aligned} (\bar{\mu}_{\sigma}, u_{\sigma} - \bar{u}_{\sigma})_{\sigma} &= \sum_{j=1}^{N_{\tau}} \sum_{K \in \mathbb{K}_h} \tau_j |K| \bar{\mu}_{K,j} (u_{K,j} - \bar{u}_{K,j}) \\ &= - \sum_{j=1}^{N_{\tau}} \sum_{K \in \mathbb{K}_h} \tau_j \left(\int_K \bar{\varphi}_{h,j} dx + \kappa |K| \bar{u}_{K,j} \right) (u_{K,j} - \bar{u}_{K,j}) \leq 0 \quad \forall u_{\sigma} \in \mathbb{U}_{\sigma, ad}. \end{aligned}$$

Case $\mathbb{U}_{\sigma} = \mathbb{U}_{\sigma,1}$. From (3.20) and using the definition of $(\cdot, \cdot)_{\sigma}$ we deduce

$$\sum_{j=1}^{N_{\tau}} \sum_{i=1}^{N_h} \tau_j \left(\int_{\Omega} \bar{\varphi}_{h,j} e_i dx + \kappa \left(\int_{\Omega} e_i dx \right) \bar{u}_{i,j} \right) (u_{i,j} - \bar{u}_{i,j}) \geq 0 \quad \forall u_{\sigma} \in \mathbb{U}_{\sigma, ad}. \quad (3.22)$$

Now we set

$$\bar{\mu}_\sigma = \sum_{j=1}^{N_\tau} \sum_{i=1}^{N_h} \bar{\mu}_{i,j} e_i \chi_j \quad \text{with} \quad \bar{\mu}_{i,j} = - \left(\frac{1}{\int_\Omega e_i \, dx} \int_\Omega \bar{\varphi}_{h,j} e_i \, dx + \kappa \bar{u}_{i,j} \right).$$

Then, the second identity of (3.19) is satisfied. We finish the proof by checking (3.18) with the aid of (3.22)

$$\begin{aligned} (\bar{\mu}_\sigma, u_\sigma - \bar{u}_\sigma)_\sigma &= \sum_{j=1}^{N_\tau} \sum_{i=1}^{N_h} \tau_j \left(\int_\Omega e_i \, dx \right) \bar{\mu}_{i,j} (u_{i,j} - \bar{u}_{i,j}) \\ &= - \sum_{j=1}^{N_\tau} \sum_{i=1}^{N_h} \left(\tau_j \int_\Omega \bar{\varphi}_{h,j} e_i \, dx + \kappa \left(\int_\Omega e_i \, dx \right) \bar{u}_{i,j} \right) (u_{i,j} - \bar{u}_{i,j}) \leq 0 \quad \forall u_\sigma \in \mathbb{U}_{\sigma,ad}. \end{aligned}$$

□

Let us introduce the following notation:

$$\|u_h\|_{l^\infty} = \begin{cases} \max_{K \in \mathbb{K}_h} |u_K| & \text{if } U_h = U_{h,0}, \\ \max_{1 \leq i \leq N_h} |u_i| & \text{if } U_h = U_{h,1}, \end{cases}$$

and $j_h : U_h \rightarrow \mathbb{R}$ is the functional defined by

$$j_h(u_h) = \begin{cases} \sum_{K \in \mathbb{K}_h} |K| |u_K| & \text{if } U_h = U_{h,0}, \\ \sum_{i=1}^{N_h} |u_i| \int_\Omega e_i \, dx & \text{if } U_h = U_{h,1}. \end{cases}$$

We have the following corollary.

COROLLARY 3.1 Let \bar{u}_σ , $\bar{\mu}_\sigma$, and $\bar{\varphi}_\sigma$ satisfy (3.17)–(3.19), and assume that $\bar{u}_\sigma \in \mathbb{U}_{\sigma,ad}$. Then, the following properties hold for every $j = 1, \dots, N_\tau$

$$(\bar{\mu}_{h,j}, u_h - \bar{u}_{h,j})_h \leq 0 \quad \forall u_h \in B_{h,\gamma}, \quad (3.23)$$

if $\mathbb{U}_\sigma = \mathbb{U}_{\sigma,0}$ then

$$\begin{cases} \bar{\mu}_{K,j} \bar{u}_{K,j} = |\bar{\mu}_{K,j}| |\bar{u}_{K,j}| \quad \forall K \in \mathbb{K}_h, \\ \text{if } j_h(\bar{u}_{h,j}) < \gamma \text{ then } \bar{\mu}_{h,j} = 0, \\ \text{if } j_h(\bar{u}_{h,j}) = \gamma \text{ and } \bar{\mu}_{h,j} \neq 0, \text{ then if } \bar{u}_{K,j} \neq 0 \Rightarrow |\bar{\mu}_{K,j}| = \|\bar{\mu}_{h,j}\|_{l^\infty}. \end{cases} \quad (3.24)$$

if $\mathbb{U}_\sigma = \mathbb{U}_{\sigma,1}$ then

$$\begin{cases} \bar{\mu}_{h,j} \bar{u}_{h,j} = |\bar{\mu}_{h,j}| |\bar{u}_{h,j}|, \\ \text{if } j_h(\bar{u}_{h,j}) < \gamma \text{ then } \bar{\mu}_{h,j} = 0, \\ \text{if } j_h(\bar{u}_{h,j}) = \gamma \text{ and } \bar{\mu}_{h,j} \neq 0, \text{ then if } \bar{u}_{i,j} \neq 0 \Rightarrow |\bar{\mu}_{i,j}| = \|\bar{\mu}_{h,j}\|_{l^\infty}. \end{cases} \quad (3.25)$$

Proof. Given $1 \leq j \leq N_\tau$ and $u_h \in B_{h,\gamma}$ we define

$$u_\sigma = \sum_{l=1}^{N_\tau} u_{h,l} \chi_l \quad \text{with} \quad u_{h,l} = \begin{cases} \bar{u}_{h,l} & \text{if } l \neq j, \\ u_h & \text{if } l = j. \end{cases}$$

Then, $u_\sigma \in \mathbb{U}_{\sigma,ad}$ and (3.18) implies

$$\tau_j(\bar{\mu}_{h,j}, u_h - \bar{u}_{h,j})_h = (\bar{\mu}_\sigma, u_\sigma - \bar{u}_\sigma)_\sigma \leq 0,$$

which proves (3.23). The rest of the proof is divided into two cases.

Case $\mathbb{U}_\sigma = \mathbb{U}_{\sigma,0}$. For

$$u_h = \sum_{K' \in \mathbb{K}_h} u_{K'} \chi_{K'} \quad \text{with} \quad u_{K'} = \begin{cases} \bar{u}_{K',j} & \text{if } K' \neq K, \\ 0 & \text{if } K' = K, \end{cases}$$

(3.23) leads to $|K| \bar{\mu}_{K,j} \bar{u}_{K,j} \geq 0$, which implies the first identity of (3.24). To establish the second statement of (3.24), given $K \in \mathbb{K}_h$ arbitrary, we define

$$u_h^\pm = \sum_{K' \in \mathbb{K}_h} u_{K'} \chi_{K'} \quad \text{with} \quad u_{K'} = \begin{cases} \bar{u}_{K',j} & \text{if } K' \neq K, \\ \bar{u}_{K,j} \pm \varepsilon & \text{if } K' = K. \end{cases}$$

Then, for ε small enough, due to the fact that $j(\bar{u}_{h,j}) < \gamma$, we have that $u_h^\pm \in B_{h,\gamma}$. Then, (3.23) leads to $\pm |K| \bar{\mu}_{K,j} \varepsilon \geq 0$, which implies that $\bar{\mu}_{K,j} = 0$ for every $K \in \mathbb{K}_h$.

Now, we assume that $j_h(\bar{u}_{h,j}) = \gamma$ and $\bar{\mu}_{h,j} \neq 0$. Let $K^0 \in \mathbb{K}_h$ be such that $|\bar{\mu}_{K^0,j}| = \max_{K' \in \mathbb{K}_h} |\bar{\mu}_{K',j}|$. If $\bar{u}_{K,j} \neq 0$ we define

$$u_h = \sum_{K' \in \mathbb{K}_h} u_{K'} \chi_{K'} \quad \text{with} \quad u_{K'} = \begin{cases} \bar{u}_{K,j} - \frac{\varepsilon}{|K|} \text{sign}(\bar{u}_{K,j}) & \text{if } K' = K, \\ \bar{u}_{K^0,j} + \frac{\varepsilon}{|K^0|} \text{sign}(\bar{u}_{K^0,j}) & \text{if } K' = K^0, \\ \bar{u}_{K,j} & \text{otherwise,} \end{cases}$$

where $0 < \varepsilon < |K| |\bar{u}_{K,j}|$. Then, $j_h(u_h) = j_h(\bar{u}_{h,j}) = \gamma$. Hence, $u_h \in B_{h,\gamma}$ and we get with (3.23) and the first statement of (3.24)

$$\varepsilon |\bar{\mu}_{K^0,j}| - \varepsilon |\bar{\mu}_{K,j}| = (\bar{\mu}_{h,j}, u_h - \bar{u}_{h,j})_h \leq 0.$$

This proves the last statement of (3.24).

Case $\mathbb{U}_\sigma = \mathbb{U}_{\sigma,1}$. Let $1 \leq i \leq N_h$ arbitrary and set

$$u_h = \sum_{i'=1}^{N_h} u_{i'} e_{i'} \quad \text{with} \quad u_{i'} = \begin{cases} \bar{u}_{i',j} & \text{if } i' \neq i, \\ 0 & \text{if } i' = i. \end{cases}$$

Then (3.23) implies

$$-\left(\int_{\Omega} e_i \, dx \right) \bar{\mu}_{i,j} \bar{u}_{i,j} = (\bar{\mu}_{h,j}, u_h - \bar{u}_{h,j})_h \leq 0,$$

which proves the first statement of (3.25).

To establish the second statement of (3.25), given $1 \leq i \leq N_h$ arbitrary, we define

$$u_h^\pm = \sum_{i'=1}^{N_h} u_{i'} e_{i'} \quad \text{with} \quad u_{i'} = \begin{cases} \bar{u}_{i',j} & \text{if } i' \neq i, \\ \bar{u}_{i,j} \pm \varepsilon & \text{if } i' = i. \end{cases}$$

Then, for ε small enough, due to the fact that $j(\bar{u}_{h,j}) < \gamma$, we have that $u_h^\pm \in B_{h,\gamma}$. Then, (3.23) leads to $\pm \left(\int_{\Omega} e_i \, dx \right) \bar{\mu}_{i,j} \varepsilon \geq 0$, which implies that $\bar{\mu}_{i,j} = 0$ for every $i = 1, \dots, N_h$.

Finally, we assume that $j_h(\bar{u}_{h,j}) = \gamma$ and $\bar{\mu}_{h,j} \neq 0$. Let $1 \leq i^0 \leq N_h$ be such that $|\bar{\mu}_{i^0,j}| = \max_{1 \leq i' \leq N_h} |\bar{\mu}_{i',j}|$. If $\bar{u}_{i,j} \neq 0$ we define

$$u_h = \sum_{i'=1}^{N_h} u_{i'} e_{i'} \quad \text{with} \quad u_{i'} = \begin{cases} \bar{u}_{i,j} - \frac{\varepsilon}{\int_{\Omega} e_i \, dx} \text{sign}(\bar{u}_{i,j}) & \text{if } i' = i, \\ \bar{u}_{i^0,j} + \frac{\varepsilon}{\int_{\Omega} e_{i^0} \, dx} \text{sign}(\bar{u}_{i^0,j}) & \text{if } i' = i^0, \\ \bar{u}_{i,j} & \text{otherwise,} \end{cases}$$

where $0 < \varepsilon < (\int_{\Omega} e_i \, dx) |\bar{u}_{i,j}|$. Then, $j_h(u_h) = j_h(\bar{u}_{h,j}) = \gamma$. Hence, $u_h \in B_{h,\gamma}$ and we get with (3.23) and the first statement of (3.25)

$$\varepsilon |\bar{\mu}_{i^0,j}| - \varepsilon |\bar{\mu}_{i,j}| = (\bar{\mu}_{h,j}, u_h - \bar{u}_{h,j})_h \leq 0.$$

This proves the last statement of (3.25). \square

COROLLARY 3.2 Let $\bar{u}_{\sigma} \in \mathbb{U}_{\sigma,ad}$ satisfy (3.19) and (3.24) or (3.25). Then, the following identities hold for every $j = 1, \dots, N_{\tau}$

$$\begin{aligned} \text{if } \mathbb{U}_{\sigma} = \mathbb{U}_{\sigma,0} \text{ then } \bar{u}_{K,j} &= -\frac{1}{\kappa} \text{sign} \left(\int_K \bar{\varphi}_{h,j} \, dx \right) \left(\frac{1}{|K|} \left| \int_K \bar{\varphi}_{h,j} \, dx \right| - \|\bar{\mu}_{h,j}\|_{l^{\infty}} \right)^+ \\ &= -\frac{1}{\kappa} \left\{ \left[\frac{1}{|K|} \int_K \bar{\varphi}_{h,j} \, dx + \|\bar{\mu}_{h,j}\|_{l^{\infty}} \right]^- + \left[\frac{1}{|K|} \int_K \bar{\varphi}_{h,j} \, dx - \|\bar{\mu}_{h,j}\|_{l^{\infty}} \right]^+ \right\}, \end{aligned} \quad (3.26)$$

$$\begin{aligned} \text{if } \mathbb{U}_{\sigma} = \mathbb{U}_{\sigma,1} \text{ then } \bar{u}_{i,j} &= -\frac{1}{\kappa} \text{sign} \left(\int_{\Omega} \bar{\varphi}_{h,j} e_i \, dx \right) \left(\frac{1}{\int_{\Omega} e_i \, dx} \left| \int_{\Omega} \bar{\varphi}_{h,j} e_i \, dx \right| - \|\bar{\mu}_{h,j}\|_{l^{\infty}} \right)^+ \\ &= -\frac{1}{\kappa} \left\{ \left[\frac{1}{\int_{\Omega} e_i \, dx} \int_{\Omega} \bar{\varphi}_{h,j} e_i \, dx + \|\bar{\mu}_{h,j}\|_{l^{\infty}} \right]^- + \left[\frac{1}{\int_{\Omega} e_i \, dx} \int_{\Omega} \bar{\varphi}_{h,j} e_i \, dx - \|\bar{\mu}_{h,j}\|_{l^{\infty}} \right]^+ \right\}. \end{aligned} \quad (3.27)$$

Moreover, the following sparsity property is fulfilled for every $j = 1, \dots, N_{\tau}$

$$\text{if } \mathbb{U}_{\sigma} = \mathbb{U}_{\sigma,0} \text{ then } \bar{u}_{K,j} = 0 \Leftrightarrow \frac{1}{|K|} \left| \int_K \bar{\varphi}_{h,j} \, dx \right| \leq \|\bar{\mu}_{h,j}\|_{l^{\infty}}, \quad \forall K \in \mathbb{K}_h, \quad (3.28)$$

$$\text{if } \mathbb{U}_{\sigma} = \mathbb{U}_{\sigma,1} \text{ then } \bar{u}_{i,j} = 0 \Leftrightarrow \frac{1}{\int_{\Omega} e_i \, dx} \left| \int_{\Omega} \bar{\varphi}_{h,j} e_i \, dx \right| \leq \|\bar{\mu}_{h,j}\|_{l^{\infty}}, \quad \forall i = 1, \dots, N_h, \quad (3.29)$$

Proof. Let us prove the first identity of (3.26). If $\|\bar{\mu}_{h,j}\|_{l^{\infty}} = 0$, then (3.19) implies that

$$\bar{u}_{K,j} = -\frac{1}{\kappa |K|} \int_K \bar{\varphi}_{h,j} \, dx \quad \forall K \in \mathbb{K}_h,$$

which coincides with (3.26). Assume that $\|\bar{\mu}_{h,j}\|_{l^{\infty}} \neq 0$. Then, from (3.24) we deduce that $j_h(\bar{u}_{h,j}) = \gamma$. Then, the third statement of (3.24) implies that $|\bar{u}_{K,j}| = \|\bar{\mu}_{h,j}\|_{l^{\infty}}$ if $\bar{u}_{K,j} \neq 0$. Now, we distinguish three cases.

i) If $\bar{u}_{K,j} > 0$, (3.19) and the first statement of (3.24) lead to

$$\bar{u}_{K,j} = -\frac{1}{\kappa} \left\{ \frac{1}{|K|} \int_K \bar{\varphi}_{h,j} \, dx + \|\bar{\mu}_{h,j}\|_{l^{\infty}} \right\},$$

which coincides with (3.26). Indeed, observe that (3.24) and the positivity of $\bar{u}_{K,j}$ imply $\bar{\mu}_{K,j} \geq 0$. Hence, we conclude with (3.19) that $\int_K \bar{\varphi}_{h,j} dx < 0$.

ii) If $\bar{u}_{K,j} = 0$, using again (3.19) we get

$$\left| \frac{1}{|K|} \int_K \bar{\varphi}_{h,j} dx \right| = |\bar{\mu}_{K,j}| \leq \|\bar{\mu}_{h,j}\|_{l^\infty}.$$

Then, the identity (3.26) holds.

iii) If $\bar{u}_{K,j} < 0$, from the first statement of (3.24) and (3.19) we infer that

$$\bar{u}_{K,j} = -\frac{1}{\kappa} \left\{ \frac{1}{|K|} \int_K \bar{\varphi}_{h,j} dx - \|\bar{\mu}_{h,j}\|_{l^\infty} \right\}.$$

Moreover, arguing as in the case i), we deduce that $\int_K \bar{\varphi}_{h,j} dx > 0$. Hence, (3.26) holds too.

The second identity of (3.26) is obvious. Following the same arguments as above, (3.27) is proved. Finally, (3.28) and (3.29) are immediate consequences of (3.26) and (3.27), respectively. \square

4. Convergence analysis and error estimates

There are two goals in this section. First we prove that the discrete problems (P_σ) provide an approximation of (P) . Second we establish error estimates in terms of $\sigma = (h, \tau)$ for the difference between the discrete and continuous optimal controls.

THEOREM 4.1 For every σ let \bar{u}_σ be a solution of (P_σ) . Then, there exists $\sigma_0 = (h_0, \tau_0)$ such that the family $\{\bar{u}_\sigma\}_\sigma$ with $h < h_0$ and $\tau < \tau_0$ is bounded in $L^\infty(Q)$. If $\bar{u}_\sigma \xrightarrow{*} \bar{u}$ in $L^\infty(Q)$ for a sequence of σ converging to zero, we have that \bar{u} is a solution of (P) , and the following convergence properties hold

$$\lim_{\sigma \rightarrow 0} \|\bar{u}_\sigma - \bar{u}\|_{L^r(0,T;L^2(\Omega))} = 0 \quad \forall r \in [1, \infty) \quad \text{and} \quad \lim_{\sigma \rightarrow 0} J_\sigma(\bar{u}_\sigma) = J(\bar{u}). \quad (4.1)$$

To prove this theorem we need the following stability property for the solution of the system (3.1).

LEMMA 4.1 Let us assume that $4|C_a|\tau < 1$. Then, given $u \in L^2(Q)$ and denoting by $y_\sigma \in \mathcal{Y}_\sigma$ the solution of (3.1), we have the stability estimate

$$\|y_\sigma\|_{L^\infty(0,T;L^2(\Omega))} + \|y_\sigma\|_{L^2(0,T;H_0^1(\Omega))} \leq C(\|u - a(\cdot, \cdot, 0)\|_{L^2(Q)} + \|y_0\|_{L^2(\Omega)}) \quad (4.2)$$

for some constant C independent of u .

Proof. For $j = 1, \dots, N_\tau$ we take $z_h = y_{h,j}$ in (3.1), which leads to

$$\begin{aligned} & \int_\Omega (y_{h,j} - y_{h,j-1})y_{h,j} dx + \tau_j b(y_{h,j}, y_{h,j}) + \int_{I_j} \int_\Omega [a(x, t, y_{h,j}) - a(x, t, 0)]y_{h,j} dx dt \\ & = \int_{I_j} \int_\Omega [u - a(x, t, 0)]y_{h,j} dx dt. \end{aligned}$$

Using (2.1) and (2.2) along with Young's inequality we deduce from the above identity

$$\begin{aligned} & \frac{1}{2} \|y_{h,j}\|_{L^2(\Omega)}^2 + \frac{1}{2} \|y_{h,j} - y_{h,j-1}\|_{L^2(\Omega)}^2 - \frac{1}{2} \|y_{h,j-1}\|_{L^2(\Omega)}^2 + \tau_j \Lambda_A \|y_{h,j}\|_{H_0^1(\Omega)}^2 + C_a \tau_j \|y_{h,j}\|_{L^2(\Omega)}^2 \\ & \leq \|u - a(\cdot, \cdot, 0)\|_{L^2(\Omega \times I_j)} \sqrt{\tau_j} \|y_{h,j}\|_{L^2(\Omega)} \leq C_1 \|u - a(\cdot, \cdot, 0)\|_{L^2(\Omega \times I_j)}^2 + \tau_j \frac{\Lambda_A}{2} \|y_{h,j}\|_{H_0^1(\Omega)}^2. \end{aligned}$$

From here we infer

$$\|y_{h,j}\|_{L^2(\Omega)}^2 + \tau_j \Lambda_A \|y_{h,j}\|_{H_0^1(\Omega)}^2 \leq 2C_1 \|u - a(\cdot, \cdot, 0)\|_{L^2(\Omega \times I_j)}^2 + 2|C_a| \tau_j \|y_{h,j}\|_{L^2(\Omega)}^2 + \|y_{h,j-1}\|_{L^2(\Omega)}^2. \quad (4.3)$$

With the discrete Gronwall's inequality and the fact that $\|y_{h,0}\|_{L^2(\Omega)} \leq \|y_0\|_{L^2(\Omega)}$ and $\tau_j \leq \tau$ for every $j = 1, \dots, N_\tau$ we get

$$\|y_{h,j}\|_{L^2(\Omega)}^2 \leq (1 - 2|C_a|\tau)^{-j} \left(\|y_0\|_{L^2(\Omega)}^2 + 2C_1 \sum_{k=0}^{j-1} (1 - 2|C_a|\tau)^k \|u - a(\cdot, \cdot, 0)\|_{L^2(\Omega \times I_{k+1})}^2 \right); \quad (4.4)$$

see, for instance, Emmrich (1999). From our assumptions $4|C_a|\tau < 1$ and $\tau \leq \rho \tau_k$ for every k , and using that

$$\frac{1}{1 - 2|C_a|\tau} = 1 + \frac{2|C_a|\tau}{1 - 2|C_a|\tau} \leq \exp\left(\frac{2|C_a|\tau}{1 - 2|C_a|\tau}\right)$$

we obtain

$$(1 - 2|C_a|\tau)^{-j} \leq \exp\left(\frac{2|C_a|\tau j}{1 - 2|C_a|\tau}\right) \leq \exp(4\rho|C_a|T).$$

Then, (4.4) yields

$$\begin{aligned} \|y_{h,j}\|_{L^2(\Omega)}^2 &\leq \exp(4\rho|C_a|T) \left(\|y_0\|_{L^2(\Omega)}^2 + 2C_1 \sum_{k=0}^{j-1} \|u - a(\cdot, \cdot, 0)\|_{L^2(\Omega \times I_{k+1})}^2 \right) \\ &\leq \exp(4\rho|C_a|T) \left(\|y_0\|_{L^2(\Omega)}^2 + 2C_1 \|u - a(\cdot, \cdot, 0)\|_{L^2(Q)}^2 \right) \end{aligned}$$

and consequently

$$\begin{aligned} \|y_\sigma\|_{L^\infty(0,T;L^2(\Omega))} &= \max_{1 \leq j \leq N_\tau} \|y_{h,j}\|_{L^2(\Omega)} \\ &\leq \exp(2\rho|C_a|T) \max\{1, \sqrt{2C_1}\} \left(\|y_0\|_{L^2(\Omega)} + \|u - a(\cdot, \cdot, 0)\|_{L^2(Q)} \right). \end{aligned} \quad (4.5)$$

Adding the inequalities (4.3) for $j = 1, \dots, N_\tau$ we deduce

$$\begin{aligned} \Lambda_A \|y_\sigma\|_{L^2(0,T;H_0^1(\Omega))}^2 &= \Lambda_A \sum_{j=1}^{N_\tau} \tau_j \|y_{h,j}\|_{H_0^1(\Omega)}^2 \\ &\leq 2C_1 \|u - a(\cdot, \cdot, 0)\|_{L^2(Q)}^2 + \left(2|C_a|T \|y_\sigma\|_{L^\infty(0,T;L^2(\Omega))}^2 + \|y_0\|_{L^2(\Omega)}^2 \right). \end{aligned}$$

Finally, (4.2) follows from this inequality and (4.5). \square

Proof of Theorem 4.1. We divide the proof into three steps.

Step I. $\{\bar{u}_\sigma\}_\sigma$ is bounded in $L^\infty(Q)$. Let us assume that τ satisfies the condition of Lemma 4.1 and $\tau \leq \tau_0$, given by Theorem 3.1. Since the null control $u_0 \equiv 0$ is admissible for every problem (P_σ) , we deduce from the optimality of \bar{u}_σ :

$$\frac{\kappa}{2} \|\bar{u}_\sigma\|_\sigma^2 \leq J_\sigma(u_0) = \frac{1}{2} \|y_\sigma^0 - y_d\|_{L^2(\Omega)}^2,$$

where y_σ^0 denotes the discrete state associated with u_0 . From Lemma 4.1 we infer that $\{y_\sigma^0\}_\sigma$ is bounded in $L^2(Q)$. Hence, with (3.4) we deduce the existence of a constant C_1 independent of \bar{u}_σ such that

$$\|\bar{u}_\sigma\|_{L^2(Q)} \leq \|\bar{u}_\sigma\|_\sigma \leq \frac{1}{\sqrt{\kappa}} \|y_\sigma^0 - y_d\|_{L^2(Q)} \leq C_1.$$

We denote by \bar{y}_σ and $\bar{\varphi}_\sigma$ the state and adjoint state associated with \bar{u}_σ . Using again Lemma 4.1 we obtain

$$\|\bar{y}_\sigma\|_{L^\infty(0,T;L^2(\Omega))} \leq C(\|\bar{u}_\sigma - a(\cdot, \cdot, 0)\|_{L^2(Q)} + \|y_0\|_{L^2(\Omega)}) \leq C(C_1 + \|a(\cdot, \cdot, 0)\|_{L^2(Q)} + \|y_0\|_{L^2(\Omega)}) = C_2.$$

Arguing as in the proof of Lemma 4.1 we deduce the stability estimate for the solution of (3.6) corresponding to \bar{u}_σ

$$\|\bar{\varphi}_\sigma\|_{L^\infty(0,T;L^2(\Omega))} \leq C_3 \|\bar{y}_\sigma - y_d\|_{L^2(Q)} \leq C_3(C_2 + \|y_d\|_{L^2(Q)}) = C_4.$$

Next we prove the estimate

$$\|\bar{u}_\sigma\|_{L^\infty(0,T;L^2(\Omega))} \leq \frac{C_4}{\kappa}. \quad (4.6)$$

We distinguish two cases according to the definition of \mathbb{U}_σ .

Case $\mathbb{U}_\sigma = \mathbb{U}_{\sigma,0}$. From (3.26) we get for every $j = 1, \dots, N_\tau$

$$\begin{aligned} \|\bar{u}_{h,j}\|_{L^2(\Omega)} &= \left(\sum_{K \in \mathbb{K}_h} |K| \bar{u}_{K,j}^2 \right)^{\frac{1}{2}} \leq \frac{1}{\kappa} \left(\sum_{K \in \mathbb{K}_h} \frac{1}{|K|} \left(\int_K \bar{\varphi}_{h,j} \, dx \right)^2 \right)^{\frac{1}{2}} \\ &\leq \frac{1}{\kappa} \left(\sum_{K \in \mathbb{K}_h} \|\bar{\varphi}_{h,j}\|_{L^2(K)}^2 \right)^{\frac{1}{2}} = \frac{1}{\kappa} \|\bar{\varphi}_{h,j}\|_{L^2(\Omega)}. \end{aligned}$$

This inequality implies (4.6)

Case $\mathbb{U}_\sigma = \mathbb{U}_{\sigma,1}$. This time we use (3.27) to deduce

$$\begin{aligned} \|\bar{u}_{h,j}\|_{L^2(\Omega)} &= \left(\int_\Omega \left(\sum_{i=1}^{N_h} \bar{u}_{i,j} e_i \right)^2 dx \right)^{\frac{1}{2}} \leq \left(\int_\Omega \sum_{i=1}^{N_h} \bar{u}_{i,j}^2 e_i dx \right)^{\frac{1}{2}} \\ &\leq \frac{1}{\kappa} \left(\sum_{i=1}^{N_h} \frac{1}{\int_\Omega e_i dx} \left(\int_\Omega \bar{\varphi}_{h,i} e_i dx \right)^2 \right)^{\frac{1}{2}} \leq \frac{1}{\kappa} \left(\sum_{i=1}^{N_h} \int_\Omega \bar{\varphi}_{h,i}^2 e_i dx \right)^{\frac{1}{2}} = \frac{1}{\kappa} \|\bar{\varphi}_{h,j}\|_{L^2(\Omega)}. \end{aligned}$$

Hence, (4.6) is satisfied as well in this case. Then combining (2.6) and (3.3) along with the estimate (4.6) we obtain

$$\|\bar{y}_\sigma\|_{L^\infty(Q)} \leq \|y_{\bar{u}_\sigma}\|_{L^\infty(Q)} + \|y_{\bar{u}_\sigma} - \bar{y}_\sigma\|_{L^\infty(Q)} \leq C_5$$

for every $\sigma = (h, \tau)$ with $h < h_0$ and $\tau < \tau_0$. From this estimate, (4.6) and (3.8) we deduce the existence of C_6 independent of σ such that $\|\bar{\varphi}_\sigma\|_{L^\infty(Q)} \leq C_6$ for the same range of σ as before. Now using again (3.26) and (3.27) we conclude that

$$\|\bar{u}_\sigma\|_{L^\infty(Q)} \leq \frac{C_6}{\kappa} \quad (4.7)$$

for $h < h_0$ and $\tau < \tau_0$.

Take a sequence such that $\bar{u}_\sigma \xrightarrow{*} \bar{u}$ in $L^\infty(Q)$ as $\sigma \rightarrow 0$.

Step II: $\bar{u} \in U_{ad}$. It is immediate to check that $\|u_h\|_{L^1(\Omega)} = j_h(u_h)$ if $u_h \in U_h = U_{h,0}$ and $\|u_h\|_{L^1(\Omega)} \leq j_h(u_h)$ if $u_h \in U_h = U_{h,1}$. Therefore, we have

$$\|\bar{u}_\sigma\|_{L^\infty(0,T;L^1(\Omega))} = \max_{1 \leq j \leq N_h} \|\bar{u}_{h,j}\|_{L^1(\Omega)} \leq \max_{1 \leq j \leq N_h} j_h(\bar{u}_{h,j}) \leq \gamma,$$

and thus $\{\bar{u}_\sigma\}_\sigma \subset U_{ad}$. Since U_{ad} is convex and closed in $L^2(Q)$, then it is weakly closed as well. Then, the weak convergence $\bar{u}_\sigma \rightharpoonup \bar{u}$ in $L^2(Q)$ implies that $\bar{u} \in U_{ad}$.

Step III: \bar{u} is a solution of (P). Let \tilde{u} be a solution of (P). For every σ we define

$$u_\sigma = \begin{cases} P_\tau P_h \tilde{u} = \sum_{j=1}^{N_\tau} \sum_{K \in \mathbb{K}_h} \frac{1}{\tau_j |K|} \int_{I_j} \int_K \tilde{u}(x,t) dx dt \chi_j \chi_K & \text{if } \mathbb{U}_\sigma = \mathbb{U}_{\sigma,0}, \\ P_\tau E_h \tilde{u} = \sum_{j=1}^{N_\tau} \sum_{i=1}^{N_h} \frac{1}{\tau_j \int_\Omega e_i dx} \int_{I_j} \int_\Omega \tilde{u}(x,t) e_i(x) dx dt \chi_j e_i & \text{if } \mathbb{U}_\sigma = \mathbb{U}_{\sigma,1}, \end{cases} \quad (4.8)$$

where P_τ is the $L^2(0,T)$ projection operator defined in the proof of Theorem 3.1, $P_h : L^2(\Omega) \rightarrow U_{h,0}$ is the $L^2(\Omega)$ projection operator, and $E_h : L^1(\Omega) \rightarrow U_{h,1}$ is the Carstensen quasi-interpolation operator; see Carstensen (1999). First we prove that $u_\sigma \in \mathbb{U}_{\sigma,ad}$. In case $U_h = U_{h,0}$ we have $u_\sigma = \sum_{j=1}^{N_\tau} u_{h,j} \chi_j$ and for every $j = 1, \dots, N_\tau$

$$j_h(u_{h,j}) = \sum_{K \in \mathbb{K}_h} |K| \left| \frac{1}{\tau_j |K|} \int_{I_j} \int_K \tilde{u}(x,t) dx \right| \leq \sum_{K \in \mathbb{K}_h} \frac{1}{\tau_j} \int_{I_j} \int_K |\tilde{u}(x,t)| dx dt = \frac{1}{\tau_j} \int_{I_j} \|\tilde{u}(t)\|_{L^1(\Omega)} dt \leq \gamma.$$

This implies that $u_\sigma \in \mathbb{U}_{\sigma,ad}$. In the case $U_h = U_{h,1}$, we have

$$\begin{aligned} j_h(u_{h,j}) &= \sum_{i=1}^{N_h} \left| \frac{1}{\tau_j \int_\Omega e_i dx} \int_{I_j} \int_\Omega \tilde{u}(x,t) e_i(x) dx dt \right| \int_\Omega e_i dx \leq \frac{1}{\tau_j} \int_{I_j} \left(\int_\Omega |\tilde{u}(x,t)| \sum_{i=1}^{N_h} e_i dx \right) dt \\ &= \frac{1}{\tau_j} \int_{I_j} \|\tilde{u}(t)\|_{L^1(\Omega)} dt \leq \gamma. \end{aligned}$$

Using that

$$\sum_{j=1}^{N_\tau} \sum_{K \in \mathbb{K}_h} \chi_j \chi_K = \sum_{j=1}^{N_\tau} \sum_{i=1}^{N_h} e_i \chi_j = 1 \quad \text{in } Q.$$

we deduce that $\|u_\sigma\|_{L^\infty(Q)} \leq \|\tilde{u}\|_{L^\infty(Q)}$ for every σ .

In the case $\mathbb{U}_\sigma = \mathbb{U}_{\sigma,0}$, we have that $P_\tau P_h : L^2(Q) \rightarrow \mathbb{U}_{\sigma,0}$ is the $L^2(Q)$ projection operator, hence $u_\sigma \rightarrow \bar{u}$ in $L^2(Q)$ when $\sigma \rightarrow 0$. If $\mathbb{U}_\sigma = \mathbb{U}_{\sigma,1}$, then we have

$$\|\tilde{u} - u_\sigma\|_{L^2(Q)} \leq \|\tilde{u} - P_\tau \tilde{u}\|_{L^2(Q)} + \|P_\tau(\tilde{u} - E_h \tilde{u})\|_{L^2(Q)} \leq \|\tilde{u} - P_\tau \tilde{u}\|_{L^2(Q)} + \|\tilde{u} - E_h \tilde{u}\|_{L^2(Q)} \rightarrow 0 \text{ as } \sigma \rightarrow 0.$$

Indeed Corollary 2.3 implies that $\tilde{u} \in H^1(Q)$. Hence, from the convergence properties of the Carstensen operator E_h we infer the converge of the last term in the above expression. The boundedness of $\{u_\sigma\}_\sigma$ in $L^\infty(Q)$ and its strong convergence to \bar{u} in $L^2(Q)$ imply the strong convergence in every $L^p(0,T;L^q(\Omega))$ space with $1 \leq p, q < \infty$.

Next we prove that $\limsup_{\sigma \rightarrow 0} J_\sigma(u_\sigma) \leq J(\bar{u})$. Let us denote by \bar{y} and y_{u_σ} the continuous states corresponding to \bar{u} and u_σ , respectively. We also denote by y_σ the discrete state associated with u_σ . Then, using the established convergence $u_\sigma \rightarrow \bar{u}$, (2.6), and (3.3) we can easily prove

$$\|\bar{y} - y_\sigma\|_{L^\infty(Q)} \leq \|\bar{y} - y_{u_\sigma}\|_{L^\infty(Q)} + \|y_{u_\sigma} - y_\sigma\|_{L^\infty(Q)} \rightarrow 0.$$

The proved convergences of $\{y_\sigma\}_\sigma$ and $\{u_\sigma\}_\sigma$ imply that $J_\sigma(u_\sigma) \rightarrow J(\bar{u})$ as $\sigma \rightarrow 0$ if $\mathbb{U}_\sigma = \mathbb{U}_{\sigma,0}$; see (3.4). For the case $\mathbb{U}_\sigma = \mathbb{U}_{\sigma,1}$ we have

$$\begin{aligned} \|u_\sigma\|_\sigma^2 &= \sum_{j=1}^{N_\tau} \sum_{i=1}^{N_h} \tau_j \left(\int_\Omega e_i dx \right) u_{i,j}^2 = \sum_{j=1}^{N_\tau} \sum_{i=1}^{N_h} \frac{1}{\tau_j \int_\Omega e_i dx} \left(\int_{I_j} \int_\Omega \bar{u} e_i dx dt \right)^2 \\ &\leq \sum_{j=1}^{N_\tau} \sum_{i=1}^{N_h} \int_{I_j} \int_\Omega |\bar{u}|^2 e_i dx dt = \|\bar{u}\|_{L^2(Q)}^2, \end{aligned}$$

which leads to the desired inequality $\limsup_{\sigma \rightarrow 0} J_\sigma(u_\sigma) \leq J(\bar{u})$.

Using the same arguments as above, we deduce that $\bar{y}_\sigma \rightarrow \bar{y}$ strongly in $L^\infty(Q)$, where \bar{y} denotes the continuous state associated with \bar{u} . Finally, from the optimality of \bar{u}_σ and the established convergence properties we obtain with (3.4)

$$\begin{aligned} J(\bar{u}) &\leq \liminf_{\sigma \rightarrow 0} \left\{ \frac{1}{2} \|\bar{y}_\sigma - y_d\|_{L^2(Q)}^2 + \frac{\kappa}{2} \|\bar{u}_\sigma\|_{L^2(Q)}^2 \right\} \leq \liminf_{\sigma \rightarrow 0} J_\sigma(\bar{u}_\sigma) \leq \limsup_{\sigma \rightarrow 0} J_\sigma(\bar{u}_\sigma) \\ &\leq \limsup_{\sigma \rightarrow 0} J_\sigma(u_\sigma) \leq J(\bar{u}) = \inf(\mathbf{P}). \end{aligned}$$

These inequalities imply that \bar{u} is a solution of (P). Moreover, since the identity $J(\bar{u}) = J(\bar{u})$ holds, we conclude that $J_\sigma(\bar{u}_\sigma) \rightarrow J(\bar{u})$. Further we have

$$\begin{aligned} J(\bar{u}) &\leq \liminf_{\sigma \rightarrow 0} \left\{ \frac{1}{2} \|\bar{y}_\sigma - y_d\|_{L^2(Q)}^2 + \frac{\kappa}{2} \|\bar{u}_\sigma\|_{L^2(Q)}^2 \right\} \leq \limsup_{\sigma \rightarrow 0} \left\{ \frac{1}{2} \|\bar{y}_\sigma - y_d\|_{L^2(Q)}^2 + \frac{\kappa}{2} \|\bar{u}_\sigma\|_{L^2(Q)}^2 \right\} \\ &\leq \limsup_{\sigma \rightarrow 0} J_\sigma(\bar{u}_\sigma) \leq \limsup_{\sigma \rightarrow 0} J_\sigma(u_\sigma) \leq J(\bar{u}) = J(\bar{u}). \end{aligned}$$

This property and the strong convergence $\bar{y}_\sigma \rightarrow \bar{y}$ in $L^2(Q)$ yield that $\|\bar{u}_\sigma\|_{L^2(Q)} \rightarrow \|\bar{u}\|_{L^2(Q)}$. Together with the weak* convergence $\bar{u}_\sigma \rightharpoonup^* \bar{u}$ in $L^\infty(Q)$ this implies the strong convergence $\bar{u}_\sigma \rightarrow \bar{u}$ in $L^r(0, T; L^2(\Omega))$ for every $r < \infty$. Thus, (4.1) is proved. \square

The following theorem can be considered as a converse of Theorem 4.1.

THEOREM 4.2 Let \bar{u} be a strict local minimum of (P) in the $L^r(0, T; L^2(\Omega))$ sense with $r \in (\frac{4n}{4-n}, \infty)$. Then, there exist strictly positive numbers τ_0 , h_0 , ε_0 , and a sequence $\{\bar{u}_\sigma\}_\sigma \subset B_{\varepsilon_0}(\bar{u})$ of local minima of (P $_\sigma$) such that (4.1) holds and

$$J_\sigma(\bar{u}_\sigma) = \min_{u_\sigma \in \mathbb{U}_{\sigma,ad} \cap B_{\varepsilon_0}(\bar{u})} J_\sigma(u_\sigma) \quad \text{for } \tau < \tau_0 \quad \text{and} \quad h < h_0, \quad (4.9)$$

where $B_{\varepsilon_0}(\bar{u})$ is the closed ball of $L^r(0, T; L^2(\Omega))$ centered at \bar{u} and radius ε_0 .

Proof. Since \bar{u} is a strict local minimum of (P) in the $L^r(0, T; L^2(\Omega))$ sense, there exists $\varepsilon_0 > 0$ such that \bar{u} is the only solution of the problem

$$(Q) \quad \inf_{u \in \mathbb{U}_{ad} \cap B_{\varepsilon_0}(\bar{u})} J(u).$$

Now, we consider the problems

$$(Q_\sigma) \quad \inf_{u_\sigma \in \mathbb{U}_{\sigma,ad} \cap B_{\varepsilon_0}(\bar{u})} J_\sigma(u_\sigma).$$

If we define u_σ by (4.8) with $\tilde{u} = \bar{u}$, then $u_\sigma \in \mathbb{U}_{\sigma,ad}$ and $u_\sigma \rightarrow \bar{u}$ in $L^r(0, T; L^2(\Omega))$. Therefore, there exist $\tau_1 > 0$ and $h_1 > 0$ such that $u_\sigma \in B_{\varepsilon_0}(\bar{u})$ for every σ with $\tau < \tau_1$ and $h < h_1$. Hence, $\mathbb{U}_{\sigma,ad} \cap B_{\varepsilon_0}(\bar{u})$ is a compact non-empty set for every $\tau < \tau_1$ and $h < h_1$. Then, the continuity of J_σ implies the existence of at least one solution \bar{u}_σ of (Q_σ) for every σ with τ and h satisfying the previous conditions. Since $\{\bar{u}_\sigma\}_\sigma$ is bounded in $L^r(0, T; L^2(\Omega))$, taking a subsequence if necessary, we can assume that $\bar{u}_\sigma \rightharpoonup \hat{u}$ in $L^r(0, T; L^2(\Omega))$ for some \hat{u} . Due to the embedding $\mathbb{U}_{\sigma,ad} \subset U_{ad}$ we deduce that $\hat{u} \in U_{ad} \cap B_{\varepsilon_0}(\bar{u})$. Moreover, we have

$$J(\hat{u}) \leq \liminf_{\sigma \rightarrow 0} J_\sigma(\bar{u}_\sigma) \leq \limsup_{\sigma \rightarrow 0} J_\sigma(\bar{u}_\sigma) \leq \limsup_{\sigma \rightarrow 0} J_\sigma(u_\sigma) \leq J(\bar{u}).$$

Since \bar{u} is the unique solution of (Q), this inequality is only possible if $\hat{u} = \bar{u}$. Consequently, the whole family $\{\bar{u}_\sigma\}_\sigma$ converges weakly to \bar{u} in $L^r(0, T; L^2(\Omega))$ as $\sigma \rightarrow 0$ and $J_\sigma(\bar{u}_\sigma) \rightarrow J(\bar{u})$. Arguing as in the proof of Theorem 4.1, we deduce the strong convergence $\bar{u}_\sigma \rightarrow \bar{u}$ in $L^r(0, T; L^2(\Omega))$. This leads to the existence of $\tau_0 \leq \tau_1$ and $h_0 \leq h_1$ such that \bar{u}_σ belongs to the interior of the ball $B_{\varepsilon_0}(\bar{u})$ for every $\sigma = (\tau, h)$ with $\tau < \tau_0$ and $h < h_0$. Hence, every of these \bar{u}_σ is a local minimum of (P_σ) satisfying (4.9). \square

The rest of this section is dedicated to the proof of the following theorem.

THEOREM 4.3 Let us assume that \bar{u} is a local solution of (P) in the $L^r(0, T; L^2(\Omega))$ sense with $r \in (\frac{4}{4-n}, \infty)$. We also assume that $J''(\bar{u})v^2 > 0 \forall v \in C_{\bar{u}} \setminus \{0\}$. Let $\{\bar{u}_\sigma\}_\sigma$ be a family of local solutions of problems (P_σ) such that $\bar{u}_\sigma \rightarrow \bar{u}$ in $L^r(0, T; L^2(\Omega))$; see Theorem 4.2. Then, there exist positive numbers δ_0 , τ_0 , and C such that the following inequality holds:

$$\|\bar{u}_\sigma - \bar{u}\|_{L^2(Q)} \leq C(h + \tau) \quad \text{for every } \sigma = (h, \tau) \text{ with } h < h_0 \text{ and } \tau < \tau_0. \quad (4.10)$$

We prove this theorem arguing by contradiction. If (4.10) does not hold, then there exists a sequence $\{\bar{u}_{\sigma_k}\}_{k=1}^\infty$ such that $\sigma_k = (h_k, \tau_k) \rightarrow 0$ as $k \rightarrow \infty$, $h_k > 0$ and $\tau_k > 0$, and

$$\|\bar{u}_{\sigma_k} - \bar{u}\|_{L^2(Q)} > k(h_k + \tau_k) \quad \forall k \geq 1. \quad (4.11)$$

We will get a contradiction for this sequence. First we prove the next lemma.

LEMMA 4.2 Let λ be as in (2.22). There exist k_0 such that

$$(J'(\bar{u}_{\sigma_k}) - J'(\bar{u}))(\bar{u}_{\sigma_k} - \bar{u}) \geq \frac{1}{2} \min\{\lambda, \kappa\} \|\bar{u}_{\sigma_k} - \bar{u}\|_{L^2(Q)}^2 \quad \forall k \geq k_0. \quad (4.12)$$

Proof. Applying the mean value theorem, we get for some $\hat{u}_k = \bar{u} + \theta_k(\bar{u}_{\sigma_k} - \bar{u})$

$$(J'(\bar{u}_{\sigma_k}) - J'(\bar{u}))(\bar{u}_{\sigma_k} - \bar{u}) = J''(\hat{u}_k)(\bar{u}_{\sigma_k} - \bar{u})^2. \quad (4.13)$$

Set $v_k = \frac{\bar{u}_{\sigma_k} - \bar{u}}{\|\bar{u}_{\sigma_k} - \bar{u}\|_{L^2(Q)}}$. Taking a subsequence, if necessary, we can suppose that $v_k \rightharpoonup v$ in $L^2(Q)$. Below we prove that $v \in C_{\bar{u}}$. Assuming that this is true, then we argue as follows. From (2.10), the fact that

$\|v_k\|_{L^2(Q)} = 1$, and (2.22) we infer

$$\begin{aligned} \lim_{k \rightarrow \infty} J''(\hat{u}_k)v_k^2 &= \lim_{k \rightarrow \infty} \left\{ \int_Q \left(1 - \frac{\partial^2 a}{\partial y^2}(x, t, y_{\hat{u}_k}) \varphi_{\hat{u}_k} \right) z_{\hat{u}_k, v_k}^2 dx dt + \kappa \right\} \\ &= \int_Q \left(1 - \frac{\partial^2 a}{\partial y^2}(x, t, \bar{y}) \bar{\varphi} \right) z_v^2 dx dt + \kappa = J''(\bar{u})v^2 + \kappa(1 - \|v\|_{L^2(Q)}^2) \geq \kappa + (\lambda - \kappa)\|v\|_{L^2(Q)}^2. \end{aligned}$$

Above, we denoted $z_{\hat{u}_k, v_k} = G'(\hat{u}_k)v_k$ and $z_v = G'(\bar{u})v$, where $G : L^1(0, T; L^2(\Omega)) \rightarrow H^{2,1}(Q) \cap L^\infty(Q)$ is the mapping associating to each control the associated state. Since $\|v\|_{L^2(Q)} \leq 1$, the above inequality proves that

$$\lim_{k \rightarrow \infty} J''(\hat{u}_k)v_k^2 \geq \min\{\lambda, \kappa\}.$$

Therefore, there exist $k_0 > 0$ such that

$$J''(\hat{u}_k)v_k^2 \geq \frac{1}{2} \min\{\lambda, \kappa\} \quad \forall k \geq k_0,$$

or equivalently

$$J''(\hat{u}_k)(\bar{u}_{\sigma_k} - \bar{u})^2 \geq \frac{1}{2} \min\{\lambda, \kappa\} \|\bar{u}_{\sigma_k} - \bar{u}\|_{L^2(Q)}^2 \quad \forall k \geq k_0.$$

This inequality along with (4.13) leads to (4.12).

Now, we verify that $v \in C_{\bar{u}}$. From the optimality of \bar{u} and the fact that $\bar{u}_{\sigma_k} \in \mathbb{U}_{\sigma_k, ad} \subset U_{ad}$ we obtain $J'(\bar{u})v_k \geq 0$. Then, passing to the limit in this inequality when $k \rightarrow \infty$, it follows that $J'(\bar{u})v \geq 0$. Let us prove the converse inequality. We consider again the approximations $u_{\sigma_k} \in \mathbb{U}_{\sigma_k, ad}$ defined as in (4.8) with $\bar{u} = \bar{u}$. Then, we have

$$\|u_{\sigma_k} - \bar{u}\|_{L^2(Q)} \leq C_1(h_k + \tau_k)\|\bar{u}\|_{H^1(Q)} \quad \forall k \geq 1. \quad (4.14)$$

Indeed, if $\mathbb{U}_{\sigma_k} = \mathbb{U}_{\sigma_k, 0}$, the above estimate follows from the fact that u_{σ_k} is the $L^2(Q)$ projection of \bar{u} . If $\mathbb{U}_{\sigma_k} = \mathbb{U}_{\sigma_k, 1}$, the estimate was proved in Casas *et al.* (2018, Lemma 6.6). From the local optimality of \bar{u}_{σ_k} we have that $J'_{\sigma_k}(\bar{u}_{\sigma_k})(u_{\sigma_k} - \bar{u}_{\sigma_k}) \geq 0$. Using this fact we get

$$\begin{aligned} J'(\bar{u})v_k &= \frac{1}{\|\bar{u}_{\sigma_k} - \bar{u}\|_{L^2(Q)}} \{ J'(\bar{u})(\bar{u}_{\sigma_k} - u_{\sigma_k}) + J'(\bar{u})(u_{\sigma_k} - \bar{u}) \} \\ &\leq \frac{1}{\|\bar{u}_{\sigma_k} - \bar{u}\|_{L^2(Q)}} \{ [J'(\bar{u}) - J'(\bar{u}_{\sigma_k})](\bar{u}_{\sigma_k} - u_{\sigma_k}) + [J'(\bar{u}_{\sigma_k}) - J'_{\sigma_k}(\bar{u}_{\sigma_k})](\bar{u}_{\sigma_k} - u_{\sigma_k}) + J'(\bar{u})(u_{\sigma_k} - \bar{u}) \} \\ &= I_{k,1} + I_{k,2} + I_{k,3}. \end{aligned}$$

Now, we estimate every I_i term. For I_1 we use the mean value theorem, the convergence $\bar{u}_{\sigma_k} \rightarrow \bar{u}$ in $L^1(0, T; L^2(\Omega))$, and (4.14) as follows

$$\begin{aligned} |I_{k,1}| &= \frac{|J''(\bar{u} + \rho_k(\bar{u}_{\sigma_k} - \bar{u}))(\bar{u}_{\sigma_k} - u_{\sigma_k}, \bar{u} - \bar{u}_{\sigma_k})|}{\|\bar{u}_{\sigma_k} - \bar{u}\|_{L^2(Q)}} \leq C_2 \|\bar{u}_{\sigma_k} - u_{\sigma_k}\|_{L^2(Q)} \\ &\leq C_2 \left\{ \|\bar{u}_{\sigma_k} - \bar{u}\|_{L^2(Q)} + \|\bar{u} - u_{\sigma_k}\|_{L^2(Q)} \right\} \rightarrow 0 \text{ as } k \rightarrow \infty. \end{aligned}$$

To estimate $I_{k,2}$ we use (2.9) and (3.5) to get

$$\begin{aligned} I_{k,2} &= \frac{1}{\|\bar{u}_{\sigma_k} - \bar{u}\|_{L^2(Q)}} \int_Q (\varphi_{\bar{u}_{\sigma_k}} - \bar{\varphi}_{\sigma_k})(\bar{u}_{\sigma_k} - u_{\sigma_k}) \, dx \, dt \\ &\quad + \frac{\kappa}{\|\bar{u}_{\sigma_k} - \bar{u}\|_{L^2(Q)}} \left[\int_Q \bar{u}_{\sigma_k} (\bar{u}_{\sigma_k} - u_{\sigma_k}) \, dx \, dt - (\bar{u}_{\sigma_k}, \bar{u}_{\sigma_k} - u_{\sigma_k})_{\sigma_k} \right]. \end{aligned} \quad (4.15)$$

To estimate the first integral in (4.15) we use (3.7) and (4.11) along with the boundedness of $\{\bar{u}_{\sigma_k} - u_{\sigma_k}\}_{k=0}^\infty$ in $L^2(Q)$ (actually $\|\bar{u}_{\sigma_k} - u_{\sigma_k}\|_{L^2(Q)} \rightarrow 0$) as follows

$$\begin{aligned} &\left| \frac{1}{\|\bar{u}_{\sigma_k} - \bar{u}\|_{L^2(Q)}} \int_Q (\varphi_{\bar{u}_{\sigma_k}} - \bar{\varphi}_{\sigma_k})(\bar{u}_{\sigma_k} - u_{\sigma_k}) \, dx \, dt \right| \leq \frac{1}{\|\bar{u}_{\sigma_k} - \bar{u}\|_{L^2(Q)}} \|\varphi_{\bar{u}_{\sigma_k}} - \bar{\varphi}_{\sigma_k}\|_{L^2(Q)} \|\bar{u}_{\sigma_k} - u_{\sigma_k}\|_{L^2(Q)} \\ &\leq C_3 \frac{h_k^2 + \tau_k}{\|\bar{u}_{\sigma_k} - \bar{u}\|_{L^2(Q)}} \|\bar{u}_{\sigma_k} - u_{\sigma_k}\|_{L^2(Q)} \rightarrow 0 \text{ as } k \rightarrow \infty. \end{aligned} \quad (4.16)$$

In the case $\mathbb{U}_{\sigma_k} = \mathbb{U}_{\sigma_k,0}$, the scalar products $(\cdot, \cdot)_{L^2(Q)}$ and $(\cdot, \cdot)_{\sigma_k}$ coincide. Hence, the last two terms of (4.15) cancel and we get from (4.16) that $|I_{k,2}| \rightarrow 0$. If $\mathbb{U}_{\sigma_k} = \mathbb{U}_{\sigma_k,1}$, we first observe that

$$(u_\sigma, P_\tau E_h u)_\sigma = \int_Q u_\sigma u \, dx \, dt \quad \forall u_\sigma \in \mathbb{U}_\sigma \text{ and } \forall u \in L^1(Q). \quad (4.17)$$

It is immediate to check this identity. Moreover, from (3.4) we have that $\|\bar{u}_{\sigma_k}\|_{L^2(Q)} \leq \|\bar{u}_{\sigma_k}\|_{\sigma_k}$. This property, (4.17) with $u_\sigma = \bar{u}_{\sigma_k}$ and $u = \bar{u}$, (4.14), and (4.11) yield

$$\begin{aligned} &\frac{1}{\|\bar{u}_{\sigma_k} - \bar{u}\|_{L^2(Q)}} \left\{ \int_Q \bar{u}_{\sigma_k} (\bar{u}_{\sigma_k} - u_{\sigma_k}) \, dx \, dt - (\bar{u}_{\sigma_k}, \bar{u}_{\sigma_k} - u_{\sigma_k})_{\sigma_k} \right\} \leq \frac{1}{\|\bar{u}_{\sigma_k} - \bar{u}\|_{L^2(Q)}} \int_Q \bar{u}_{\sigma_k} (\bar{u} - u_{\sigma_k}) \, dx \, dt \\ &\leq \frac{\|\bar{u} - u_{\sigma_k}\|_{L^2(Q)}}{\|\bar{u}_{\sigma_k} - \bar{u}\|_{L^2(Q)}} \|\bar{u}_{\sigma_k}\|_{L^2(Q)} \leq C_1 \frac{h_k + \tau_k}{\|\bar{u}_{\sigma_k} - \bar{u}\|_{L^2(Q)}} \|\bar{u}_{\sigma_k}\|_{L^2(Q)} \rightarrow 0 \text{ as } k \rightarrow \infty. \end{aligned} \quad (4.18)$$

From (4.15), (4.16), and (4.18) we infer that $\lim_{k \rightarrow \infty} I_{k,2} \leq 0$. The estimate of the last term $I_{k,3}$ is an immediate consequence of (2.9), (4.11) and (4.14)

$$|I_{k,3}| \leq \|\bar{\varphi} + \kappa \bar{u}\|_{L^2(Q)} \frac{\|u_{\sigma_k} - \bar{u}\|_{L^2(Q)}}{\|\bar{u}_{\sigma_k} - \bar{u}\|_{L^2(Q)}} \leq C_1 \|\bar{\varphi} + \kappa \bar{u}\|_{L^2(Q)} \frac{h + \tau}{\|\bar{u}_{\sigma_k} - \bar{u}\|_{L^2(Q)}} \rightarrow 0 \text{ as } k \rightarrow \infty.$$

Thus, we have that $J'(\bar{u})v = \lim_{k \rightarrow \infty} J'(\bar{u})v_k \leq 0$, and consequently $J'(\bar{u})v = 0$, which is the first condition to have $v \in C_{\bar{u}}$.

Now, take $t \in I_\gamma$. This means that $\|\bar{u}(t)\|_{L^1(\Omega)} = \gamma$. Since $\bar{u}_{\sigma_k} \in U_{ad}$ we have that $\|\bar{u}_{\sigma_k}\|_{L^1(\Omega)} \leq \gamma$. As a consequence, we get with (2.20) an the convexity of j

$$\begin{aligned} j'(\bar{u}(t); v(t)) &\leq \liminf_{k \rightarrow \infty} j'(\bar{u}(t); v_k(t)) = \liminf_{k \rightarrow \infty} \frac{1}{\|\bar{u}_{\sigma_k} - \bar{u}\|_{L^2(Q)}} j'(\bar{u}(t); \bar{u}_{\sigma_k}(t) - \bar{u}(t)) \\ &= \liminf_{k \rightarrow \infty} \frac{1}{\|\bar{u}_{\sigma_k} - \bar{u}\|_{L^2(Q)}} \liminf_{\rho \rightarrow 0} \frac{j(\bar{u}(t) + \rho(\bar{u}_{\sigma_k}(t) - \bar{u}(t))) - j(\bar{u}(t))}{\rho} \\ &\leq \liminf_{k \rightarrow \infty} \frac{1}{\|\bar{u}_{\sigma_k} - \bar{u}\|_{L^2(Q)}} [j(\bar{u}_{\sigma_k}(t)) - j(\bar{u}(t))] = \liminf_{k \rightarrow \infty} \frac{1}{\|\bar{u}_{\sigma_k} - \bar{u}\|_{L^2(Q)}} [\|\bar{u}_{\sigma_k}(t)\|_{L^1(\Omega)} - \gamma] \leq 0. \end{aligned}$$

It remains to prove that $j'(\bar{u}(t); v(t)) = 0$ if $t \in I_\gamma^+$. The inequality $j'(\bar{u}(t); v(t)) \leq 0$ for $t \in I_\gamma^+$ implies

$$-\int_{\Omega_{\bar{u}(t)}^+} v \, dx + \int_{\Omega_{\bar{u}(t)}^-} v \, dx \geq \int_{\Omega_{\bar{u}(t)}^0} |v| \, dx. \quad (4.19)$$

Using (2.15), (2.18), and (4.19) we obtain

$$\begin{aligned} 0 &= J'(\bar{u})v = \int_Q (\bar{\varphi} + \kappa \bar{u})v \, dx \, dt = - \int_Q \bar{\mu}v \, dx \, dt = - \int_{I_\gamma^+} \int_\Omega \bar{\mu}v \, dx \, dt \\ &= - \int_{I_\gamma^+} \left[\int_{\Omega_{\bar{u}(t)}^+} \|\bar{\mu}(t)\|_{L^\infty(\Omega)} v \, dx - \int_{\Omega_{\bar{u}(t)}^-} \|\bar{\mu}(t)\|_{L^\infty(\Omega)} v \, dx + \int_{\Omega_{\bar{u}(t)}^0} \bar{\mu}v \, dx \right] dt \\ &\geq \int_{I_\gamma^+} \int_{\Omega_{\bar{u}(t)}^0} [\|\bar{\mu}(t)\|_{L^\infty(\Omega)} |v| - \bar{\mu}v] \, dx \, dt. \end{aligned}$$

This inequality is possible if and only if $\|\bar{\mu}(t)\|_{L^\infty(\Omega)} |v| = \bar{\mu}v$ in $\Omega_{\bar{u}(t)}^0 \times (0, T)$. Now, this latter identity and the fact that $\bar{\mu}(t) = 0$ if $t \notin I_\gamma^+$ we infer

$$\begin{aligned} 0 &\geq \int_{I_\gamma^+} \|\bar{\mu}(t)\|_{L^\infty(\Omega)} j'(\bar{u}(t); v(t)) \, dt = \int_{I_\gamma^+} \|\bar{\mu}(t)\|_{L^\infty(\Omega)} \left[\int_{\Omega_{\bar{u}(t)}^+} v(t) \, dx - \int_{\Omega_{\bar{u}(t)}^-} v(t) \, dx + \int_{\Omega_{\bar{u}(t)}^0} |v(t)| \, dx \right] dt \\ &= \int_{I_\gamma^+} \int_\Omega \bar{\mu}v \, dx \, dt = \int_Q \bar{\mu}v \, dx \, dt = -J'(\bar{u})v = 0, \end{aligned}$$

which implies that $j'(\bar{u}(t); v(t)) = 0$ for almost every $t \in I_\gamma^+$. This concludes the proof of $v \in C_{\bar{u}}$. \square

Proof of Theorem 4.3. Let us take k_0 big enough so that (3.7) and (4.12) hold. The goal is to prove that (4.11) is not possible. To this end, we take $u_{\sigma_k} \in \mathbb{U}_{\sigma_k, ad}$ as in the proof of Lemma 4.2. Using the optimality of \bar{u}_{σ_k} we get

$$\begin{aligned} 0 &\leq J'(\bar{u}_{\sigma_k})(u_{\sigma_k} - \bar{u}_{\sigma_k}) \\ &= J'(\bar{u}_{\sigma_k})(\bar{u} - \bar{u}_{\sigma_k}) + J'(\bar{u})(u_{\sigma_k} - \bar{u}) + [J'(\bar{u}_{\sigma_k}) - J'(\bar{u})](u_{\sigma_k} - \bar{u}) + [J'_{\sigma_k}(\bar{u}_{\sigma_k}) - J'(\bar{u}_{\sigma_k})](u_{\sigma_k} - \bar{u}_{\sigma_k}). \end{aligned}$$

We also have that $J'(\bar{u})(\bar{u}_{\sigma_k} - \bar{u}) \geq 0$. Adding these two inequalities and using (4.12) we infer

$$\begin{aligned} \frac{1}{2} \min\{\lambda, \kappa\} \|\bar{u}_{\sigma_k} - \bar{u}\|_{L^2(Q)}^2 &\leq [J'(\bar{u}_{\sigma_k}) - J'(\bar{u})](\bar{u}_{\sigma_k} - \bar{u}) \leq J'(\bar{u})(u_{\sigma_k} - \bar{u}) \\ &+ [J'(\bar{u}_{\sigma_k}) - J'(\bar{u})](u_{\sigma_k} - \bar{u}) + [J'_{\sigma_k}(\bar{u}_{\sigma_k}) - J'(\bar{u}_{\sigma_k})](u_{\sigma_k} - \bar{u}_{\sigma_k}) = I_{k,1} + I_{k,2} + I_{k,3}. \end{aligned} \quad (4.20)$$

To estimate $I_{k,1}$ we use the property

$$\|u_{\sigma_k} - \bar{u}\|_{H^1(Q)^*} \leq C_1 (h_k^2 + \tau_k^2) \|\bar{u}\|_{H^1(Q)}; \quad (4.21)$$

see Casas *et al.* (2018). With this inequality we get

$$|I_{k,1}| \leq \|\bar{\varphi} + \kappa \bar{u}\|_{H^1(Q)} \|u_{\sigma_k} - \bar{u}\|_{H^1(Q)^*} \leq C_1 (h_k^2 + \tau_k^2) \|\bar{\varphi} + \kappa \bar{u}\|_{H^1(Q)}. \quad (4.22)$$

For the estimate of $I_{k,2}$ we use the mean value theorem, the convergence $\bar{u}_{\sigma_k} \rightarrow \bar{u}$ in $L^1(0, T; L^2(\Omega))$, and (4.14) to get

$$\begin{aligned} |I_{k,2}| &= |J''(\bar{u} + \rho_k(\bar{u}_{\sigma_k} - \bar{u}))(u_{\sigma_k} - \bar{u}, \bar{u}_{\sigma_k} - \bar{u})| \leq C_2 \|u_{\sigma_k} - \bar{u}\|_{L^2(Q)} \|\bar{u}_{\sigma_k} - \bar{u}\|_{L^2(Q)} \\ &\leq C_3 (h + \tau) \|\bar{u}_{\sigma_k} - \bar{u}\|_{L^2(Q)}. \end{aligned} \quad (4.23)$$

To deal with $I_{k,3}$ we apply (2.9) and (3.5) to deduce

$$I_{k,3} = \int_Q (\bar{\varphi}_{\sigma_k} - \varphi_{\bar{u}_{\sigma_k}})(u_{\sigma_k} - \bar{u}_{\sigma_k}) \, dx \, dt + \kappa \left[(\bar{u}_{\sigma_k}, u_{\sigma_k} - \bar{u}_{\sigma_k})_{\sigma_k} - \int_Q \bar{u}_{\sigma_k} (u_{\sigma_k} - \bar{u}_{\sigma_k}) \, dx \, dt \right]. \quad (4.24)$$

With (3.7) and (4.14) we obtain

$$\begin{aligned} & \left| \int_Q (\bar{\varphi}_{\sigma_k} - \varphi_{\bar{u}_{\sigma_k}})(u_{\sigma_k} - \bar{u}_{\sigma_k}) \, dx \, dt \right| \leq \| \bar{\varphi}_{\sigma_k} - \varphi_{\bar{u}_{\sigma_k}} \|_{L^2(Q)} \| u_{\sigma_k} - \bar{u}_{\sigma_k} \|_{L^2(Q)} \\ & \leq C_4 (h_k^2 + \tau_k) \left(\| u_{\sigma_k} - \bar{u} \|_{L^2(Q)} + \| \bar{u} - \bar{u}_{\sigma_k} \|_{L^2(Q)} \right) \\ & \leq C_5 (h_k^2 + \tau_k) (h_k + \tau_k) + C_4 (h_k^2 + \tau_k) \| \bar{u} - \bar{u}_{\sigma_k} \|_{L^2(Q)}. \end{aligned} \quad (4.25)$$

The last two terms of (4.24) cancel if $\mathbb{U}_{\sigma_k} = \mathbb{U}_{\sigma_k,0}$. In the case $\mathbb{U}_{\sigma_k} = \mathbb{U}_{\sigma_k,1}$, (3.4), (4.17), (4.14), and (4.21) yield

$$\begin{aligned} & (\bar{u}_{\sigma_k}, u_{\sigma_k} - \bar{u}_{\sigma_k})_{\sigma_k} - \int_Q \bar{u}_{\sigma_k} (u_{\sigma_k} - \bar{u}_{\sigma_k}) \, dx \, dt \leq \int_Q \bar{u}_{\sigma_k} (\bar{u} - u_{\sigma_k}) \, dx \, dt \\ & = \int_Q (\bar{u}_{\sigma_k} - \bar{u})(\bar{u} - u_{\sigma_k}) \, dx \, dt + \int_Q \bar{u} (\bar{u} - u_{\sigma_k}) \, dx \, dt \\ & \leq \| \bar{u}_{\sigma_k} - \bar{u} \|_{L^2(Q)} \| \bar{u} - u_{\sigma_k} \|_{L^2(Q)} + \| \bar{u} \|_{H^1(Q)} \| \bar{u} - u_{\sigma_k} \|_{H^1(Q)^*} \\ & \leq C_6 (h_k + \tau_k) \| \bar{u}_{\sigma_k} - \bar{u} \|_{L^2(Q)} + C_7 (h_k^2 + \tau_k^2) \| \bar{u} \|_{H^1(Q)}. \end{aligned} \quad (4.26)$$

The estimates (4.24)-(4.16) lead to

$$|I_{k,3}| \leq C_8 (h_k^2 + \tau_k^2) + C_9 (h_k + \tau_k) \| \bar{u}_{\sigma_k} - \bar{u} \|_{L^2(Q)}. \quad (4.27)$$

Finally, (4.20), (4.22), (4.23), and (4.27) imply

$$\| \bar{u}_{\sigma_k} - \bar{u} \|_{L^2(Q)} \leq C_{10} (h_k + \tau_k) \quad \forall k \geq k_0,$$

which contradicts (4.11). \square

5. Numerical Examples

Let Ω be $(0,1)^n$, $n = 1$ or $n = 2$, $A = -\Delta$, $a \equiv 0$, $y_0 \equiv 0$, $T = 1$, $\kappa = 10^{-4}$, and

$$y_d(x, t) = \exp(-20[(x - 0.2)^2 + (t - 0.2)^2]) + \exp(-20[(x - 0.7)^2 + (t - 0.9)^2]) \text{ if } n = 1,$$

or

$$\begin{aligned} y_d(x, t) &= \exp(-20[(x_1 - 0.2)^2 + (x_2 - 0.2)^2 + (t - 0.2)^2]) \\ &\quad + \exp(-20[(x_1 - 0.7)^2 + (x_2 - 0.7)^2 + (t - 0.9)^2]) \text{ if } n = 2. \end{aligned}$$

Notice that all the results obtained in the paper are also valid for dimension $n = 1$. For dimension 1, these data correspond to the problem presented in (Casas *et al.*, 2017a, Remark 2.11) and also studied in Casas *et al.* (2017b, 2018). The problem in dimension 2 was introduced in Casas *et al.* (2018).

To discretize the problems, we use two families of uniform partitions in space and time, with $h_i = 2^{-i}\sqrt{2^n - 1}$ and $\tau_j = 2^{-j}$, and denote $\sigma_{i,j} = (h_i, \tau_j)$. The discrete problems are solved using a projected gradient algorithm with the Barzilai-Borwein strategy as line search; see (Barzilai & Borwein, 1988, eq. (5)). Projection strategies onto the L^1 -ball can be found in Condat (2016).

5.1 Sparsity patterns

In Figure 1 we show the solutions obtained for the one-dimensional problem as the bound parameter γ varies in $\{0.5, 1, 2, 3\}$. To discretize the problem we use the control space $\mathbb{U}_{\sigma,1}$ at the discretization level $i = j = 10$. We also plot at the left hand side of the graph the norm in $L^1(\Omega)$ of $\bar{u}_\sigma(\cdot, t)$ for all $t \in [0, 1]$, (this norm is computed with the approximation $j_h(u_{h,j})$). We use a dark green line for the norm and a magenta line for the bound. Notice that when the control constraint is attained, the solution exhibits a sparsity pattern that varies with time. We have coloured in grey the zero-level set of \bar{u}_σ to emphasize this behaviour. For $\gamma = 0.5$ and $\gamma = 1$, we have that the control constraint is active for all $t \in [0, 1]$ (green line and magenta line coincide). For $\gamma = 2$, we have that $\|\bar{u}_\sigma(\cdot, t)\|_{L^1(\Omega)} < \gamma$ if $t \in J_1 = (0.4814, 0.5723)$ and if $t > 0.9980$; for $\gamma = 3$, $\|\bar{u}_\sigma(\cdot, t)\|_{L^1(\Omega)} < \gamma$ if $t \in J_2 = (0.4502, 0.6182)$ and if $t > 0.9971$. Black lines are drawn to separate these regions. As soon as the norm constraint is not active we do not observe sparsity, in the sense that there are no subintervals in space where the control is identically zero. This behavior is consistent with the optimality condition expressed in (2.18).

The sparsity behavior obtained by means of the constraint imposed by $u \in U_{ad}$ should also be compared to sparsity phenomena implied by non-smooth cost-functionals, as considered in Casas *et al.* (2017a), for example. The functional in that paper, which is closest to the situation of the present one is given by $u \rightarrow \|u\|_{L^2(0,T;L^1(\Omega))}$, ie. it considers the L^2 -norm in time, compared to the L^∞ -norm used here. In both cases the L^1 norm in space is used. In (Casas *et al.*, 2017a, Figure 1) a numerical result with the same desired state as in Figure 1 of the present paper is presented. It lies in the nature of these two different sparsity enhancing approaches, that the solution in (Casas *et al.*, 2017a, Figure 1) also exhibits intervals of sparsity in the regions corresponding to J_1, J_2 .

In Figure 2 we show, at nine different instants of time, the solution obtained for the two-dimensional problem, using the control space $\mathbb{U}_{\sigma,0}$ at the discretization level $i = j = 7$. The control constraint parameter is set to $\gamma = 2$. The norm of the optimal control in $L^1(\Omega)$ is also reported at the indicated time instances. Again, the solution exhibits a sparsity pattern that varies with time, and there is not sparsity if the control constraint is inactive. The subdomains where $\bar{u}_\sigma(x, t)$ vanishes are coloured in grey.

5.2 Convergence rates

We show convergence rates for the problem in dimension 1. In this case we take the bound $\gamma = 4$. Since we do not have the analytic solution, we denote $I = 13$ and take as reference solution the one obtained for $\sigma_{I,I}$.

Three tests are carried out for each of the three discretizations of the control proposed in Section 3.2. In the first test, we take $h_i = \tau_i$, $i = 8, 9, 10$; in the second one, we take a fixed fine discretization in time given by τ_I , $I = 13$, and solve for h_i , $i = 8, 9, 10$; finally, we fix the discretization parameter in space to h_I , $I = 13$, and solve for τ_i , $i = 8, 9, 10$. We measure the experimental order of convergence (EOC) between two consecutive simultaneous refinement levels by setting

$$EOC = \log_2 \|\bar{u}_{\sigma_{I,I}} - \bar{u}_{\sigma_{i-1,i-1}}\|_{L^2(Q)} - \log_2 \|\bar{u}_{\sigma_{I,I}} - \bar{u}_{\sigma_{i,i}}\|_{L^2(Q)},$$

and analogously for the refinement in space and in time, respectively.

Results are shown in Table 1 for simultaneous refinement, Table 2 for refinement in space, and Table 3 for refinement in time. Except for a superconvergence phenomenon in the space discretization when we use continuous piecewise linear in space approximations of the control, all the experimental results are as predicted in Theorem 4.3.




FIG. 1. 1D problem. Continuous piecewise linear approximation in space, piecewise constant approximation in time, of the optimal control for different values of γ . The norm in $L^1(\Omega)$ at every instant of time is also shown with a dark green line located on the plane $x = -0.1$ together with the magenta line $z = \gamma$. In grey, the zero-level set of \bar{u}_σ .




FIG. 2. 2D problem. Piecewise constant approximation of the optimal control. In grey, the level sets $\bar{u}_\sigma(\cdot, t_j) = 0$.

$h_i = \tau_i$	$\mathbb{U}_{\sigma,0}$		$\mathbb{U}_{\sigma,1}$	
	$\ \bar{u}_{\sigma_{l,l}} - \bar{u}_{\sigma_{i,i}}\ _{L^2(Q)}$	<i>EOC</i>	$\ \bar{u}_{\sigma_{l,l}} - \bar{u}_{\sigma_{i,i}}\ _{L^2(Q)}$	<i>EOC</i>
2^{-8}	$2.01\text{E}-1$	—	$1.76\text{E}-1$	—
2^{-9}	$1.02\text{E}-1$	0.98	$8.93\text{E}-2$	0.98
2^{-10}	$5.11\text{E}-2$	0.99	$4.49\text{E}-2$	0.99

Table 1. Experimental order of convergence. Simultaneous refinement in space and time.

h_i	$\mathbb{U}_{\sigma,0}$		$\mathbb{U}_{\sigma,1}$	
	$\ \bar{u}_{\sigma_{l,l}} - \bar{u}_{\sigma_{i,i}}\ _{L^2(Q)}$	<i>EOC</i>	$\ \bar{u}_{\sigma_{l,l}} - \bar{u}_{\sigma_{i,i}}\ _{L^2(Q)}$	<i>EOC</i>
2^{-8}	$9.86\text{E}-2$	—	$1.10\text{E}-02$	—
2^{-9}	$4.93\text{E}-2$	1.00	$3.87\text{E}-03$	1.51
2^{-10}	$2.45\text{E}-2$	1.01	$1.34\text{E}-03$	1.53

Table 2. Experimental order of convergence. Refinement in space.

τ_i	$\mathbb{U}_{\sigma,0}$		$\mathbb{U}_{\sigma,1}$	
	$\bar{u}_{\sigma_{l,l}} - \bar{u}_{\sigma_{i,i}}$	<i>EOC</i>	$\bar{u}_{\sigma_{l,l}} - \bar{u}_{\sigma_{i,i}}$	<i>EOC</i>
2^{-8}	$1.76\text{E}-1$	—	$1.76\text{E}-1$	—
2^{-9}	$8.93\text{E}-2$	0.98	$8.93\text{E}-2$	0.98
2^{-10}	$4.49\text{E}-2$	0.99	$4.49\text{E}-2$	0.99

Table 3. Experimental order of convergence. Refinement in time.

References

- AKMAN, T., YÜCEL, H. & KARASÖZEN, B. (2014) A priori error analysis of the upwind symmetric interior penalty Galerkin (SIPG) method for the optimal control problems governed by unsteady convection diffusion equations. *Comput. Optim. Appl.*, **57**, 703–729.
- BARZILAI, J. & BORWEIN, J. M. (1988) Two-point step size gradient methods. *IMA J. Numer. Anal.*, **8**, 141–148.
- BRENNER, S. C. & SCOTT, L. R. (2008) *The mathematical theory of finite element methods*. Texts in Applied Mathematics, vol. 15, third edn. New York: Springer.
- CARSTENSEN, C. (1999) Quasi-interpolation and a posteriori error analysis in finite element methods. *M2AN Math. Model. Numer. Anal.*, **33**, 1187–1202.
- CASAS, E., HERZOG, R. & WACHSMUTH, G. (2012) Approximation of sparse controls in semilinear equations by piecewise linear functions. *Numer. Math.*, **122**, 645–669.
- CASAS, E., HERZOG, R. & WACHSMUTH, G. (2017a) Analysis of spatio-temporally sparse optimal control problems of semilinear parabolic equations. *ESAIM Control Optim. Calc. Var.*, **23**, 263–295.

- CASAS, E., MATEOS, M. & RÖSCH, A. (2017b) Finite element approximation of sparse parabolic control problems. *Math. Control Relat. Fields*, **7**, 393–417.
- CASAS, E., MATEOS, M. & RÖSCH, A. (2018) Improved approximation rates for a parabolic control problem with an objective promoting directional sparsity. *Comput. Optim. Appl.*, **70**, 239–266.
- CASAS, E., MATEOS, M. & RÖSCH, A. (2019) Error estimates for semilinear parabolic control problems in the absence of Tikhonov term. *SIAM J. Control Optim.*, **57**, 2515–2540.
- CASAS, E. & KUNISCH, K. (2021) Optimal control of semilinear parabolic equations with non-smooth pointwise-integral control constraints in time-space. *Appl. Math. Optim.*, To appear.
- CHRISTOF, C. & VEXLER, B. (2021) New regularity results and finite element error estimates for a class of parabolic optimal control problems with pointwise state constraints. *ESAIM Control Optim. Calc. Var.*, **27**, Paper No. 4, 39.
- CHRYSAFINOS, K. & KARATZAS, E. N. (2012) Symmetric error estimates for discontinuous Galerkin approximations for an optimal control problem associated to semilinear parabolic PDE's. *Discrete Contin. Dyn. Syst. Ser. B*, **17**, 1473–1506.
- CONDAT, L. (2016) Fast projection onto the simplex and the l_1 ball. *Math. Program.*, **158**, 575–585.
- DI BENEDETTO, E. (1986) On the local behaviour of solutions of degenerate parabolic equations with measurable coefficients. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, **13**, 487–535.
- EMMRICH, E. (1999) Discrete versions of Gronwall's lemma and their application to the numerical analysis of parabolic problems. *Technical Report 637*. Fachbereich 3 Preprint Reihe Mathematik, Berlin TU.
- GONG, W., HINZE, M. & ZHOU, Z. (2014) A priori error analysis for finite element approximation of parabolic optimal control problems with pointwise control. *SIAM J. Control Optim.*, **52**, 97–119.
- GUNZBURGER, M. D. & MANSERVISI, S. (1999) The velocity tracking problem for Navier-Stokes flows with bounded distributed controls. *SIAM J. Control Optim.*, **37**, 1913–1945.
- HOPPE, F. & NEITZEL, I. (2020) Optimal control of quasilinear parabolic PDEs with state constraints. In Revision. Available as INS Preprint No. 2004.
- HUANG, F., ZHENG, Z. & PENG, Y. (2018) Error estimates of the space-time spectral method for parabolic control problems. *Comput. Math. Appl.*, **75**, 335–348.
- LADYZHENSKAYA, O., SOLONNIKOV, V. & URAL'TSEVA, N. (1988) *Linear and Quasilinear Equations of Parabolic Type*. Providence: American Mathematical Society.
- LEYKEKHMAN, D. & VEXLER, B. (2013) Optimal a priori error estimates of parabolic optimal control problems with pointwise control. *SIAM J. Numer. Anal.*, **51**, 2797–2821.
- MEIDNER, D. & VEXLER, B. (2011) A priori error analysis of the Petrov-Galerkin Crank-Nicolson scheme for parabolic optimal control problems. *SIAM J. Control Optim.*, **49**, 2183–2211.
- MEIDNER, D. & VEXLER, B. (2018) Optimal error estimates for fully discrete Galerkin approximations of semilinear parabolic equations. *ESAIM: M2AN*, **52**, 2307–2325.

- PIEPER, K. (2015) Finite element discretization and efficient numerical solution of elliptic and parabolic sparse control problems. *Ph.D. thesis*, Technische Universität München.
- RÖSCH, A. & WACHSMUTH, G. (2017) Mass lumping for the optimal control of elliptic partial differential equations. *SIAM J. Numer. Anal.*, **55**, 1412–1436.
- VON DANIELS, N., HINZE, M. & VIERLING, M. (2015) Crank-Nicolson time stepping and variational discretization of control-constrained parabolic optimal control problems. *SIAM J. Control Optim.*, **53**, 1182–1198.