

|                        |                |         |       |             |
|------------------------|----------------|---------|-------|-------------|
| Phyton (Horn, Austria) | <b>Vol. 43</b> | Fasc. 1 | 1–240 | 21. 7. 2003 |
|------------------------|----------------|---------|-------|-------------|

## CONTENTS

(Engl.) = Paper in English with German summary  
(Germ.) = Paper in German with English summary

|  |                       |
|--|-----------------------|
| LAUS G. & KEPLINGER K.: Alkaloids of Peruvian <i>Uncaria guianensis</i> ( <i>Rubiaceae</i> ). (Engl.)  | 1                     |
| NAYDENOV K., TREMBLAY F. & GANCHEV P.: Karyotypic diversity in European Black Pine ( <i>Pinus nigra</i> ARN.) from Bulgarian provenances. (Engl.)  | 9                     |
| TRIGAS P. & IATROU G.: <i>Asperula</i> (sect. <i>Cynanchicae</i> ) <i>brachyphylla</i> , spec. nova ( <i>Rubiaceae</i> ) from the island of Evvia (Greece). (Engl.)  | 29                    |
| TOMASELLI M., BERNARDO L. & PASSALACQUA N.: The vegetation of the Ranunculo-Nardion in the Southern Apennines (S.-Italy). (Engl.)  | 39                    |
| CASTRO-DÍEZ P., MILLA-GUTIÉRREZ R. & MONTERRAT-MARTÍ G. 2003. Comparison of methods to study phenological patterns. The case of <i>Halimium atriplicifolium</i> ( <i>Cistaceae</i> ). (Engl.)  | 59                    |
| KÖCKINGER H.: <i>Saxifraga styriaca</i> spec. nova ( <i>Saxifragaceae</i> ) – an endemic species of the eastern Niedere Tauern Mts. (Styria, Austria). (Germ.)   | 79                    |
| VRACHNAKIS Th.: Trichomes of <i>Origanum dictamnus</i> L. ( <i>Labiatae</i> ). (Engl.)   | 109                   |
| RAJPUT K. S.: Structure of cambium and its derivatives in the compressed stem of <i>Canavalia ensiformis</i> (L.) DC. ( <i>Fabaceae</i> ) (Engl.)  | 135                   |
| WONISCH A., TAUSZ M., MÜLLER M., SOJA G. & GRILL D.: Ozone-induced long-term effects on chromosomal aberration rates in root-tip meristems of spruce trees do not correspond to changes in tissue antioxidant status. (Engl.)        | 147                   |
| KIKUTA S. B.: Ultrasound acoustic emissions from bark samples differing in anatomical characteristics. (Engl.)   | 161                   |
| SUZUKI L. S., HERRIG V., FERRARESE M. L. L., RODRIGUES J. D. & FERRARESE-FILHO O.: Simultaneous effects of ferulic and vanillic acids on peroxidase and phenylalanine ammonia-lyase in soybean ( <i>Glycine max</i> ) roots. (Engl.) | 179                   |
| MORENO D. A., VILLORA G. & ROMERO L.: Age-related changes in micronutrients and related enzymex in red pepper. (Engl.)   | 187                   |
| BODHIPADMA K. & LEUNG D. W. M.: Histological changes during direct somatic embryogenesis from immature zygotic embryo explants of <i>Capsicum annum</i> L. cv. Sweet Banana. (Engl.)   | 197                   |
| SACCHETTI G., BALLERO M., SERAFINI M., MUZZOLI M., TOSI B. & POLI F.: Morphological and histochemical investigation on glandular trichomes of <i>Orobanche ramosa</i> subsp. <i>nana</i> ( <i>Orobanchaceae</i> ). (Engl.)           | 207                   |
| HAUSMANN N., WERHAHN W., HUCHZERMAYER B., BRAUN H. P. & PAPENBROCK J.: How to document the purity of mitochondria prepared from green tissue of pea, tobacco and <i>Arabidopsis thaliana</i> . (Engl.)                               | 215                   |
| MÜLLER M., STABENTHEINER E., BORZAN Ž & GUTTENBERGER H.: Further contributions to the discussion of the origin of a particular oak from the northern Dalmatia called the “green oak”. (Engl.)  | 231                   |
| Recensiones  | 37, 57, 108, 134, 160 |
| Instructions for Contributors  | 239                   |

New taxa and/or new combinations as well as typifications:

|  |    |
|--|----|
| <i>Asperula brachyphylla</i> TRIGAS & IATROU, spec. nova                                     | 30 |
| <i>Saxifraga styriaca</i> KÖCKINGER, spec. nova  | 82 |
| <i>Saxifraga androsacea</i> × <i>S. styriaca</i> = <i>S. × melzeri</i> KÖCKINGER, hybr. nova | 85 |