

Phyton (Horn, Austria)	Vol. 47	Fasc. 1–2	1–337	19. 12. 2007
------------------------	----------------	-----------	-------	--------------

CONTENTS

TEPPNER H.: Polyad development and karyology in <i>Inga</i> and <i>Calliandra</i> (<i>Mimosaceae-Ingae</i>): A reply to a recent paper in Flora.	1
DENGLER J. & BOCH S.: Taxonomy, ecology, and distribution of six remarkable plant taxa (<i>Spermatophyta</i> , <i>Bryophyta</i> , and <i>Lichenes</i>) of the Estonian dry grassland flora.	47
SLOVÁK M. & MARHOLD K.: Morphological evaluation of <i>Picris hieracioides</i> L. (<i>Compositae-Lactuceae</i>) in Slovakia.	73
UHLEMANN I., KIRSCHNER J., ØLLGAARD H. & ŠTĚPÁNEK J.: Four new species of <i>Taraxacum</i> sect. <i>Ruderalia</i> (<i>Asteraceae-Cichorieae</i>) from Central Europe and Scandinavia.	103
REBOREDO F. & SILVARES C.: Fasciation phenomena and mineral balance in <i>Spartium junceum</i> L.	123
TOMASELLI M., DOWGIALLO G., PETRAGLIA A. & CONSIGLIO L.: The summit vegetation of the Sirino-Papa massif (Southern Apennines, S. Italy). . . .	133
GARGANO D., PASSALACQUA N. G. & BERNARDO L.: Bogs and mires in Mediterranean areas: the vegetation of the marshlands of the Lacina Plain (Calabria, S. Italy).	161
BÜRCEL J.: Hybridisation in <i>Saxifraga</i> subsection <i>Kabschia</i> (<i>Saxifragaceae</i>) from the Central Himalaya.	191
GROB V., PFEIFER E. & RUTISHAUSER R.: Morphology, development and regeneration of <i>Thelethylax minutiflora</i> , a Madagascan river-weed (<i>Podostemaceae</i>).	205
OBERMAYER W. & MAYRHOFER H.: Hunting for <i>Cetrelia chicitae</i> (lichenized Ascomycetes) in the eastern European Alps (including an attempt for a morphological characterization of all taxa of the genus <i>Cetrelia</i> in Central Europe).	231
TEPPNER H. & STABENTHEINER E.: Anther opening, polyad presentation, pollenkitt and pollen adhesive in four <i>Calliandra</i> species (<i>Mimosaceae-Ingae</i>).	291
WETSCHNIG W., KNIRSCH W., ALI S. S. & PFOSSER M.: Systematic position of three little known and frequently misplaced species of <i>Hyacinthaceae</i> from Madagascar.	321
New taxa and/or new combinations as well as typifications:	
<i>Saxifraga</i> × <i>hetenbeliana</i> BÜRCEL nothospec. nova (<i>Saxifragaceae</i>)	201
<i>Taraxacum capillosum</i> H. ØLLGAARD & UHLEMANN spec. nova (<i>Asteraceae</i>) . .	105
<i>Taraxacum delectum</i> UHLEMANN spec. nova (<i>Asteraceae</i>)	110
<i>Taraxacum ottonis</i> UHLEMANN spec. nova (<i>Asteraceae</i>)	111
<i>Taraxacum subborgvallii</i> UHLEMANN, ŠTĚPÁNEK & KIRSCHNER spec. nova (<i>Asteraceae</i>)	114