

Phyton (Horn, Austria)	<b>Vol. 51</b>	Fasc. 2	177–327	20. 12. 2011
------------------------	----------------	---------	---------	--------------

## CONTENTS

KIRSCHNER J., KIRSCHNEROVÁ L. & BARTISH I.: Conservation status of two isolated populations of <i>Gentiana verna</i> ( <i>Gentianaceae</i> ) in the Czech Republic: Insights from an allozyme analysis. . . . .	177
BRULLO C., BRULLO S., GIUSSO DEL GALDO G. & SCIANDRELLO S.: Taxonomical, morphological and geobotanical considerations on <i>Seseli tortuosum</i> L. ( <i>Apiaceae</i> ) from Sicily. . . . .	201
SPETA F.: Contribution to the knowledge of <i>Othocallis monanthos</i> (C. KOCH) SPETA ( <i>Hyacinthaceae-Hyacintheae</i> ). . . . .	211
SPETA F.: Morphological and karyological studies on <i>Othocallis morrisii</i> (MEIKLE) SPETA ( <i>Hyacinthaceae</i> ) from the island of Cyprus. . . . .	217
PRAMOD S., THOMAS V. & RAO K. S.: Structural and dimensional changes in the cambium of tapping panel dryness affected bark of <i>Hevea brasiliensis</i> . . . . .	231
GIANNAKOULA A. F. & ILIAS I. F.: Responses of the photosynthetic apparatus to plant growth regulators in two sunflower cultivars ( <i>Helianthus annuus</i> L.). . . . .	245
IBRAHIM A. H., ABDEL-FATTAH G. M., EMAM F. M., ABD EL-AZIZ M. H. & SHOKR A. E.: Arbuscular mycorrhizal fungi and spermine alleviate the adverse effects of salinity stress on electrolyte leakage and productivity of wheat plants. . . . .	261
KOWALSKI T. & DROŻYŃSKA K.: Mycobiota in needles and shoots of <i>Pinus nigra</i> following infection by <i>Dothistroma septosporum</i> . . . . .	277
RAJPUT K. S. & RAO K. S.: Development of vascular cambium in the petiole of <i>Terminalia catappa</i> L. ( <i>Combretaceae</i> ). . . . .	289
SWIDRAK I., GRUBER A. & OBERHUBER W.: Cambial activity and xylem cell development in <i>Pinus cembra</i> and <i>Pinus sylvestris</i> at their climatic limits in the Eastern Alps in 2007. . . . .	299
BRAUN M. & GOTTSBERGER G.: Floral biology and breeding system of <i>Anaxagorea dolichocarpa</i> ( <i>Annonaceae</i> ), with observations on the interval between anthesis and fruit formation. . . . .	315
New taxa and/or new combinations as well as typifications:	
<i>Scilla morrisii</i> (MEIKLE) SPETA var. <i>veneris</i> (SPETA) SPETA stat. nov. ( <i>Hyacinthaceae</i> ) . . . . .	226
<i>Seseli tortuosum</i> L. subsp. <i>maritimum</i> (GUSS.) C. BRULLO & al. stat. nov. ( <i>Apiaceae</i> ) . . . . .	208
<i>Seseli tortuosum</i> L. subsp. <i>tortuosum</i> ( <i>Apiaceae</i> ) . . . . .	205
<i>Seseli tortuosum</i> L. var. <i>maritimum</i> GUSS., lectotypification, epitype ( <i>Apiaceae</i> ) . . . . .	208

New syntaxa:

—