

Long
zero-sum
free and
n-zero-sum
free
sequences

Li

1. Introduction
and
Notation

2. On long
zero-sum
free
sequences
over \mathbb{Z}_n

3. long
n-zero-sum
free
sequences
over \mathbb{Z}_n

4. Subsums
of a long
zero-sum
free
sequence.

Bibliography

Long zero-sum free and n-zero-sum free sequences over finite cyclic groups

Yuanlin Li

Brock University

January 2016

Content

Long
zero-sum
free and
 n -zero-sum
free
sequences

Li

1. Introduction
and
Notation

2. On long
zero-sum
free
sequences
over \mathbb{Z}_n

3. long
 n -zero-sum
free
sequences
over \mathbb{Z}_n

4. Subsums
of a long
zero-sum
free
sequence.

Bibliography

1. Introduction and Notation
2. On long zero-sum free sequences over \mathbb{Z}_n
3. long n -zero-sum free sequences over \mathbb{Z}_n
4. Subsums of a long zero-sum free sequence.
5. Bibliography

Introduction and Notation

Long
zero-sum
free and
 n -zero-sum
free
sequences

Li

G – Additive finite abelian group

$\mathbb{Z}_n = \mathbb{Z}/n\mathbb{Z}$ – Cyclic group of order n

$\mathcal{F}(G)$ – Free abelian monoid with basis G .

1. Introduction
and
Notation

2. On long
zero-sum
free
sequences
over \mathbb{Z}_n

3. long
 n -zero-sum
free
sequences
over \mathbb{Z}_n

4. Subsums
of a long
zero-sum
free
sequence.

Bibliography

Introduction and Notation

Long
zero-sum
free and
 n -zero-sum
free
sequences

Li

1. Introduction
and
Notation

2. On long
zero-sum
free
sequences
over \mathbb{Z}_n

3. long
 n -zero-sum
free
sequences
over \mathbb{Z}_n

4. Subsums
of a long
zero-sum
free
sequence.

Bibliography

G – Additive finite abelian group

$\mathbb{Z}_n = \mathbb{Z}/n\mathbb{Z}$ – Cyclic group of order n

$\mathcal{F}(G)$ – Free abelian monoid with basis G .

An element of $\mathcal{F}(G)$ is called a sequence of G , and A sequence of length l over G can be written in the form $S = g_1 \cdot \dots \cdot g_l$ for some $g_1, \dots, g_l \in G$.

Introduction and Notation

Long
zero-sum
free and
 n -zero-sum
free
sequences

Li

G – Additive finite abelian group

$\mathbb{Z}_n = \mathbb{Z}/n\mathbb{Z}$ – Cyclic group of order n

$\mathcal{F}(G)$ – Free abelian monoid with basis G .

An element of $\mathcal{F}(G)$ is called a sequence of G , and A sequence of length l over G can be written in the form $S = g_1 \cdot \dots \cdot g_l$ for some $g_1, \dots, g_l \in G$. S is called a *zero-sum sequence* if the sum of all elements of S is zero, and a *zero-sum free sequence* if S does not contain a nonempty zero-sum subsequence.

1. Introduction
and
Notation

2. On long
zero-sum
free
sequences
over \mathbb{Z}_n

3. long
 n -zero-sum
free
sequences
over \mathbb{Z}_n

4. Subsums
of a long
zero-sum
free
sequence.

Bibliography

Introduction and Notation

Long
zero-sum
free and
 n -zero-sum
free
sequences

Li

1. Introduction
and
Notation

2. On long
zero-sum
free
sequences
over \mathbb{Z}_n

3. long
 n -zero-sum
free
sequences
over \mathbb{Z}_n

4. Subsums
of a long
zero-sum
free
sequence.

Bibliography

G – Additive finite abelian group

$\mathbb{Z}_n = \mathbb{Z}/n\mathbb{Z}$ – Cyclic group of order n

$\mathcal{F}(G)$ – Free abelian monoid with basis G .

An element of $\mathcal{F}(G)$ is called a sequence of G , and a sequence of length l over G can be written in the form $S = g_1 \cdot \dots \cdot g_l$ for some $g_1, \dots, g_l \in G$. S is called a *zero-sum sequence* if the sum of all elements of S is zero, and a *zero-sum free* sequence if S does not contain a nonempty zero-sum subsequence. If S is a zero-sum sequence, but no proper nontrivial subsequence of S has sum zero, then S is called a *minimal zero-sum sequence*.

Introduction and Notation

Long
zero-sum
free and
 n -zero-sum
free
sequences

Li

1. Introduction
and
Notation

2. On long
zero-sum
free
sequences
over \mathbb{Z}_n

3. long
 n -zero-sum
free
sequences
over \mathbb{Z}_n

4. Subsums
of a long
zero-sum
free
sequence.

Bibliography

G – Additive finite abelian group

$\mathbb{Z}_n = \mathbb{Z}/n\mathbb{Z}$ – Cyclic group of order n

$\mathcal{F}(G)$ – Free abelian monoid with basis G .

An element of $\mathcal{F}(G)$ is called a sequence of G , and a sequence of length l over G can be written in the form $S = g_1 \cdot \dots \cdot g_l$ for some $g_1, \dots, g_l \in G$. S is called a *zero-sum sequence* if the sum of all elements of S is zero, and a *zero-sum free* sequence if S does not contain a nonempty zero-sum subsequence. If S is a zero-sum sequence, but no proper nontrivial subsequence of S has sum zero, then S is called a *minimal zero-sum sequence*.

Introduction and Notation

Definition 1.1

Let $G = \mathbb{Z}_n$, for a sequence $S = (n_1g) \cdot \dots \cdot (n_lg)$, $1 \leq n_1, \dots, n_l \leq n$, the index of S is defined by $\text{ind}(S) = \min\{\|S\|_g \mid g \in \mathbb{Z}_n, \text{ with } \mathbb{Z}_n = \langle g \rangle\}$, where $\|S\|_g = \frac{n_1 + \dots + n_l}{\text{ord}(g)}$.

Long
zero-sum
free and
 n -zero-sum
free
sequences

Li

1. Introduction
and
Notation

2. On long
zero-sum
free
sequences
over \mathbb{Z}_n

3. long
 n -zero-sum
free
sequences
over \mathbb{Z}_n

4. Subsums
of a long
zero-sum
free
sequence.

Bibliography

Introduction and Notation

Definition 1.1

Let $G = \mathbb{Z}_n$, for a sequence $S = (n_1g) \cdot \dots \cdot (n_lg)$, $1 \leq n_1, \dots, n_l \leq n$, the index of S is defined by $\text{ind}(S) = \min\{\|S\|_g \mid g \in \mathbb{Z}_n, \text{ with } \mathbb{Z}_n = \langle g \rangle\}$, where $\|S\|_g = \frac{n_1 + \dots + n_l}{\text{ord}(g)}$.

The index of a sequence is a crucial invariant in the investigation of (minimal) zero-sum sequences (resp. of zero-sum free sequences) over cyclic groups. Recently, there are considerable publications on this subject (see for example, [4, 5, 6, 7, 8, 10, 11, 12]).

Long
zero-sum
free and
 n -zero-sum
free
sequences

Li

1. Introduction
and
Notation

2. On long
zero-sum
free
sequences
over \mathbb{Z}_n

3. long
 n -zero-sum
free
sequences
over \mathbb{Z}_n

4. Subsums
of a long
zero-sum
free
sequence.

Bibliography

Introduction and Notation

Definition 1.1

Let $G = \mathbb{Z}_n$, for a sequence $S = (n_1g) \cdot \dots \cdot (n_lg)$, $1 \leq n_1, \dots, n_l \leq n$, the index of S is defined by $\text{ind}(S) = \min\{\|S\|_g \mid g \in \mathbb{Z}_n, \text{ with } \mathbb{Z}_n = \langle g \rangle\}$, where $\|S\|_g = \frac{n_1 + \dots + n_l}{\text{ord}(g)}$.

The index of a sequence is a crucial invariant in the investigation of (minimal) zero-sum sequences (resp. of zero-sum free sequences) over cyclic groups. Recently, there are considerable publications on this subject (see for example, [4, 5, 6, 7, 8, 10, 11, 12]). We say that two sequences S_1 and S_2 are *equivalent* and write $S_1 \sim S_2$ if S_2 can be obtained from S_1 through multiplication by an integer coprime to n and rearrangement of terms.

Long
zero-sum
free and
 n -zero-sum
free
sequences

Li

1. Introduction
and
Notation

2. On long
zero-sum
free
sequences
over \mathbb{Z}_n

3. long
 n -zero-sum
free
sequences
over \mathbb{Z}_n

4. Subsums
of a long
zero-sum
free
sequence.

Bibliography

Introduction and Notation

Definition 1.1

Let $G = \mathbb{Z}_n$, for a sequence $S = (n_1g) \cdot \dots \cdot (n_lg)$, $1 \leq n_1, \dots, n_l \leq n$, the index of S is defined by $\text{ind}(S) = \min\{\|S\|_g \mid g \in \mathbb{Z}_n, \text{ with } \mathbb{Z}_n = \langle g \rangle\}$, where $\|S\|_g = \frac{n_1 + \dots + n_l}{\text{ord}(g)}$.

The index of a sequence is a crucial invariant in the investigation of (minimal) zero-sum sequences (resp. of zero-sum free sequences) over cyclic groups. Recently, there are considerable publications on this subject (see for example, [4, 5, 6, 7, 8, 10, 11, 12]). We say that two sequences S_1 and S_2 are *equivalent* and write $S_1 \sim S_2$ if S_2 can be obtained from S_1 through multiplication by an integer coprime to n and rearrangement of terms. Clearly, $\text{ind}(S_1) = \text{ind}(S_2)$.

Long
zero-sum
free and
 n -zero-sum
free
sequences

Li

1. Introduction
and
Notation

2. On long
zero-sum
free
sequences
over \mathbb{Z}_n

3. long
 n -zero-sum
free
sequences
over \mathbb{Z}_n

4. Subsums
of a long
zero-sum
free
sequence.

Bibliography

Introduction and Notation

Definition 1.1

Let $G = \mathbb{Z}_n$, for a sequence $S = (n_1g) \cdot \dots \cdot (n_lg)$, $1 \leq n_1, \dots, n_l \leq n$, the index of S is defined by $\text{ind}(S) = \min\{\|S\|_g \mid g \in \mathbb{Z}_n, \text{ with } \mathbb{Z}_n = \langle g \rangle\}$, where $\|S\|_g = \frac{n_1 + \dots + n_l}{\text{ord}(g)}$.

The index of a sequence is a crucial invariant in the investigation of (minimal) zero-sum sequences (resp. of zero-sum free sequences) over cyclic groups. Recently, there are considerable publications on this subject (see for example, [4, 5, 6, 7, 8, 10, 11, 12]). We say that two sequences S_1 and S_2 are *equivalent* and write $S_1 \sim S_2$ if S_2 can be obtained from S_1 through multiplication by an integer coprime to n and rearrangement of terms. Clearly, $\text{ind}(S_1) = \text{ind}(S_2)$.

Long
zero-sum
free and
 n -zero-sum
free
sequences

Li

1. Introduction
and
Notation

2. On long
zero-sum
free
sequences
over \mathbb{Z}_n

3. long
 n -zero-sum
free
sequences
over \mathbb{Z}_n

4. Subsums
of a long
zero-sum
free
sequence.

Bibliography

Introduction and Notation

Long
zero-sum
free and
 n -zero-sum
free
sequences

Li

$\sigma(S)$ –the sum of all terms of S .

$h(S)$ –the maximum multiplicity of a term in S .

$\Sigma(S)$ –the set of all subsums of S , and $\Sigma_k(S)$ – the set of k -term subsums of S , where $k \in \mathbb{N}$.

1.
Introduction
and
Notation

2. On long
zero-sum
free
sequences
over \mathbb{Z}_n

3. long
 n -zero-sum
free
sequences
over \mathbb{Z}_n

4. Subsums
of a long
zero-sum
free
sequence.

Bibliography

Introduction and Notation

Long zero-sum free and n -zero-sum free sequences

Li

$\sigma(S)$ –the sum of all terms of S .

$h(S)$ –the maximum multiplicity of a term in S .

$\Sigma(S)$ –the set of all subsums of S , and $\Sigma_k(S)$ – the set of k -term subsums of S , where $k \in \mathbb{N}$.

Definition 1.2

Let S be a minimal zero-sum (resp. zero-sum free) sequence of elements over abelian group G . An element g_0 in S is called splittable if there exist two elements $x, y \in G$ such that $x + y = g_0$ and $Sg_0^{-1}xy$ is a minimal zero-sum (resp. zero-sum free) sequence as well; otherwise, g_0 is called unsplittable. S is called splittable if at least one of the elements of S is splittable; otherwise, it is called unsplittable.

1. Introduction and Notation

2. On long zero-sum free sequences over \mathbb{Z}_n

3. long n -zero-sum free sequences over \mathbb{Z}_n

4. Subsums of a long zero-sum free sequence.

Bibliography

Introduction and Notation

Long zero-sum free and n -zero-sum free sequences

Li

1. Introduction and Notation

2. On long zero-sum free sequences over \mathbb{Z}_n

3. long n -zero-sum free sequences over \mathbb{Z}_n

4. Subsums of a long zero-sum free sequence.

Bibliography

$\sigma(S)$ –the sum of all terms of S .

$h(S)$ –the maximum multiplicity of a term in S .

$\Sigma(S)$ –the set of all subsums of S , and $\Sigma_k(S)$ – the set of k -term subsums of S , where $k \in \mathbb{N}$.

Definition 1.2

Let S be a minimal zero-sum (resp. zero-sum free) sequence of elements over abelian group G . An element g_0 in S is called splittable if there exist two elements $x, y \in G$ such that $x + y = g_0$ and $Sg_0^{-1}xy$ is a minimal zero-sum (resp. zero-sum free) sequence as well; otherwise, g_0 is called unsplitable. S is called splittable if at least one of the elements of S is splittable; otherwise, it is called unsplitable.

Clearly, if S is a minimal zero-sum sequence and S' is obtained from S by splitting some elements of S , then $|S| \leq |S'|$ and $\text{ind}(S) \leq \text{ind}(S')$.

Introduction and Notation

Long
zero-sum
free and
 n -zero-sum
free
sequences

Li

$\sigma(S)$ –the sum of all terms of S .

$h(S)$ –the maximum multiplicity of a term in S .

$\Sigma(S)$ –the set of all subsums of S , and $\Sigma_k(S)$ – the set of k -term subsums of S , where $k \in \mathbb{N}$.

Definition 1.2

Let S be a minimal zero-sum (resp. zero-sum free) sequence of elements over abelian group G . An element g_0 in S is called splittable if there exist two elements $x, y \in G$ such that $x + y = g_0$ and $Sg_0^{-1}xy$ is a minimal zero-sum (resp. zero-sum free) sequence as well; otherwise, g_0 is called unsplitable. S is called splittable if at least one of the elements of S is splittable; otherwise, it is called unsplitable.

Clearly, if S is a minimal zero-sum sequence and S' is obtained from S by splitting some elements of S , then $|S| \leq |S'|$ and $\text{ind}(S) \leq \text{ind}(S')$.

1. Introduction and Notation

2. On long zero-sum free sequences over \mathbb{Z}_n

3. long n -zero-sum free sequences over \mathbb{Z}_n

4. Subsums of a long zero-sum free sequence.

Bibliography

Introduction and Notation

Long
zero-sum
free and
 n -zero-sum
free
sequences

Li

In this talk we will focus on the following three aspects in zero-sum theory.

1. Introduction
and
Notation

2. On long
zero-sum
free
sequences
over \mathbb{Z}_n

3. long
 n -zero-sum
free
sequences
over \mathbb{Z}_n

4. Subsums
of a long
zero-sum
free
sequence.

Bibliography

Introduction and Notation

Long
zero-sum
free and
 n -zero-sum
free
sequences

Li

In this talk we will focus on the following three aspects in zero-sum theory.

1. long zero-sum free sequences.

1. Introduction
and
Notation

2. On long
zero-sum
free
sequences
over \mathbb{Z}_n

3. long
 n -zero-sum
free
sequences
over \mathbb{Z}_n

4. Subsums
of a long
zero-sum
free
sequence.

Bibliography

Introduction and Notation

Long
zero-sum
free and
 n -zero-sum
free
sequences

Li

In this talk we will focus on the following three aspects in zero-sum theory.

- 1 long zero-sum free sequences.
- 2 n -zero-sum free sequences.

1. Introduction
and
Notation

2. On long
zero-sum
free
sequences
over \mathbb{Z}_n

3. long
 n -zero-sum
free
sequences
over \mathbb{Z}_n

4. Subsums
of a long
zero-sum
free
sequence.

Bibliography

Introduction and Notation

Long
zero-sum
free and
 n -zero-sum
free
sequences

Li

In this talk we will focus on the following three aspects in zero-sum theory.

- 1 long zero-sum free sequences.
- 2 n -zero-sum free sequences.
- 3 Subsums of a long zero-sum free sequence (When may those subsums form an interval?).

1. Introduction
and
Notation

2. On long
zero-sum
free
sequences
over \mathbb{Z}_n

3. long
 n -zero-sum
free
sequences
over \mathbb{Z}_n

4. Subsums
of a long
zero-sum
free
sequence.

Bibliography

Introduction and Notation

Long
zero-sum
free and
 n -zero-sum
free
sequences

Li

In this talk we will focus on the following three aspects in zero-sum theory.

- 1 long zero-sum free sequences.
- 2 n -zero-sum free sequences.
- 3 Subsums of a long zero-sum free sequence (When may those subsums form an interval?).

1. Introduction
and
Notation

2. On long
zero-sum
free
sequences
over \mathbb{Z}_n

3. long
 n -zero-sum
free
sequences
over \mathbb{Z}_n

4. Subsums
of a long
zero-sum
free
sequence.

Bibliography

long zero-sum free sequences

Long
zero-sum
free and
 n -zero-sum
free
sequences

Li

A zero-sum free sequence in \mathbb{Z}_n of length $\geq cn$ with c a constant is referred as to a long zero-sum free sequence. The study of long zero-sum free sequences in \mathbb{Z}_n has attracted considerable attention recently. There are several related results on the structure of zero-sum free sequences.

1. Introduction
and
Notation

2. On long
zero-sum
free
sequences
over \mathbb{Z}_n

3. long
 n -zero-sum
free
sequences
over \mathbb{Z}_n

4. Subsums
of a long
zero-sum
free
sequence.

Bibliography

long zero-sum free sequences

Long
zero-sum
free and
 n -zero-sum
free
sequences

Li

A zero-sum free sequence in \mathbb{Z}_n of length $\geq cn$ with c a constant is referred as to a long zero-sum free sequence. The study of long zero-sum free sequences in \mathbb{Z}_n has attracted considerable attention recently. There are several related results on the structure of zero-sum free sequences.

W. Gao [2] characterized the zero-sum free sequences of lengths roughly greater than $\frac{2n}{3}$.

1. Introduction
and
Notation

2. On long
zero-sum
free
sequences
over \mathbb{Z}_n

3. long
 n -zero-sum
free
sequences
over \mathbb{Z}_n

4. Subsums
of a long
zero-sum
free
sequence.

Bibliography

long zero-sum free sequences

Long
zero-sum
free and
 n -zero-sum
free
sequences

Li

A zero-sum free sequence in \mathbb{Z}_n of length $\geq cn$ with c a constant is referred as to a long zero-sum free sequence. The study of long zero-sum free sequences in \mathbb{Z}_n has attracted considerable attention recently. There are several related results on the structure of zero-sum free sequences.

W. Gao [2] characterized the zero-sum free sequences of lengths roughly greater than $\frac{2n}{3}$. S. Savchev and F. Chen [8], and P. Yuan [10] independently proved that each zero-sum free sequence S in \mathbb{Z}_n with $|S| > \frac{n}{2}$ has index less than 1.

1. Introduction
and
Notation

2. On long
zero-sum
free
sequences
over \mathbb{Z}_n

3. long
 n -zero-sum
free
sequences
over \mathbb{Z}_n

4. Subsums
of a long
zero-sum
free
sequence.

Bibliography

long zero-sum free sequences

Long
zero-sum
free and
 n -zero-sum
free
sequences

Li

A zero-sum free sequence in \mathbb{Z}_n of length $\geq cn$ with c a constant is referred as to a long zero-sum free sequence. The study of long zero-sum free sequences in \mathbb{Z}_n has attracted considerable attention recently. There are several related results on the structure of zero-sum free sequences.

W. Gao [2] characterized the zero-sum free sequences of lengths roughly greater than $\frac{2n}{3}$. S. Savchev and F. Chen [8], and P. Yuan [10] independently proved that each zero-sum free sequence S in \mathbb{Z}_n with $|S| > \frac{n}{2}$ has index less than 1.

We now consider the general structure of the zero-sum free sequences S in \mathbb{Z}_n of length $\frac{n}{3} + 1 < |S| \leq \frac{n}{2}$, and our first main result (Theorem 2.2) shows that by removing at most two elements from S , the index of the remaining sequence is not more than $1 - \frac{4}{n}$.

1. Introduction
and
Notation

2. On long
zero-sum
free
sequences
over \mathbb{Z}_n

3. long
 n -zero-sum
free
sequences
over \mathbb{Z}_n

4. Subsums
of a long
zero-sum
free
sequence.

Bibliography

long zero-sum free sequences

Long
zero-sum
free and
 n -zero-sum
free
sequences

Li

1. Introduction
and
Notation

2. On long
zero-sum
free
sequences
over \mathbb{Z}_n

3. long
 n -zero-sum
free
sequences
over \mathbb{Z}_n

4. Subsums
of a long
zero-sum
free
sequence.

Bibliography

The following lemma is essential to approach our main results.

Lemma 2.1

[11, 12] Let $n \geq 13$ be an odd integer, and S be an unsplittable minimal zero-sum sequence of length $|S| > \lfloor \frac{n}{3} \rfloor + 2$ over \mathbb{Z}_n . If $\text{ind}(S) \geq 2$, then

$$S \sim g^\alpha \left(\frac{n+s}{2} g \right)^{2t} \left(\left(\frac{n-s}{2} + 1 \right) g \right),$$

where g is a generator of \mathbb{Z}_n , $t \geq 1$, s is odd with $s \geq 3$ and $2\alpha + 2ts + 2 - s = n$. Moreover, $\text{ind}(S) = 2$.

long zero-sum free sequences

The First Main Result.

Theorem 2.2

[3] Let $n \geq 50$ be an odd integer, and S be a zero-sum free sequence of length $\lfloor \frac{n}{3} \rfloor + 1 < |S| \leq \lfloor \frac{n}{2} \rfloor$ over \mathbb{Z}_n . Then, by removing at most two elements from S , the remaining sequence is equivalent to a sequence whose index is not more than $1 - 4/n$.

Proof of Theorem 2.2.

Long zero-sum free and n -zero-sum free sequences

Li

1. Introduction and Notation

2. On long zero-sum free sequences over \mathbb{Z}_n

3. long n -zero-sum free sequences over \mathbb{Z}_n

4. Subsums of a long zero-sum free sequence.

Bibliography

long zero-sum free sequences

The First Main Result.

Theorem 2.2

[3] Let $n \geq 50$ be an odd integer, and S be a zero-sum free sequence of length $\lfloor \frac{n}{3} \rfloor + 1 < |S| \leq \lfloor \frac{n}{2} \rfloor$ over \mathbb{Z}_n . Then, by removing at most two elements from S , the remaining sequence is equivalent to a sequence whose index is not more than $1 - 4/n$.

Proof of Theorem 2.2.

Let S be a zero-sum free sequence over \mathbb{Z}_n with $l = |S| > \lfloor \frac{n}{3} \rfloor + 1$. Then $S_1 = S(-\sigma(S))$ is a minimal zero-sum sequence of length $|S_1| > \lfloor \frac{n}{3} \rfloor + 2$. By splitting S_1 if necessary, we eventually obtain an unsplittable minimal zero-sum sequence S' .

Long zero-sum free and n -zero-sum free sequences

Li

1. Introduction and Notation

2. On long zero-sum free sequences over \mathbb{Z}_n

3. long n -zero-sum free sequences over \mathbb{Z}_n

4. Subsums of a long zero-sum free sequence.

Bibliography

long zero-sum free sequences

The First Main Result.

Theorem 2.2

[3] Let $n \geq 50$ be an odd integer, and S be a zero-sum free sequence of length $\lfloor \frac{n}{3} \rfloor + 1 < |S| \leq \lfloor \frac{n}{2} \rfloor$ over \mathbb{Z}_n . Then, by removing at most two elements from S , the remaining sequence is equivalent to a sequence whose index is not more than $1 - 4/n$.

Proof of Theorem 2.2.

Let S be a zero-sum free sequence over \mathbb{Z}_n with $l = |S| > \lfloor \frac{n}{3} \rfloor + 1$. Then $S_1 = S(-\sigma(S))$ is a minimal zero-sum sequence of length $|S_1| > \lfloor \frac{n}{3} \rfloor + 2$. By splitting S_1 if necessary, we eventually obtain an unsplittable minimal zero-sum sequence S' . Then, we have $|S_1| \leq |S'|$ and

$$\text{ind}(S) < \text{ind}(S_1) \leq \text{ind}(S'). \quad (1)$$

Long zero-sum free and n -zero-sum free sequences

Li

1. Introduction and Notation

2. On long zero-sum free sequences over \mathbb{Z}_n

3. long n -zero-sum free sequences over \mathbb{Z}_n

4. Subsums of a long zero-sum free sequence.

Bibliography

long zero-sum free sequences

The First Main Result.

Theorem 2.2

[3] Let $n \geq 50$ be an odd integer, and S be a zero-sum free sequence of length $\lfloor \frac{n}{3} \rfloor + 1 < |S| \leq \lfloor \frac{n}{2} \rfloor$ over \mathbb{Z}_n . Then, by removing at most two elements from S , the remaining sequence is equivalent to a sequence whose index is not more than $1 - 4/n$.

Proof of Theorem 2.2.

Let S be a zero-sum free sequence over \mathbb{Z}_n with $l = |S| > \lfloor \frac{n}{3} \rfloor + 1$. Then $S_1 = S(-\sigma(S))$ is a minimal zero-sum sequence of length $|S_1| > \lfloor \frac{n}{3} \rfloor + 2$. By splitting S_1 if necessary, we eventually obtain an unsplittable minimal zero-sum sequence S' . Then, we have $|S_1| \leq |S'|$ and

$$\text{ind}(S) < \text{ind}(S_1) \leq \text{ind}(S'). \quad (1)$$

Long zero-sum free and n -zero-sum free sequences

Li

1. Introduction and Notation

2. On long zero-sum free sequences over \mathbb{Z}_n

3. long n -zero-sum free sequences over \mathbb{Z}_n

4. Subsums of a long zero-sum free sequence.

Bibliography

long zero-sum free sequences

Long zero-sum free and n -zero-sum free sequences

Li

1. Introduction and Notation

2. On long zero-sum free sequences over \mathbb{Z}_n

3. long n -zero-sum free sequences over \mathbb{Z}_n

4. Subsums of a long zero-sum free sequence.

Bibliography

- ① **Case 1:** If $\text{ind}(S') \leq 1$, then $\text{ind}(S) < 1$ by (1). Thus $S \sim (n_1 h) \cdot \dots \cdot (n_l h)$ with $\text{ord}(h) = n$ and $\text{ind}(S) = \|S\|_h = \frac{n_1 + \dots + n_l}{n}$. If $n_i + n_j \geq 4$ for some $1 \leq i \neq j \leq l$, then $\frac{\sum_{k=1}^l n_k - n_i - n_j}{n} \leq \frac{\sum_{k=1}^l n_k - 4}{n} < 1 - \frac{4}{n}$. If $n_i + n_j \leq 3$ for all i, j , then there is at most one i such that $n_i = 2$, and for all other $j \neq i$, $n_j = 1$. So $\sum_{k=1}^l n_k \leq \lfloor \frac{n}{2} \rfloor + 1$ (as $l \leq \lfloor \frac{n}{2} \rfloor$), and thus $\text{ind}(S) \leq \frac{\lfloor \frac{n}{2} \rfloor + 1}{n} < 1 - \frac{4}{n}$.

long zero-sum free sequences

Long zero-sum free and n -zero-sum free sequences

Li

1. Introduction and Notation

2. On long zero-sum free sequences over \mathbb{Z}_n

3. long n -zero-sum free sequences over \mathbb{Z}_n

4. Subsums of a long zero-sum free sequence.

Bibliography

- ① **Case 1:** If $\text{ind}(S') \leq 1$, then $\text{ind}(S) < 1$ by (1). Thus $S \sim (n_1 h) \cdot \dots \cdot (n_l h)$ with $\text{ord}(h) = n$ and $\text{ind}(S) = \|S\|_h = \frac{n_1 + \dots + n_l}{n}$. If $n_i + n_j \geq 4$ for some $1 \leq i \neq j \leq l$, then $\frac{\sum_{k=1}^l n_k - n_i - n_j}{n} \leq \frac{\sum_{k=1}^l n_k - 4}{n} < 1 - \frac{4}{n}$. If $n_i + n_j \leq 3$ for all i, j , then there is at most one i such that $n_i = 2$, and for all other $j \neq i$, $n_j = 1$. So $\sum_{k=1}^l n_k \leq \lfloor \frac{n}{2} \rfloor + 1$ (as $l \leq \lfloor \frac{n}{2} \rfloor$), and thus $\text{ind}(S) \leq \frac{\lfloor \frac{n}{2} \rfloor + 1}{n} < 1 - \frac{4}{n}$.
- ② **Case 2:.** If $\text{ind}(S') > 1$, then by Lemma 2.1, $S' \sim g^\alpha (\frac{n+s}{2} g)^{2t} ((\frac{n-s}{2} + 1)g) \sim (2g)^\alpha (sg)^{2t} ((n-s+2)g) = T$.

long zero-sum free sequences

Long zero-sum free and n -zero-sum free sequences

Li

1. Introduction and Notation

2. On long zero-sum free sequences over \mathbb{Z}_n

3. long n -zero-sum free sequences over \mathbb{Z}_n

4. Subsums of a long zero-sum free sequence.

Bibliography

- ① **Case 1:** If $\text{ind}(S') \leq 1$, then $\text{ind}(S) < 1$ by (1). Thus $S \sim (n_1 h) \cdot \dots \cdot (n_l h)$ with $\text{ord}(h) = n$ and $\text{ind}(S) = \|S\|_h = \frac{n_1 + \dots + n_l}{n}$. If $n_i + n_j \geq 4$ for some $1 \leq i \neq j \leq l$, then $\frac{\sum_{k=1}^l n_k - n_i - n_j}{n} \leq \frac{\sum_{k=1}^l n_k - 4}{n} < 1 - \frac{4}{n}$. If $n_i + n_j \leq 3$ for all i, j , then there is at most one i such that $n_i = 2$, and for all other $j \neq i$, $n_j = 1$. So $\sum_{k=1}^l n_k \leq \lfloor \frac{n}{2} \rfloor + 1$ (as $l \leq \lfloor \frac{n}{2} \rfloor$), and thus $\text{ind}(S) \leq \frac{\lfloor \frac{n}{2} \rfloor + 1}{n} < 1 - \frac{4}{n}$.
- ② **Case 2:.** If $\text{ind}(S') > 1$, then by Lemma 2.1, $S' \sim g^\alpha (\frac{n+s}{2} g)^{2t} ((\frac{n-s}{2} + 1)g) \sim (2g)^\alpha (sg)^{2t} ((n-s+2)g) = T$. By removing 3 elements from T , including sg and $(n-s+2)g$, the remaining sequence has index not more than $1 - 4/n$. If we remove two elements from S such that, including the elements sg and $(n-s+2)g$, at least three terms of T are left out, then the remaining sequence of S has index not more than $1 - 4/n$.

long zero-sum free sequences

Long zero-sum free and n -zero-sum free sequences

Li

1. Introduction and Notation

2. On long zero-sum free sequences over \mathbb{Z}_n

3. long n -zero-sum free sequences over \mathbb{Z}_n

4. Subsums of a long zero-sum free sequence.

Bibliography

- ① **Case 1:** If $\text{ind}(S') \leq 1$, then $\text{ind}(S) < 1$ by (1). Thus $S \sim (n_1 h) \cdot \dots \cdot (n_l h)$ with $\text{ord}(h) = n$ and $\text{ind}(S) = \|S\|_h = \frac{n_1 + \dots + n_l}{n}$. If $n_i + n_j \geq 4$ for some $1 \leq i \neq j \leq l$, then $\frac{\sum_{k=1}^l n_k - n_i - n_j}{n} \leq \frac{\sum_{k=1}^l n_k - 4}{n} < 1 - \frac{4}{n}$. If $n_i + n_j \leq 3$ for all i, j , then there is at most one i such that $n_i = 2$, and for all other $j \neq i$, $n_j = 1$. So $\sum_{k=1}^l n_k \leq \lfloor \frac{n}{2} \rfloor + 1$ (as $l \leq \lfloor \frac{n}{2} \rfloor$), and thus $\text{ind}(S) \leq \frac{\lfloor \frac{n}{2} \rfloor + 1}{n} < 1 - \frac{4}{n}$.
- ② **Case 2:.** If $\text{ind}(S') > 1$, then by Lemma 2.1, $S' \sim g^\alpha (\frac{n+s}{2} g)^{2t} ((\frac{n-s}{2} + 1)g) \sim (2g)^\alpha (sg)^{2t} ((n-s+2)g) = T$. By removing 3 elements from T , including sg and $(n-s+2)g$, the remaining sequence has index not more than $1 - 4/n$. If we remove two elements from S such that, including the elements sg and $(n-s+2)g$, at least three terms of T are left out, then the remaining sequence of S has index not more than $1 - 4/n$.

Lower bound on $h(S)$

Let S be a zero-sum free sequence over \mathbb{Z}_n . An extensively used result of Bovey et al [1] states that if $|S| = l > n/2$, then $h(S) \geq 2l - n + 1$. S. Savchev and F. Chen [8], and P. Yuan [10] gave a more precisely lower bound on $h(S)$ for zero-sum free sequences of length $l > n/2$ independently. We now give a lower bound for $h(S)$ when the length of S is between $\lfloor \frac{n}{3} \rfloor + 1$ and $\lfloor \frac{n}{2} \rfloor$.

Theorem 2.3

[3] Let $n \geq 13$ be an odd integer and l be an integer satisfying $\lfloor \frac{n}{3} \rfloor + 1 < l \leq \lfloor \frac{n}{2} \rfloor$. Let S be a zero-sum free sequence of length l over \mathbb{Z}_n , and i be an positive integer such that $1 \leq i \leq 3$ and $4l - 4 - n \equiv i \pmod{3}$. If $\text{ind}(S) > 1$, then

- (a) $h(S) \geq 3l - 2 - n$, if $\frac{2n+5-i}{5} \leq l \leq \lfloor \frac{n}{2} \rfloor$;
(b) $h(S) \geq \lfloor \frac{4l-4-n}{3} \rfloor + 1$, if $\lfloor \frac{n}{3} \rfloor + 1 < l < \frac{2n+5-i}{5}$.

Moreover, these estimates are best possible.

Long zero-sum free and n -zero-sum free sequences

Li

1. Introduction and Notation

2. On long zero-sum free sequences over \mathbb{Z}_n

3. long n -zero-sum free sequences over \mathbb{Z}_n

4. Subsums of a long zero-sum free sequence.

Bibliography

Lower bound on $h(S)$

Proof. By the assumption and Theorem 2.2, we may assume

$$S = (2g)^u (sg)^v W g_1 g_2, \quad \|S(g_1 g_2)^{-1}\|_g \leq 1 - \frac{4}{n}, \quad \text{ind}(S) = \|S\|_g,$$

where $s \geq 3$, $W|S$, $1 \geq |W| \geq 0$ and $2g, sg \notin W$. We claim that $g \notin S$.

Long
zero-sum
free and
 n -zero-sum
free
sequences

Li

1. Introduction
and
Notation

2. On long
zero-sum
free
sequences
over \mathbb{Z}_n

3. long
 n -zero-sum
free
sequences
over \mathbb{Z}_n

4. Subsums
of a long
zero-sum
free
sequence.

Bibliography

Lower bound on $h(S)$

Proof. By the assumption and Theorem 2.2, we may assume

$$S = (2g)^u (sg)^v W g_1 g_2, \quad \|S(g_1 g_2)^{-1}\|_g \leq 1 - \frac{4}{n}, \quad \text{ind}(S) = \|S\|_g,$$

where $s \geq 3$, $W|S$, $1 \geq |W| \geq 0$ and $2g, sg \notin W$. We claim that $g \notin S$. Then

$$n - 4 \geq n \|S(g_1 g_2)^{-1}\|_g \geq 2u + 3(l - 2 - u) = 3l - 6 - u. \quad (2)$$

$$n - 4 \geq n \|S(g_1 g_2)^{-1}\|_g \geq 2u + 3v + 4(l - 2 - u - v) = 4l - 8 - 2u - v \quad (3)$$

The above inequalities yield that $u \geq 3l - 2 - n$ and $2u + v \geq 4l - 4 - n$, respectively. Since $h(S) \geq \max\{u, v\}$, we have $h(S) \geq \max\{3l - 2 - n, \lceil \frac{4l - 4 - n}{3} \rceil\}$.

Long zero-sum free and n -zero-sum free sequences

Li

1. Introduction and Notation

2. On long zero-sum free sequences over Z_n

3. long n -zero-sum free sequences over Z_n

4. Subsums of a long zero-sum free sequence.

Bibliography

Lower bound on $h(S)$

Proof. By the assumption and Theorem 2.2, we may assume

$$S = (2g)^u (sg)^v W g_1 g_2, \quad \|S(g_1 g_2)^{-1}\|_g \leq 1 - \frac{4}{n}, \quad \text{ind}(S) = \|S\|_g,$$

where $s \geq 3$, $W|S$, $1 \geq |W| \geq 0$ and $2g, sg \notin W$. We claim that $g \notin S$. Then

$$n - 4 \geq n \|S(g_1 g_2)^{-1}\|_g \geq 2u + 3(l - 2 - u) = 3l - 6 - u. \quad (2)$$

$$n - 4 \geq n \|S(g_1 g_2)^{-1}\|_g \geq 2u + 3v + 4(l - 2 - u - v) = 4l - 8 - 2u - v \quad (3)$$

The above inequalities yield that $u \geq 3l - 2 - n$ and $2u + v \geq 4l - 4 - n$, respectively. Since $h(S) \geq \max\{u, v\}$, we have $h(S) \geq \max\{3l - 2 - n, \lceil \frac{4l - 4 - n}{3} \rceil\}$.

We next show that the above estimates are best possible by finding some extremal sequences S with $h(S) = 3l - 2 - n$ and $h(S) = \lceil \frac{4l - 4 - n}{3} \rceil$, respectively. This can be done by the case by case argument on $i = 1, 2, 3$.

Long zero-sum free and n -zero-sum free sequences

Li

1. Introduction and Notation

2. On long zero-sum free sequences over Z_n

3. long n -zero-sum free sequences over Z_n

4. Subsums of a long zero-sum free sequence.

Bibliography

Lower bound on $h(S)$

Proof. By the assumption and Theorem 2.2, we may assume

$$S = (2g)^u (sg)^v W g_1 g_2, \quad \|S(g_1 g_2)^{-1}\|_g \leq 1 - \frac{4}{n}, \quad \text{ind}(S) = \|S\|_g,$$

where $s \geq 3$, $W|S$, $1 \geq |W| \geq 0$ and $2g, sg \notin W$. We claim that $g \notin S$. Then

$$n - 4 \geq n \|S(g_1 g_2)^{-1}\|_g \geq 2u + 3(l - 2 - u) = 3l - 6 - u. \quad (2)$$

$$n - 4 \geq n \|S(g_1 g_2)^{-1}\|_g \geq 2u + 3v + 4(l - 2 - u - v) = 4l - 8 - 2u - v \quad (3)$$

The above inequalities yield that $u \geq 3l - 2 - n$ and $2u + v \geq 4l - 4 - n$, respectively. Since $h(S) \geq \max\{u, v\}$, we have $h(S) \geq \max\{3l - 2 - n, \lceil \frac{4l - 4 - n}{3} \rceil\}$.

We next show that the above estimates are best possible by finding some extremal sequences S with $h(S) = 3l - 2 - n$ and $h(S) = \lceil \frac{4l - 4 - n}{3} \rceil$, respectively. This can be done by the case by case argument on $i = 1, 2, 3$.

Long zero-sum free and n -zero-sum free sequences

Li

1. Introduction and Notation

2. On long zero-sum free sequences over Z_n

3. long n -zero-sum free sequences over Z_n

4. Subsums of a long zero-sum free sequence.

Bibliography

long n -zero-sum free sequences

The n -zero-sum free sequences in \mathbb{Z}_n play an important role in the investigation of the structure of zero-sum free sequences. Savchev and Chen [9] characterized n -zero-sum free sequences of length $|S| \geq \frac{3n}{2} - 1$. Our main theorem below characterizes those sequences S with $|S| = n + l$ where $l \geq n/p + p - 2$ and p is the smallest prime divisor of $|G| = n$.

Long
zero-sum
free and
 n -zero-sum
free
sequences

Li

1. Introduction
and
Notation

2. On long
zero-sum
free
sequences
over \mathbb{Z}_n

3. long
 n -zero-sum
free
sequences
over \mathbb{Z}_n

4. Subsums
of a long
zero-sum
free
sequence.

Bibliography

long n -zero-sum free sequences

Long zero-sum free and n -zero-sum free sequences

Li

1. Introduction and Notation

2. On long zero-sum free sequences over \mathbb{Z}_n

3. long n -zero-sum free sequences over \mathbb{Z}_n

4. Subsums of a long zero-sum free sequence.

Bibliography

The n -zero-sum free sequences in \mathbb{Z}_n play an important role in the investigation of the structure of zero-sum free sequences. Savchev and Chen [9] characterized n -zero-sum free sequences of length $|S| \geq \frac{3n}{2} - 1$. Our main theorem below characterizes those sequences S with $|S| = n + l$ where $l \geq n/p + p - 2$ and p is the smallest prime divisor of $|G| = n$.

Theorem 3.1

Let G be a finite abelian group of order n , and $l \geq n/p + p - 2$ be an integer where p is the smallest prime divisor of n . Let S be a n -zero-sum free sequence over G of length $|S| = n + l$. Then, there is an element $g \in G$ such that

$$-g + S = 0^h T S'$$

with $h = h(S) \geq l + 1$, T is a zero-sum sequence of length $|T| \leq |G| - h - 1$, and S' is zero-sum free of length $|S'| \geq l + 1$.

long n -zero-sum free sequences

The n -zero-sum free sequences in \mathbb{Z}_n play an important role in the investigation of the structure of zero-sum free sequences. Savchev and Chen [9] characterized n -zero-sum free sequences of length $|S| \geq \frac{3n}{2} - 1$. Our main theorem below characterizes those sequences S with $|S| = n + l$ where $l \geq n/p + p - 2$ and p is the smallest prime divisor of $|G| = n$.

Theorem 3.1

Let G be a finite abelian group of order n , and $l \geq n/p + p - 2$ be an integer where p is the smallest prime divisor of n . Let S be a n -zero-sum free sequence over G of length $|S| = n + l$. Then, there is an element $g \in G$ such that

$$-g + S = 0^h T S'$$

with $h = h(S) \geq l + 1$, T is a zero-sum sequence of length $|T| \leq |G| - h - 1$, and S' is zero-sum free of length $|S'| \geq l + 1$.

Long zero-sum free and n -zero-sum free sequences

Li

1. Introduction and Notation

2. On long zero-sum free sequences over \mathbb{Z}_n

3. long n -zero-sum free sequences over \mathbb{Z}_n

4. Subsums of a long zero-sum free sequence.

Bibliography

long n -zero-sum free sequences

Long
zero-sum
free and
 n -zero-sum
free
sequences

Li

1. Introduction
and
Notation

2. On long
zero-sum
free
sequences
over \mathbb{Z}_n

3. long
 n -zero-sum
free
sequences
over \mathbb{Z}_n

4. Subsums
of a long
zero-sum
free
sequence.

Bibliography

Corollary 3.2

Let p be the smallest prime dividing n , and S be an n -zero-sum free sequence over \mathbb{Z}_n of length $|S| = n + \ell$, where $\ell \geq n/p + p - 2$ is an integer. Then, there exists $g \in \mathbb{Z}_n$ such that

$$-g + S = 0^h T S'$$

with $h \geq \ell + 1$, T is a zero-sum sequence of length $|T| \leq n - h - 1$, and S' is zero-sum free with $|S'| \geq \ell + 1$.

Subsums

Long
zero-sum
free and
 n -zero-sum
free
sequences

Li

1. Introduction
and
Notation

2. On long
zero-sum
free
sequences
over \mathbb{Z}_n

3. long
 n -zero-sum
free
sequences
over \mathbb{Z}_n

4. Subsums
of a long
zero-sum
free
sequence.

Bibliography

Let S be a long zero-sum free sequence in \mathbb{Z}_n . We investigate when subsums of S (with $|S| \geq (n + 2)/3$) form an interval:
Question: Let S be a long zero-sum free sequence over \mathbb{Z}_n . Does there exist a sequence T , such that $T \sim S$ and subsums of T form an interval ? (such T is referred as to a smooth sequence).

Subsums

Long
zero-sum
free and
 n -zero-sum
free
sequences

Li

1. Introduction
and
Notation

2. On long
zero-sum
free
sequences
over \mathbb{Z}_n

3. long
 n -zero-sum
free
sequences
over \mathbb{Z}_n

4. Subsums
of a long
zero-sum
free
sequence.

Bibliography

Let S be a long zero-sum free sequence in \mathbb{Z}_n . We investigate when subsums of S (with $|S| \geq (n + 2)/3$) form an interval:

Question: Let S be a long zero-sum free sequence over \mathbb{Z}_n . Does there exist a sequence T , such that $T \sim S$ and subsums of T form an interval ? (such T is referred as to a smooth sequence).

In [8], S. Savchev and F. Chen showed that if S is a zero-sum free sequence of length $|S| > \frac{n}{2}$ over \mathbb{Z}_n , then $ind(S) < 1$. Thus there exists a sequence T , such that $T \sim S$ and $\sigma(T) < n$ (as positive integers)(clearly, $1 \in T$ as $|T| = |S| > n/2$). Moreover, $\sum(T) = [1, \sigma(T)]$ is an interval (i.e., T is smooth). Next we will show that if S is a zero-sum free sequence over \mathbb{Z}_n of length $|S| \geq \frac{n+2}{3}$ such that $1 \in S$ and $\sigma(S) < n$ (as positive integers), then $\sum(S)$ is an interval except for one special case.

Subsums

Long zero-sum free and n -zero-sum free sequences

Li

1. Introduction and Notation

2. On long zero-sum free sequences over \mathbb{Z}_n

3. long n -zero-sum free sequences over \mathbb{Z}_n

4. Subsums of a long zero-sum free sequence.

Bibliography

Let S be a long zero-sum free sequence in \mathbb{Z}_n . We investigate when subsums of S (with $|S| \geq (n + 2)/3$) form an interval:

Question: Let S be a long zero-sum free sequence over \mathbb{Z}_n . Does there exist a sequence T , such that $T \sim S$ and subsums of T form an interval ? (such T is referred as to a smooth sequence).

In [8], S. Savchev and F. Chen showed that if S is a zero-sum free sequence of length $|S| > \frac{n}{2}$ over \mathbb{Z}_n , then $ind(S) < 1$. Thus there exists a sequence T , such that $T \sim S$ and $\sigma(T) < n$ (as positive integers)(clearly, $1 \in T$ as $|T| = |S| > n/2$). Moreover, $\sum(T) = [1, \sigma(T)]$ is an interval (i.e., T is smooth). Next we will show that if S is a zero-sum free sequence over \mathbb{Z}_n of length $|S| \geq \frac{n+2}{3}$ such that $1 \in S$ and $\sigma(S) < n$ (as positive integers), then $\sum(S)$ is an interval except for one special case.

Subsums

Long
zero-sum
free and
 n -zero-sum
free
sequences

Li

1. Introduction
and
Notation

2. On long
zero-sum
free
sequences
over \mathbb{Z}_n

3. long
 n -zero-sum
free
sequences
over \mathbb{Z}_n

4. Subsums
of a long
zero-sum
free
sequence.

Bibliography

Theorem 4.1

[3] Let S be a sequence with positive integer terms of length $|S| = t \geq \frac{n+2}{3}$, and $\sigma(S) < n$. If $1 \in S$, then $\Sigma(S)$ is an interval except for the case when $S = S_0 n_1$, where $\Sigma(S_0)$ is an interval and $n_1 > \sigma(S_0) + 1$.

Subsums

Long
zero-sum
free and
 n -zero-sum
free
sequences

Li

1. Introduction
and
Notation

2. On long
zero-sum
free
sequences
over \mathbb{Z}_n

3. long
 n -zero-sum
free
sequences
over \mathbb{Z}_n

4. Subsums
of a long
zero-sum
free
sequence.

Bibliography

Theorem 4.1

[3] Let S be a sequence with positive integer terms of length $|S| = t \geq \frac{n+2}{3}$, and $\sigma(S) < n$. If $1 \in S$, then $\Sigma(S)$ is an interval except for the case when $S = S_0n_l$, where $\Sigma(S_0)$ is an interval and $n_l > \sigma(S_0) + 1$.

We remark that the sequence S in the above theorem can be regarded as a zero-sum free sequence S over \mathbb{Z}_n of length $|S| \geq \frac{n+2}{3}$ such that $1 \in S$ and $\sigma(S) < n$ (as positive integers). Such S has $\text{ind}(S) < 1$ and it is almost smooth.

Subsums

Long
zero-sum
free and
 n -zero-sum
free
sequences

Li

1. Introduction
and
Notation

2. On long
zero-sum
free
sequences
over \mathbb{Z}_n

3. long
 n -zero-sum
free
sequences
over \mathbb{Z}_n

4. Subsums
of a long
zero-sum
free
sequence.

Bibliography

Theorem 4.1

[3] Let S be a sequence with positive integer terms of length $|S| = t \geq \frac{n+2}{3}$, and $\sigma(S) < n$. If $1 \in S$, then $\Sigma(S)$ is an interval except for the case when $S = S_0n_l$, where $\Sigma(S_0)$ is an interval and $n_l > \sigma(S_0) + 1$.

We remark that the sequence S in the above theorem can be regarded as a zero-sum free sequence S over \mathbb{Z}_n of length $|S| \geq \frac{n+2}{3}$ such that $1 \in S$ and $\sigma(S) < n$ (as positive integers). Such S has $\text{ind}(S) < 1$ and it is almost smooth.

Long
zero-sum
free and
n-zero-sum
free
sequences

Li

1. Introduction
and
Notation

2. On long
zero-sum
free
sequences
over \mathbb{Z}_n

3. long
 n -zero-sum
free
sequences
over \mathbb{Z}_n

4. Subsums
of a long
zero-sum
free
sequence.

Bibliography

Thank you!

AUTHOR: Yuanlin Li

ADDRESS: Department of Mathematics and Statistics
Brock University
Canada

EMAIL: yli@brocku.ca

Bibliography I

Long
zero-sum
free and
 n -zero-sum
free
sequences

Li

1. Introduction
and
Notation

2. On long
zero-sum
free
sequences
over \mathbb{Z}_n

3. long
 n -zero-sum
free
sequences
over \mathbb{Z}_n

4. Subsums
of a long
zero-sum
free
sequence.

Bibliography



J.D. Bovey, P. Erdős, I. Niven, Conditions for a zero sum modulo n , *Canad. Math. Bull.* 18 (1) (1975) 27 C29.



W. Gao, Zero sums in finite cyclic groups, *Integers*(2000), A12, 7 pp.



W. Gao, Y. Li, P. Yuan and J. Zhuang, On the structure of long zero-sum free sequences and n -zero-sum free sequences over finite cyclic groups, *Arch. Math* 105(2015), 361-370.



A. Geroldinger, Additive group theory and non-unique factorizations, in *Combinatorial number Theory and Additive Group Theory*, eds. A. Geroldinger and I. Ruzsa, *Advanced Courses in Mathematics CRM Barcelona* (Birkhäuser, Basel, 2009), pp. 1-86.

Bibliography II

Long
zero-sum
free and
 n -zero-sum
free
sequences

Li






1. Introduction
and
Notation

2. On long
zero-sum
free
sequences
over \mathbb{Z}_n

3. long
 n -zero-sum
free
sequences
over \mathbb{Z}_n

4. Subsums
of a long
zero-sum
free
sequence.

Bibliography

-  Y. Li and J. Peng, Minimal zero-sum sequences of length four over finite cyclic groups II, *Int. J. Number Theory*, 9(2013), 845-866.
-  Y. Li, C. Plyley, P. Yuan and X. Zeng, Minimal zero sum sequences of length four over finite cyclic groups, *J. Number Theory*. 130(2010), 2033-2048.
-  J. Peng, Y. Li, Minimal zero-sum sequences of length five over finite cyclic groups, *Ars Combinatoria*, 112(2013), 373-384.
-  S. Savchev and F. Chen, Long zero-free sequences in finite cyclic groups, *Discrete Math.* 307(2007), 2671-2679.
-  S. Savchev and F. Chen, Long n -zero-free sequences in finite cyclic groups, *Discrete Math.* 308(2008), 1-8.

Bibliography III

Long
zero-sum
free and
 n -zero-sum
free
sequences

Li




1. Introduction
and
Notation

2. On long
zero-sum
free
sequences
over \mathbb{Z}_n

3. long
 n -zero-sum
free
sequences
over \mathbb{Z}_n

4. Subsums
of a long
zero-sum
free
sequence.

Bibliography

-  P. Yuan, On the index of minimal zero-sum sequences over finite cyclic groups, *J. Combin. Theory Ser. A* 114(2007), no. 8, 1545-1551.
-  P. Yuan and Y. Li, Long unsplittable zero-sum sequences over a finite cyclic group, *Int. J. Number Theory*, Accepted, 2015.
-  X. Zeng, P. Yuan and Y. Li, On the structure of long unsplittable minimal zero-sum sequences, Submitted, 2015.